

Participants with Drug-Related Adverse Events  
(Incidence  $\geq 6\%$  in One or More Treatment Groups)  
Weeks 0 to 12  
All Participants as Treated

| Low Dose                                                                 |               | Placebo   |               | Total     |               |
|--------------------------------------------------------------------------|---------------|-----------|---------------|-----------|---------------|
| n                                                                        | (%)           | n         | (%)           | n         | (%)           |
| Participants in population <sup>84</sup>                                 |               | 86        |               | 170       |               |
| with one or more drug-related adverse events <sup>73</sup>               | (86.9)        | 44        | (51.2)        | 117       | (68.8)        |
| with no drug-related adverse events <sup>11</sup>                        | (13.1)        | 42        | (48.8)        | 53        | (31.2)        |
|                                                                          |               |           |               |           |               |
| <b>Cardiac disorders<sup>7</sup></b>                                     | <b>(8.3)</b>  | <b>6</b>  | <b>(7.0)</b>  | <b>13</b> | <b>(7.6)</b>  |
| <b>Gastrointestinal disorders<sup>8</sup></b>                            | <b>(9.5)</b>  | <b>4</b>  | <b>(4.7)</b>  | <b>12</b> | <b>(7.1)</b>  |
| <b>General disorders and administration site conditions<sup>43</sup></b> | <b>(51.2)</b> | <b>18</b> | <b>(20.9)</b> | <b>61</b> | <b>(35.9)</b> |
| Application site pruritus <sup>22</sup>                                  | (26.2)        | 6         | (7.0)         | 28        | (16.5)        |
| Application site dermatitis <sup>9</sup>                                 | (10.7)        | 5         | (5.8)         | 14        | (8.2)         |
| Application site erythema <sup>12</sup>                                  | (14.3)        | 3         | (3.5)         | 15        | (8.8)         |
| Application site irritation <sup>9</sup>                                 | (10.7)        | 3         | (3.5)         | 12        | (7.1)         |
| <b>Nervous system disorders<sup>12</sup></b>                             | <b>(14.3)</b> | <b>5</b>  | <b>(5.8)</b>  | <b>17</b> | <b>(10.0)</b> |
| Dizziness <sup>6</sup>                                                   | (7.1)         | 2         | (2.3)         | 8         | (4.7)         |
| <b>Psychiatric disorders<sup>9</sup></b>                                 | <b>(10.7)</b> | <b>2</b>  | <b>(2.3)</b>  | <b>11</b> | <b>(6.5)</b>  |
| <b>Skin and subcutaneous tissue disorders<sup>39</sup></b>               | <b>(46.4)</b> | <b>17</b> | <b>(19.8)</b> | <b>56</b> | <b>(32.9)</b> |
| Erythema <sup>13</sup>                                                   | (15.5)        | 9         | (10.5)        | 22        | (12.9)        |
| Pruritus <sup>21</sup>                                                   | (25.0)        | 7         | (8.1)         | 28        | (16.5)        |
| Rash <sup>11</sup>                                                       | (13.1)        | 3         | (3.5)         | 14        | (8.2)         |
| Skin irritation <sup>6</sup>                                             | (7.1)         | 2         | (2.3)         | 8         | (4.7)         |
| Blister <sup>5</sup>                                                     | (6.0)         | 0         | (0.0)         | 5         | (2.9)         |

Every participant is counted a single time for each applicable row and column.

A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.

Adverse event terms are from MedDRA Version 24.0.Source: [CDISCpilot: adam-adsl; adae]