

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose		Placebo		Total	
n	(%)	n	(%)	n	(%)
Participants in population ⁸⁴		86		170	
with one or more drug-related adverse events ⁷³	(86.9)	44	(51.2)	117	(68.8)
with no drug-related adverse events ¹¹	(13.1)	42	(48.8)	53	(31.2)
Cardiac disorders⁷	(8.3)	6	(7.0)	13	(7.6)
Myocardial infarction ¹	(1.2)	2	(2.3)	3	(1.8)
Sinus bradycardia ²	(2.4)	2	(2.3)	4	(2.4)
Atrial fibrillation ⁰	(0.0)	1	(1.2)	1	(0.6)
Atrioventricular block first degree ⁰	(0.0)	1	(1.2)	1	(0.6)
Atrioventricular block second degree ⁰	(0.0)	1	(1.2)	1	(0.6)
Bradycardia ⁰	(0.0)	1	(1.2)	1	(0.6)
Cardiac failure congestive ⁰	(0.0)	1	(1.2)	1	(0.6)
Sinus arrhythmia ⁰	(0.0)	1	(1.2)	1	(0.6)
Atrial flutter ¹	(1.2)	0	(0.0)	1	(0.6)
Bundle branch block right ¹	(1.2)	0	(0.0)	1	(0.6)
Palpitations ¹	(1.2)	0	(0.0)	1	(0.6)
Supraventricular extrasystoles ¹	(1.2)	0	(0.0)	1	(0.6)
Ventricular extrasystoles ¹	(1.2)	0	(0.0)	1	(0.6)
Wolff-parkinson-white syndrome ¹	(1.2)	0	(0.0)	1	(0.6)
Congenital, familial and genetic disorders¹	(1.2)	0	(0.0)	1	(0.6)
Ventricular septal defect ¹	(1.2)	0	(0.0)	1	(0.6)

Ear and labyrinth disorders ²	(2.4)	0	(0.0)	2	(1.2)
Tinnitus ¹	(1.2)	0	(0.0)	1	(0.6)
Vertigo ¹	(1.2)	0	(0.0)	1	(0.6)
Eye disorders ¹	(1.2)	0	(0.0)	1	(0.6)
Vision blurred ¹	(1.2)	0	(0.0)	1	(0.6)
Gastrointestinal disorders ⁸	(9.5)	4	(4.7)	12	(7.1)
Diarrhoea ³	(3.6)	3	(3.5)	6	(3.5)
Dyspepsia ¹	(1.2)	1	(1.2)	2	(1.2)
Gastrooesophageal reflux disease ⁰	(0.0)	1	(1.2)	1	(0.6)

Participants with Drug-Related Adverse Events
 (Incidence > 0 % in One or More Treatment Groups)
 Weeks 0 to 12
 All Participants as Treated

Low Dose		Placebo		Total	
n	(%)	n	(%)	n	(%)
Abdominal pain ¹	(1.2)	0	(0.0)	1	(0.6)
Nausea ³	(3.6)	0	(0.0)	3	(1.8)
Vomiting ²	(2.4)	0	(0.0)	2	(1.2)
General disorders and administration site conditions ⁴³	(51.2)	18	(20.9)	61	(35.9)
Application site pruritus ²²	(26.2)	6	(7.0)	28	(16.5)
Application site dermatitis ⁹	(10.7)	5	(5.8)	14	(8.2)
Application site erythema ¹²	(14.3)	3	(3.5)	15	(8.8)
Application site irritation ⁹	(10.7)	3	(3.5)	12	(7.1)
Application site induration ⁰	(0.0)	1	(1.2)	1	(0.6)
Application site reaction ⁰	(0.0)	1	(1.2)	1	(0.6)
Application site vesicles ⁴	(4.8)	1	(1.2)	5	(2.9)
Asthenia ⁰	(0.0)	1	(1.2)	1	(0.6)
Chills ¹	(1.2)	1	(1.2)	2	(1.2)
Fatigue ²	(2.4)	1	(1.2)	3	(1.8)
Application site bleeding ¹	(1.2)	0	(0.0)	1	(0.6)
Application site desquamation ¹	(1.2)	0	(0.0)	1	(0.6)

Application site discolouration1	(1.2)	0	(0.0)	1	(0.6)
Application site swelling1	(1.2)	0	(0.0)	1	(0.6)
Application site urticaria2	(2.4)	0	(0.0)	2	(1.2)
Application site warmth1	(1.2)	0	(0.0)	1	(0.6)
Malaise1	(1.2)	0	(0.0)	1	(0.6)
Oedema2	(2.4)	0	(0.0)	2	(1.2)
Pain1	(1.2)	0	(0.0)	1	(0.6)
Injury, poisoning and procedural complications2	(2.4)	0	(0.0)	2	(1.2)
Fall1	(1.2)	0	(0.0)	1	(0.6)
Skin laceration1	(1.2)	0	(0.0)	1	(0.6)
Wound1	(1.2)	0	(0.0)	1	(0.6)
Investigations2	(2.4)	4	(4.7)	6	(3.5)
Blood creatine phosphokinase increased0	(0.0)	1	(1.2)	1	(0.6)
Electrocardiogram st segment depression1	(1.2)	1	(1.2)	2	(1.2)

Participants with Drug-Related Adverse Events
 (Incidence > 0 % in One or More Treatment Groups)
 Weeks 0 to 12
 All Participants as Treated

Low Dose		Placebo		Total	
n	(%)	n	(%)	n	(%)
Heart rate increased0	(0.0)	1	(1.2)	1	(0.6)
Heart rate irregular0	(0.0)	1	(1.2)	1	(0.6)
Body temperature increased1	(1.2)	0	(0.0)	1	(0.6)
Metabolism and nutrition disorders0	(0.0)	3	(3.5)	3	(1.8)
Decreased appetite0	(0.0)	1	(1.2)	1	(0.6)
Food craving0	(0.0)	1	(1.2)	1	(0.6)
Increased appetite0	(0.0)	1	(1.2)	1	(0.6)

Musculoskeletal and connective tissue disorders 0	(0.0)	1	(1.2)	1	(0.6)
Shoulder pain0	(0.0)	1	(1.2)	1	(0.6)
Nervous system disorders 12	(14.3)	5	(5.8)	17	(10.0)
Dizziness6	(7.1)	2	(2.3)	8	(4.7)
Headache1	(1.2)	2	(2.3)	3	(1.8)
Somnolence0	(0.0)	1	(1.2)	1	(0.6)
Balance disorder1	(1.2)	0	(0.0)	1	(0.6)
Complex partial seizures1	(1.2)	0	(0.0)	1	(0.6)
Coordination abnormal1	(1.2)	0	(0.0)	1	(0.6)
Lethargy1	(1.2)	0	(0.0)	1	(0.6)
Paraesthesia oral1	(1.2)	0	(0.0)	1	(0.6)
Stupor1	(1.2)	0	(0.0)	1	(0.6)
Syncope4	(4.8)	0	(0.0)	4	(2.4)
Transient ischaemic attack1	(1.2)	0	(0.0)	1	(0.6)
Psychiatric disorders 9	(10.7)	2	(2.3)	11	(6.5)
Confusional state2	(2.4)	1	(1.2)	3	(1.8)
Irritability1	(1.2)	1	(1.2)	2	(1.2)
Agitation2	(2.4)	0	(0.0)	2	(1.2)
Anxiety3	(3.6)	0	(0.0)	3	(1.8)
Depressed mood1	(1.2)	0	(0.0)	1	(0.6)
Restlessness1	(1.2)	0	(0.0)	1	(0.6)
Renal and urinary disorders 1	(1.2)	0	(0.0)	1	(0.6)
Enuresis1	(1.2)	0	(0.0)	1	(0.6)
Reproductive system and breast disorders 0	(0.0)	1	(1.2)	1	(0.6)

Participants with Drug-Related Adverse Events
(Incidence > 0 % in One or More Treatment Groups)
Weeks 0 to 12
All Participants as Treated

Low Dose	Placebo	Total
----------	---------	-------

n	(%)	n	(%)	n	(%)
Pelvic pain0	(0.0)	1	(1.2)	1	(0.6)
Respiratory, thoracic and mediastinal disorders0	(0.0)	2	(2.3)	2	(1.2)
Dyspnoea0	(0.0)	1	(1.2)	1	(0.6)
Emphysema0	(0.0)	1	(1.2)	1	(0.6)
Skin and subcutaneous tissue disorders39	(46.4)	17	(19.8)	56	(32.9)
Erythema13	(15.5)	9	(10.5)	22	(12.9)
Pruritus21	(25.0)	7	(8.1)	28	(16.5)
Rash11	(13.1)	3	(3.5)	14	(8.2)
Skin irritation6	(7.1)	2	(2.3)	8	(4.7)
Cold sweat0	(0.0)	1	(1.2)	1	(0.6)
Hyperhidrosis4	(4.8)	1	(1.2)	5	(2.9)
Skin ulcer0	(0.0)	1	(1.2)	1	(0.6)
Blister5	(6.0)	0	(0.0)	5	(2.9)
Dermatitis contact1	(1.2)	0	(0.0)	1	(0.6)
Pruritus generalised1	(1.2)	0	(0.0)	1	(0.6)
Rash erythematous2	(2.4)	0	(0.0)	2	(1.2)
Rash pruritic1	(1.2)	0	(0.0)	1	(0.6)
Skin exfoliation1	(1.2)	0	(0.0)	1	(0.6)
Urticaria1	(1.2)	0	(0.0)	1	(0.6)
Vascular disorders2	(2.4)	1	(1.2)	3	(1.8)
Hypotension1	(1.2)	1	(1.2)	2	(1.2)
Orthostatic hypotension0	(0.0)	1	(1.2)	1	(0.6)
Hypertension1	(1.2)	0	(0.0)	1	(0.6)

Every participant is counted a single time for each applicable row and column.

A system organ class or specific adverse event appears on this report only if its incidence in one or more of the columns meets the incidence criterion in the report title, after rounding.

Adverse event terms are from MedDRA Version 24.0.Source: [CDISCPilot: adam-ads!; adae]