

## BFGS ALGORITHM

MERIEL RYU

Suppose that  $n$  is a positive integer and  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  is of class  $C^2$ . We investigate the notion of BFGS algorithm in order to find the minimum of  $f$ . This will be done with basics from vector calculus such as Taylor's theorem and Lagrange multiplier method. Throughout the article,  $x \in \mathbb{R}^n$  is regarded as a column vector.

### 1. NEWTON'S METHOD

We recall the classical Newton's method. Suppose we want to find a root of a differentiable function  $g : \mathbb{R} \rightarrow \mathbb{R}$ . First, we choose an initial value  $x_0 \in \mathbb{R}$ . The tangent line through  $(x_0, g(x_0))$  is given by

$$y = g'(x_0)(x - x_0) + g(x_0) \quad (1.1)$$

The root of (1.1) is taken as  $x_1$ , that is,

$$x_1 = x_0 - \frac{g(x_0)}{g'(x_0)}.$$

In general, the sequence  $\{x_k\}_{k=1}^{\infty}$  is determined by the following recurrence relation:

$$x_{k+1} = x_k - \frac{g(x_k)}{g'(x_k)} \quad (k = 0, 1, 2, \dots).$$

If the initial value  $x_0$  is close enough to a root of  $g(x)$  and  $g'(x_0) \neq 0$ , then  $x_k$  converges to the root of  $g(x)$  as  $k$  goes to the infinity.

Newton's method can be used to find a minimum or maximum of a  $C^2$  function. In this case, we apply Newton's method to the derivative because the derivative is zero at a minimum or maximum. Suppose that we want to find a minimum of a  $C^2$  function  $g : \mathbb{R} \rightarrow \mathbb{R}$ . The second order Taylor approximation of  $g$  is

$$g(x+d) \approx g(x) + g'(x)d + \frac{1}{2}g''(x)d^2.$$

This approximation is valid only if  $d$  is small enough. If we put

$$q(d) = g(x) + g'(x)d + \frac{1}{2}g''(x)d^2,$$

then  $q(d)$  is minimized when

$$q'(d) = 0.$$

Since

$$q'(d) = g'(x) + g''(x)d,$$

it follows that  $q(d)$  attains a minimum at

$$d = -\frac{g'(x)}{g''(x)}.$$

We note that  $q'(d) = 0$  does not imply that  $q$  is minimized at  $d$ . It is well known that  $g''(x)$  is positive at a local minimum. Since  $q(d)$  is a quadratic function, positiveness of  $g''(x)$  guarantees the existence of a minimum by the convexity of  $q(d)$ . We choose an initial value  $x_0 \in \mathbb{R}$  and let

$$x_{k+1} = x_k - \frac{g'(x_k)}{g''(x_k)} \quad (k = 0, 1, 2, \dots).$$

The same argument can be applied when we want to find a maximum of  $g(x)$  because a maximum of  $g(x)$  is actually a minimum of  $-g(x)$ .

Now, we use Newton's method in order to find a minimum point of a  $C^2$  function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ . Suppose  $x, d \in \mathbb{R}^n$ . The second order Taylor approximation of  $f$  is

$$f(x + d) \approx f(x) + \sum_{i=1}^n d_i \frac{\partial f}{\partial x_i}(x) + \frac{1}{2} \sum_{i,j=1}^n d_i d_j \frac{\partial^2 f}{\partial x_i \partial x_j}(x). \quad (1.2)$$

This approximation is valid only if  $d$  is sufficiently close to  $0 \in \mathbb{R}^n$ . If we denote the gradient of  $f$  and the Hessian of  $f$  by  $\nabla f$  and  $Hf$ , respectively, then (1.2) can be written as follows:

$$f(x + d) \approx f(x) + d^T \nabla f(x) + \frac{1}{2} d^T Hf(x) d.$$

If we set

$$q(d) = f(x) + d^T \nabla f(x) + \frac{1}{2} d^T Hf(x) d, \quad (1.3)$$

then  $q(d)$  is minimized when

$$\nabla q(d) = 0.$$

Since  $\nabla q(d) = \nabla f(x) + Hf(x)d$ , it follows that  $q(d)$  attains a minimum at

$$d = -Hf^{-1}(x) \nabla f(x).$$

In the same manner as the one-dimensional case,  $\nabla q(d) = 0$  does not guarantee that  $d$  is a minimum point of  $q$ . It is well known that the Hessian  $Hf(x)$  is positive-definite at a local minimum  $x$ . Hence, if  $Hf(x)$  is positive-definite, then the point  $d \in \mathbb{R}^n$  satisfying  $\nabla q(d) = 0$  is the minimum of  $q$  because  $q$  is a convex quadratic function. We choose an initial value  $x_0 \in \mathbb{R}^n$  and let

$$\delta_k = -Hf^{-1}(x_k) \nabla f(x_k),$$

$$x_{k+1} = x_k + \delta_k \quad (k = 0, 1, 2, \dots).$$

If the Hessian  $Hf(x_k)$  at  $x_k$  is positive definite, then so is  $Hf^{-1}(x_k)$  and  $\delta_k$  becomes the downhill direction from  $x_k$ . We discuss it in the rest of the section. For  $x \in \mathbb{R}^n$ , a direction  $d \in \mathbb{R}^n$  is called a *downhill direction* if there is a positive real number  $\alpha'$  such that

$$0 < \alpha < \alpha' \quad \Rightarrow \quad f(x + \alpha d) < f(x).$$

**Lemma 1.1.** *Suppose that  $x, d \in \mathbb{R}^n$ . If  $d$  satisfies*

$$d^T \nabla f(x) < 0,$$

*then  $d$  is a downhill direction. In particular,  $-\nabla f(x)$  is a downhill direction.*

*Proof.* By Taylor's theorem,

$$f(x + \alpha d) = f(x) + \alpha d^T \nabla f(x) + R_2(x, \alpha d),$$

where

$$\frac{|R_2(x, \alpha d)|}{|\alpha| \|d\|} \rightarrow 0 \quad \text{as} \quad \alpha \rightarrow 0.$$

Hence, there is a positive real number  $\alpha'$  such that

$$0 < \alpha < \alpha' \quad \Rightarrow \quad \frac{|R_2(x, \alpha d)|}{|\alpha| \|d\|} < |d^T \nabla f(x)|.$$

This implies that

$$0 < \alpha < \alpha' \quad \Rightarrow \quad f(x + \alpha d) - f(x) = \alpha d^T \nabla f(x) + R_2(x, \alpha d) < 0$$

because  $d^T \nabla f(x) < 0$ .  $\square$

## 2. QUASI-NEWTON METHOD

If  $n$  is large, then Hessian is impractical to compute. Quasi-Newton method uses approximate Hessian  $H_k$ . Instead of (1.3), we apply the same argument to

$$q(d) = f(x) + d^T \nabla f(x) + \frac{1}{2} d^T H_k d.$$

If  $x_0 \in \mathbb{R}^n$  is an initial value, then we have

$$\delta_k = -H_k^{-1} \nabla f(x_k)$$

and

$$x_{k+1} = x_k + \delta_k \quad (k = 0, 1, 2, \dots).$$

There are several methods to update  $H_k$  and in the next section we will investigate BFGS updates, which is one of quasi-Newton methods. In this section, we discuss four desired properties of  $H_k$  in quasi-Newton methods. We start with the secant condition. Since the approximation of  $f(x_{k+1})$  is

$$f(x_k + \delta_k) \approx q(\delta_k) = f(x_k) + \delta_k^T \nabla f(x_k) + \frac{1}{2} \delta_k^T H_k \delta_k$$

and  $\nabla q(d) = \nabla f(x) + H_k d$ , it follows that

$$\begin{aligned} \nabla f(x_{k+1}) - \nabla f(x_k) &\approx \nabla q(\delta_k) - \nabla q(0) \\ &= \nabla f(x_k) + H_{k+1} \delta_k - \nabla f(x_k) \\ &= H_{k+1} \delta_k. \end{aligned}$$

For each  $k = 0, 1, 2, \dots$ , we let

$$\gamma_k = \nabla f(x_{k+1}) - \nabla f(x_k).$$

We say that  $H_k$  satisfies *secant condition* if

$$\gamma_k = H_{k+1} \delta_k. \quad (2.1)$$

The first and the second desired properties of  $H_k$  are as follows:

- (1) For a convex quadratic  $C^2$  function  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ ,  $H_k$  converges to the Hessian  $Hf$  of  $f$  as  $k$  goes to the infinity.
- (2)  $H_k$  satisfies the secant condition.

Since  $f$  is a  $C^2$  function, its Hessian is well-defined and symmetric at  $x \in \mathbb{R}^n$ . It is known that if  $x^*$  is a local minimum point, then the Hessian  $Hf(x^*)$  is

positive-definite at  $x^*$ . The following is the third desired property of  $H_k$ :

(3)  $H_k$  is nonsingular, symmetric and positive definite.

We note that the third property guarantees that  $\delta_k = -H_k^{-1}\nabla f(x_k)$  is the downhill direction from  $x_k$ .

The last property is as follows:

(4) We minimize the change in  $H_k$ . This is done by minimizing  $\|H_{k+1} - H_k\|$  in some norm or alternatively minimizing  $\|H_{k+1}^{-1} - H_k^{-1}\|$ .

### 3. BFGS UPDATES

We start the section with the Frobenius norm. We need some notation. If  $n$  is a positive integer, we denote the set of  $n \times n$  real matrix by  $M(n)$ . The Frobenius norm  $\|\cdot\|_F$  is defined by

$$\|A\|_F = \sqrt{\text{tr}(A^T A)} = \sqrt{\sum_{i,j=1}^n |A(i,j)|^2} \quad (A \in M(n)).$$

When  $W \in M(n)$ , a weighted Frobenius norm  $\|\cdot\|_W$  is define by

$$\|A\|_W = \sqrt{\frac{1}{2}\text{tr}(W A W A^T)} \quad (A \in M(n)).$$

If  $W = M^T M$  for some  $M \in M(n)$  and  $W$  is nonsingular, BFGS update is obtained by solving

$$\begin{aligned} & \min_{H \in M(n)} \|H^{-1} - H_k^{-1}\|_W, \\ \text{where} \quad & \delta_k = H^{-1}\gamma_k \quad \text{and} \quad H^T = H. \end{aligned} \quad (3.1)$$

For convenience, we drop the index  $k$  and write  $\delta$  and  $\gamma$  instead of  $\delta_k$  and  $\gamma_k$ . If we set

$$E = H^{-1} - H_k^{-1},$$

then (3.1) can be written as follows:

$$\begin{aligned} & \min_{E \in M(n)} \|E\|_W^2, \\ \text{where} \quad & E\gamma = \delta - H_k^{-1}\gamma \quad \text{and} \quad E^T = E. \end{aligned} \quad (3.2)$$

If we define a function  $L(E, z, N) : M(n) \times \mathbb{R}^n \times M(n) \rightarrow \mathbb{R}$  by

$$L(E, z, N) = \text{tr} \left[ \frac{1}{2} W E W E^T + z^T (E\gamma - (\delta - H_k^{-1}\gamma)) + N(E - E^T) \right],$$

then there are  $\lambda \in \mathbb{R}^n$  and  $\Gamma \in M(n)$  such that

$$\frac{\partial L}{\partial E(s,t)}(E, \lambda, \Gamma) = 0 \quad (3.3)$$

for all  $1 \leq s, t \leq n$ . Lagrange multiplier method tells us that  $E$  satisfying (3.3) is a minimum point with respect to the weighted Frobenius norm  $\|\cdot\|_W$  in the restrictions

$$E\gamma = \delta - H_k^{-1}\gamma \quad \text{and} \quad E^T = E.$$

We first solve (3.3). Since

$$\frac{1}{2}\text{tr}(WEWE^\top) = \frac{1}{2} \sum_{i,j,k,l=1}^n W(i,j)W(k,l)E(j,k)E(i,l),$$

it follows that

$$\frac{\partial}{\partial E(s,t)} \frac{1}{2}\text{tr}(WEWE^\top) = \sum_{i,l=1}^n W(s,i)E(i,l)W(l,t) = WEW(s,t). \quad (3.4)$$

Since

$$\frac{\partial}{\partial E(s,t)} \text{tr}(\lambda^\top(E\gamma - (\delta - H_k^{-1}\gamma))) = \frac{\partial}{\partial E(s,t)} \text{tr}(\lambda^\top E\gamma)$$

and

$$\lambda^\top E\gamma = \sum_{i,j=1}^n \lambda(i)E(i,j)\gamma(j),$$

it follows that

$$\frac{\partial}{\partial E(s,t)} \text{tr}(\lambda^\top(E\gamma - (\delta - H_k^{-1}\gamma))) = \lambda(s)\gamma(t) = \lambda\gamma^\top(s,t). \quad (3.5)$$

Finally, from

$$\text{tr}(\Gamma(E - E^\top)) = \sum_{i,j=1}^n \Gamma(i,j)(E(j,i) - E(i,j)),$$

we obtain

$$\frac{\partial}{\partial E(s,t)} \text{tr}(\Gamma(E - E^\top)) = \Gamma(t,s) - \Gamma(s,t) = (\Gamma^\top - \Gamma)(s,t). \quad (3.6)$$

From (3.4), (3.5) and (3.6), (3.3) becomes

$$WEW + \lambda\gamma^\top + \Gamma^\top - \Gamma = 0$$

and (3.2) can be written as follows:

$$WEW + \lambda\gamma^\top + \Gamma^\top - \Gamma = 0, \quad (3.7)$$

$$E\gamma = \delta - H_k^{-1}\gamma \quad (3.8)$$

and

$$E^\top = E. \quad (3.9)$$

Since  $W = M^\top M$  is nonsingular, (3.7) can be written as follows:

$$E = -W^{-1}(\lambda\gamma^\top + \Gamma^\top - \Gamma)W^{-1}. \quad (3.10)$$

Since  $W = M^\top M$  is symmetric, (3.9) implies

$$\lambda\gamma^\top + \Gamma^\top - \Gamma = \gamma\lambda^\top + \Gamma - \Gamma^\top$$

and this implies

$$\Gamma^\top - \Gamma = \frac{1}{2}(\gamma\lambda^\top - \lambda\gamma^\top).$$

Hence, (3.10) becomes

$$E = -\frac{1}{2}W^{-1}(\gamma\lambda^\top + \lambda\gamma^\top)W^{-1}. \quad (3.11)$$

If we substitute it to (3.8), then we obtain

$$\gamma_k\lambda^\top W^{-1}\gamma + \lambda\gamma^\top W^{-1}\gamma = -2W(\delta - H_k^{-1}\gamma).$$

Since  $\gamma^\top W^{-1} \gamma$  is a scalar, we have

$$\lambda = \frac{-2W\delta + 2WH_k^{-1}\gamma - \gamma\lambda^\top W^{-1}\gamma}{\gamma^\top W^{-1}\gamma}. \quad (3.12)$$

In order to represent  $\lambda$  in (3.12) with respect to  $W$ ,  $H_k$ ,  $\delta$  and  $\gamma$ , we multiply both sides of (3.12) by  $\gamma^\top W^{-1}$ :

$$\gamma^\top W^{-1}\lambda = \frac{-2\gamma^\top \delta + 2\gamma^\top H_k^{-1}\gamma - \gamma^\top W^{-1}\gamma\lambda^\top W^{-1}\gamma}{\gamma^\top W^{-1}\gamma}. \quad (3.13)$$

We transpose (3.13):

$$\begin{aligned} \lambda^\top W^{-1}\gamma &= \frac{-2\delta^\top \gamma + 2\gamma^\top H_k^{-1}\gamma - \gamma^\top W^{-1}\lambda\gamma^\top W^{-1}\gamma}{\gamma^\top W^{-1}\gamma} \\ &= \frac{-2\delta^\top \gamma + 2\gamma^\top H_k^{-1}\gamma}{\gamma^\top W^{-1}\gamma} - \lambda^\top W^{-1}\gamma \end{aligned}$$

and this implies

$$\lambda^\top W^{-1}\gamma = \frac{-\delta^\top \gamma + \gamma^\top H_k^{-1}\gamma}{\gamma^\top W^{-1}\gamma}. \quad (3.14)$$

We substitute (3.14) to (3.12):

$$\lambda = \frac{-2W\delta + 2WH_k^{-1}\gamma}{\gamma^\top W^{-1}\gamma} + \frac{\gamma(\delta^\top - \gamma^\top H_k^{-1})\gamma}{(\gamma^\top W^{-1}\gamma)^2}.$$

Again, we substitute this to (3.11):

$$\begin{aligned} E &= -\frac{1}{2}W^{-1} \left( \frac{-2\gamma\delta^\top W + 2\gamma\gamma^\top H_k^{-1}W}{\gamma^\top W^{-1}\gamma} + \frac{\gamma\gamma^\top(\delta - H_k^{-1}\gamma)\gamma^\top}{(\gamma^\top W^{-1}\gamma)^2} \right. \\ &\quad \left. + \frac{-2W\delta\gamma^\top + 2WH_k^{-1}\gamma\gamma^\top}{\gamma^\top W^{-1}\gamma} + \frac{\gamma(\delta^\top - \gamma^\top H_k^{-1})\gamma\gamma^\top}{(\gamma^\top W^{-1}\gamma)^2} \right) W^{-1}. \end{aligned}$$

Since

$$\gamma(\delta^\top - \gamma^\top H_k^{-1})\gamma = \gamma\gamma^\top(\delta - H_k^{-1}\gamma),$$

we have

$$\begin{aligned} E &= \frac{W^{-1}\gamma(\delta^\top - \gamma^\top H_k^{-1})}{\gamma^\top W^{-1}\gamma} + \frac{(\delta - H_k^{-1}\gamma)\gamma^\top W^{-1}}{\gamma^\top W^{-1}\gamma} \\ &\quad - \frac{W^{-1}\gamma\gamma^\top(\delta - H_k^{-1}\gamma)\gamma^\top W^{-1}}{(\gamma^\top W^{-1}\gamma)^2}. \end{aligned} \quad (3.15)$$

We have not chosen the weight  $W$  yet. Different weights  $W$  lead to different updates. Here, we will put  $W = H_{k+1}$ . In the next proposition, we will see that this choice is possible. A matrix  $A \in M(n)$  is said to have a *Cholesky factorization* if there is a lower triangular matrix  $L$  whose diagonal entries are all positive such that  $X = LL^\top$ .

**Proposition 3.1.** *A matrix  $A \in M(n)$  has a Cholesky factorization  $A = LL^\top$  if and only if  $A$  is symmetric and positive definite. Moreover, if the Cholesky factorization exists, then it is unique.*

*Proof.* We only prove that every symmetric positive definite matrix has a Cholesky factorization. The proof will be done by mathematical induction on  $n$ . When  $n = 1$  and  $A = [a]$ , obviously  $L = [\sqrt{a}]$ . We assume the proposition is true for  $n$  and prove a symmetric positive matrix  $A \in M(n+1)$  has a Cholesky factorization. Let

$$A = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n+1} \\ a_{21} & a_{22} & \cdots & a_{2n+1} \\ \vdots & \vdots & & \vdots \\ a_{n+11} & a_{n+12} & \cdots & a_{n+1n+1} \end{bmatrix} \quad \text{and} \quad A' = \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{n1} & a_{n2} & \cdots & a_{nn} \end{bmatrix}.$$

By assumption  $A'$  has a Cholesky factorization  $A' = LL^\top$ . When

$$L = \begin{bmatrix} l_{11} & 0 & \cdots & 0 \\ l_{21} & l_{22} & \cdots & 0 \\ \vdots & \vdots & & \vdots \\ l_{n1} & l_{n2} & \cdots & l_{nn} \end{bmatrix},$$

we can find  $m_1, m_2, \dots, m_{n+1}$  such that

$$M = \begin{bmatrix} l_{11} & 0 & \cdots & 0 & 0 \\ l_{21} & l_{22} & \cdots & 0 & 0 \\ \vdots & \vdots & & \vdots & \vdots \\ l_{n1} & l_{n2} & \cdots & l_{nn} & 0 \\ m_1 & m_2 & \cdots & m_n & m_{n+1} \end{bmatrix}$$

has the requiring property.  $\square$

If  $W = H_{k+1}$ , then  $W^{-1} = H_{k+1}^{-1} = E + H_k^{-1}$ . By (3.15) and the secant condition (2.1), we have

$$\begin{aligned} H_{k+1}^{-1} &= H_k^{-1} + \frac{\delta(\delta^\top - \gamma^\top H_k^{-1})}{\gamma^\top \delta} + \frac{(\delta - H_k^{-1}\gamma)\delta^\top}{\gamma^\top \delta} - \frac{\delta\gamma^\top(\delta - H_k^{-1}\gamma)\delta^\top}{(\gamma^\top \delta)^2} \\ &= H_k^{-1} - \frac{\delta\gamma^\top H_k^{-1}}{\gamma^\top \delta} - \frac{H_k^{-1}\gamma\delta^\top}{\gamma^\top \delta} + \frac{\delta\delta^\top}{\gamma^\top \delta} + \frac{(\gamma^\top H_k^{-1}\gamma)\delta\delta^\top}{(\gamma^\top \delta)^2} \\ &= H_k^{-1} - \frac{\delta\gamma^\top H_k^{-1}}{\gamma^\top \delta} - \frac{H_k^{-1}\gamma\delta^\top}{\gamma^\top \delta} + \frac{1}{\gamma^\top \delta} \left( 1 + \frac{\gamma^\top H_k^{-1}\gamma}{\gamma^\top \delta} \right) \delta\delta^\top. \end{aligned} \quad (3.16)$$

In order to drive the update of  $H_k$  from that of  $H_k^{-1}$ , we introduce the Sherman-Morrison formula:

**Lemma 3.2.** *Suppose that  $A \in M(n)$  is nonsingular and that  $u, v \in \mathbb{R}^n$ . Then  $A + uv^\top$  is nonsingular if and only if  $1 + v^\top A^{-1}u$  is nonzero. If  $A + uv^\top$  is nonsingular, its inverse is*

$$(A + uv^\top)^{-1} = A^{-1} - \frac{A^{-1}uv^\top A^{-1}}{1 + v^\top A^{-1}u}.$$

*Proof.* Direct computations show that

$$(A + uv^\top) \left( A^{-1} - \frac{A^{-1}uv^\top A^{-1}}{1 + v^\top A^{-1}u} \right) = I_n$$

and that

$$\left( A^{-1} - \frac{A^{-1}uv^\top A^{-1}}{1 + v^\top A^{-1}u} \right) (A + uv^\top) = I_n.$$

By the uniqueness of the inverse, the result follows.  $\square$

If  $A + uv^\top + vu^\top$  is nonsingular, then we can obtain its inverse by applying the Sherman-Morrison formula to

$$(A + uv^\top)^{-1} \quad \text{and} \quad ((A + uv^\top) + vu^\top)^{-1}.$$

Direct computations show that

$$\begin{aligned} (A + uv^\top + vu^\top)^{-1} &= A^{-1} + \frac{(u^\top A^{-1}u)A^{-1}vv^\top A^{-1} + (v^\top A^{-1}v)A^{-1}uu^\top}{(1 + v^\top A^{-1}u)(1 + u^\top A^{-1}v) - (u^\top A^{-1}u)(v^\top A^{-1}v)} \\ &\quad + \frac{-(1 + v^\top A^{-1}u)A^{-1}vu^\top A^{-1} - (1 + u^\top A^{-1}v)A^{-1}uv^\top A^{-1}}{(1 + v^\top A^{-1}u)(1 + u^\top A^{-1}v) - (u^\top A^{-1}u)(v^\top A^{-1}v)}. \end{aligned} \quad (3.17)$$

By (3.17), we obtain

$$\begin{aligned} &\left( H_k^{-1} - \frac{H_k^{-1}\gamma\delta^\top}{\gamma^\top\delta} - \frac{\delta\gamma^\top H_k^{-1}}{\gamma^\top\delta} \right)^{-1} \\ &= H_k - \frac{(\gamma^\top H_k^{-1}\gamma)H_k\delta\delta^\top H_k}{(\gamma^\top H_k^{-1}\gamma)(\delta^\top H_k\delta)} - \frac{(\delta^\top H_k\delta)\gamma\gamma^\top}{(\gamma^\top H_k^{-1}\gamma)(\delta^\top H_k\delta)}. \end{aligned}$$

Applying the Sherman-Morrison formula once again, we have

$$\begin{aligned} H_{k+1} &= \left( H_k^{-1} - \frac{\delta\gamma^\top H_k^{-1}}{\gamma^\top\delta} - \frac{H_k^{-1}\gamma\delta^\top}{\gamma^\top\delta} + \frac{1}{\gamma^\top\delta} \left( 1 + \frac{\gamma^\top H_k^{-1}\gamma}{\gamma^\top\delta} \right) \delta\delta^\top \right)^{-1} \\ &= H_k + \frac{\gamma\gamma^\top}{\gamma^\top\delta} - \frac{H_k\delta\delta^\top H_k}{\delta^\top H_k\delta}. \end{aligned} \quad (3.18)$$

If  $u, v \in \mathbb{R}^n$ , then the rank of  $u \times v^\top \in M(n)$  is 1. The update of the form

$$H_{k+1} = H_k + uv^\top$$

is called a rank 1 update. Our BFGS update is expressed as a sum of rank 1 updates.

If  $H_k$  is symmetric, then so is  $H_{k+1}$ . Now, we want to see that if  $H_k$  is positive definite, then so is  $H_{k+1}$ . For any  $x \in \mathbb{R}^n \setminus \{0\}$ , we have

$$\begin{aligned} x^\top H_{k+1}x &= x^\top H_kx + x^\top \frac{\gamma\gamma^\top}{\gamma^\top\delta}x - x^\top \frac{H_k\delta\delta^\top H_k}{\delta^\top H_k\delta}x \\ &= \frac{(x^\top H_kx)(\delta^\top H_k\delta) - (x^\top H_k\delta)(\delta^\top H_kx)}{\delta^\top H_k\delta} + \frac{(x^\top\gamma)^2}{\gamma^\top\delta}. \end{aligned}$$

If we define a map  $\langle \cdot, \cdot \rangle_{H_k} : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  by

$$\langle x, y \rangle_{H_k} = x^\top H_k y,$$

then  $\langle \cdot, \cdot \rangle_{H_k}$  is an inner product since  $H_k$  is positive definite and symmetric. By Cauchy-Schwarz inequality, we have

$$(x^\top H_kx)(\delta^\top H_k\delta) - (x^\top H_k\delta)(\delta^\top H_kx) = \langle x, x \rangle_{H_k} \langle \delta, \delta \rangle_{H_k} - \langle x, \delta \rangle_{H_k}^2 \geq 0.$$

Thus, positiveness of  $H_k$  implies

$$\frac{(x^\top H_kx)(\delta^\top H_k\delta) - (x^\top H_k\delta)(\delta^\top H_kx)}{\delta^\top H_k\delta} > 0.$$



It remains to show that

$$\frac{(x^\top \gamma)^2}{\gamma^\top \delta} > 0.$$

In (1.2), if the length of the vector  $d$  is large, then the Taylor approximation is not accurate. Exact line search and inexact line search are common methods to fix this problem. Exact line search is a method to choose  $\alpha$  which minimizes  $f(x_k + \alpha\delta)$  but it is not considered cost effective. Inexact line search is a method to try different  $\alpha$  until the step length  $\alpha$  is acceptable (not to be too long or not to be too short) so that  $x_k$  would converge to some value well. Wolfe conditions provide upper and lower bound on the admissible step length values when we perform inexact line search. We say  $\alpha$  satisfies the Wolfe conditions if the following two inequalities hold:

$$(1) \ f(x_k + \alpha\delta) \leq f(x_k) + c_1 \alpha \delta^\top \nabla f(x_k) \quad (0 < c_1 < 1).$$

$$(2) \ -\delta^\top \nabla f(x_k + \alpha\delta) \leq -c_2 \delta^\top \nabla f(x_k) \quad (c_1 < c_2 < 1).$$

The first inequality (1) ensures that the step length decreases  $f$  sufficiently and the second inequality (2) ensures that the slope has been reduced sufficiently.

If we perform exact line search, then we have

$$\frac{d}{d\alpha} f(x_k + \alpha\delta) = 0.$$

Since

$$\frac{d}{d\alpha} f(x_k + \alpha\delta) = \delta^\top \nabla f(x_{k+1}),$$

we have

$$\gamma^\top \delta = \delta^\top \gamma = \delta^\top (\nabla f(x_{k+1}) - \nabla f(x_k)) = -\delta^\top \nabla f(x_k) > 0$$

by Lemma 1.1.

If we perform inexact line search, then we have

$$\gamma^\top \delta = \delta^\top \gamma = \delta^\top (\nabla f(x_{k+1}) - \nabla f(x_k)) = -\delta^\top \nabla f(x_k) > 0$$

by Lemma 1.1 and the second Wolfe condition.

## REFERENCES

- [1] Quasi-Newton optimization: Origin of the BFGS update Steven G. Johnson, notes for 18.335 at MIT April 25, 2019
- [2] Continuous optimisation: Quasi-Newton methods, Honour School of Mathematics, Oxford University Hilary Term 2005, Dr. Raphael Hauser
- [3] Wolfe Conditions, Wikipedia