

Introduction to dual methods





#### Primal problem

$$\begin{array}{ll} f_0(x) \to \min_{x \in \mathbb{R}^n} \\ \text{s.t.} & f_i(x) \leq 0, \ i=1,\ldots,m \\ & h_i(x) = 0, \ i=1,\ldots,p \end{array}$$

#### Dual problem

$$\begin{split} g(\lambda,\nu) &= \min_{x \in \mathcal{D}} L(x,\lambda,\nu) = \\ \min_{x \in \mathcal{D}} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right) &\to \max_{\lambda \in \mathbb{R}^m, \nu \in \mathbb{R}^p} \\ &\text{s.t. } \lambda \succeq 0 \end{split}$$

 Shadow Prices. In economics and resource allocation problems, dual variables can be interpreted as shadow prices, providing economic insights into resource utilization and constraints.

େ♥େଉ

#### Primal problem

$$\begin{aligned} f_0(x) &\to \min_{x \in \mathbb{R}^n} \\ \text{s.t.} \quad f_i(x) \leq 0, \ i=1,\dots,m \\ h_i(x) = 0, \ i=1,\dots,p \end{aligned}$$

#### Dual problem

$$\begin{split} g(\lambda,\nu) &= \min_{x \in \mathcal{D}} L(x,\lambda,\nu) = \\ \min_{x \in \mathcal{D}} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right) &\to \max_{\lambda \in \mathbb{R}^m, \nu \in \mathbb{R}^p} \\ &\text{s.t. } \lambda \succeq 0 \end{split}$$

- Shadow Prices. In economics and resource allocation problems, dual variables can be interpreted as shadow prices, providing economic insights into resource utilization and constraints.
- Market Equilibrium. Dual problems often represent market equilibrium conditions, making them essential for economic modeling and analysis.



#### Primal problem

$$\begin{aligned} f_0(x) &\to \min_{x \in \mathbb{R}^n} \\ \text{s.t.} & f_i(x) \leq 0, \ i=1,\dots,m \\ & h_i(x) = 0, \ i=1,\dots,p \end{aligned}$$

#### Dual problem

$$\begin{split} g(\lambda,\nu) &= \min_{x \in \mathcal{D}} L(x,\lambda,\nu) = \\ \min_{x \in \mathcal{D}} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right) &\to \max_{\lambda \in \mathbb{R}^m, \nu \in \mathbb{R}^p} \\ &\text{s.t. } \lambda \succeq 0 \end{split}$$

- Shadow Prices. In economics and resource allocation problems, dual variables can be interpreted as shadow prices, providing economic insights into resource utilization and constraints.
- Market Equilibrium. Dual problems often represent market equilibrium conditions, making them essential for economic modeling and analysis.
- **Dual Problems Provide Bounds.** Dual problems often offer bounds on the optimal value of the primal problem. This can be useful for assessing the quality of approximate solutions.

**♥ ೧ Ø** 

#### Primal problem

$$\begin{aligned} f_0(x) &\to \min_{x \in \mathbb{R}^n} \\ \text{s.t.} & f_i(x) \leq 0, \ i=1,\ldots,m \\ & h_i(x) = 0, \ i=1,\ldots,p \end{aligned}$$

#### Dual problem

$$\begin{split} g(\lambda,\nu) &= \min_{x \in \mathcal{D}} L(x,\lambda,\nu) = \\ \min_{x \in \mathcal{D}} \left( f_0(x) + \sum_{i=1}^m \lambda_i f_i(x) + \sum_{i=1}^p \nu_i h_i(x) \right) &\to \max_{\lambda \in \mathbb{R}^m, \nu \in \mathbb{R}^p} \\ &\text{s.t. } \lambda \succeq 0 \end{split}$$

- Shadow Prices. In economics and resource allocation problems, dual variables can be interpreted as shadow prices, providing economic insights into resource utilization and constraints.
- Market Equilibrium. Dual problems often represent market equilibrium conditions, making them essential for economic modeling and analysis.
- **Dual Problems Provide Bounds.** Dual problems often offer bounds on the optimal value of the primal problem. This can be useful for assessing the quality of approximate solutions.
- **Duality Gap.** The difference between the primal and dual solutions (duality gap) provides valuable information about the solution's optimality.




**Conjugate functions** 





## **Conjugate functions**




Recall that given  $f:\mathbb{R}^n\to\mathbb{R},$  the function defined by

$$f^*(y) = \max_x \left[ y^T x - f(x) \right]$$


is called its conjugate.

### **Geometrical intution**



 $\underset{x,y,z}{\mapsto} \min$  Conjugate functions

### **Geometrical intution**



# Conjugate function properties

Recall that given  $f: \mathbb{R}^n \to \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x \left[ y^T x - f(x) \right]$$

is called its conjugate.

Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x \left[ f(x) - y^T x \right]$$



# Conjugate function properties

Recall that given  $f: \mathbb{R}^n \to \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x \left[ y^T x - f(x) \right]$$

is called its conjugate.

Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x \left[ f(x) - y^T x \right]$$

• If f is closed and convex, then  $f^{**} = f$ . Also,

$$x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x) \Leftrightarrow x \in \arg\min_{z} \left[ f(z) - y^T z \right]$$



# Conjugate function properties

Recall that given  $f: \mathbb{R}^n \to \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x \left[ y^T x - f(x) \right]$$

is called its conjugate.

Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x \left[ f(x) - y^T x \right]$$

• If f is closed and convex, then  $f^{**} = f$ . Also,

$$x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x) \Leftrightarrow x \in \arg\min_{z} \left[ f(z) - y^T z \right]$$

• If f is strictly convex, then

$$\nabla f^*(y) = \arg\min_{z} \left[ f(z) - y^T z \right]$$

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that f is convex and closed.

• **Proof of**  $\Leftarrow$ : Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^Tz - f(z)$  over z. But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{ and } \quad \partial f^*(y) = \operatorname{cl}(\operatorname{conv}(\bigcup_{z \in M_+} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that f is convex and closed.

• Proof of  $\Leftarrow$ : Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^Tz - f(z)$  over z. But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{ and } \quad \partial f^*(y) = \operatorname{cl}(\operatorname{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

• **Proof of**  $\Rightarrow$ : From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .



We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that f is convex and closed.

• Proof of  $\Leftarrow$ : Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^Tz - f(z)$  over z. But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{ and } \quad \partial f^*(y) = \operatorname{cl}(\operatorname{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

• **Proof of**  $\Rightarrow$ : From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .



We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that f is convex and closed.

• Proof of  $\Leftarrow$ : Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^Tz - f(z)$  over z. But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{ and } \quad \partial f^*(y) = \operatorname{cl}(\operatorname{conv}(\bigcup_{z \in M} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

• **Proof of**  $\Rightarrow$ : From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .

Clearly  $y \in \partial f(x) \Leftrightarrow x \in \arg\min_z \{f(z) - y^T z\}$ 

Lastly, if f is strictly convex, then we know that  $f(z) - y^T z$  has a unique minimizer over z, and this must be  $\nabla f^*(y)$ .

 $f \to \min_{x,y,z}$  Conjugate functions

**Dual ascent** 



Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

#### Consider the problem:

$$\min_{x} \quad f(x) \quad \text{subject to} \quad Ax = b$$

Dual ascent

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_{x} \quad f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_{u} \quad -f^*(-A^Tu) - b^Tu$$

where  $f^{*}$  is the conjugate of f. Defining  $g(u)=-f^{*}(-A^{T}u)-b^{T}u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Dual ascent

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_{x} \quad f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_{u} \quad -f^*(-A^Tu) - b^Tu$$

where  $f^*$  is the conjugate of f. Defining  $g(u) = -f^*(-A^Tu) - b^Tu$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b \quad \text{where} \quad x \in \arg\min_{z} \left[ f(z) + u^T Az \right]$$

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_{x} \quad f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_{u} \quad -f^*(-A^T u) - b^T u$$

where  $f^{*}$  is the conjugate of f. Defining  $g(u)=-f^{*}(-A^{T}u)-b^{T}u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b$$
 where  $x \in \arg\min_{z} \left[ f(z) + u^T Az 
ight]$ 

Dual ascent method for maximizing dual objective: • Step sizes  $\alpha_k$ , k=1,2,3

 $\begin{aligned} \mathbf{i} \\ x_k &\in \arg\min_x \left[ f(x) + (u_{k-1})^T Ax \right] \\ u_k &= u_{k-1} + \alpha_k (Ax_k - b) \end{aligned}$ 

• Step sizes  $\alpha_k$  ,  $k=1,2,3,\ldots$  , are chosen in standard ways.

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_{x} \quad f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

Dual ascent

$$\max_{u} \quad -f^*(-A^Tu) - b^Tu$$

where  $f^{*}$  is the conjugate of f. Defining  $g(u)=-f^{*}(-A^{T}u)-b^{T}u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b$$
 where  $x \in rg \min \left[ f(z) + u^T Az 
ight]$ 


Dual ascent method for maximizing dual objective: • Step sizes  $\alpha_k$ , k=1,2,3,..., are chosen in standard ways.

 $\begin{aligned} \mathbf{i} \\ x_k &\in \arg\min_x \left[ f(x) + (u_{k-1})^T A x \right] \\ u_k &= u_{k-1} + \alpha_k (A x_k - b) \end{aligned}$ 

Proximal gradients and acceleration can be applied as

they would usually.

# ${\bf Slopes} \ {\bf of} \ f \ {\bf and} \ f^*$



Assume that f is a closed and convex function. Then f is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

Assume that f is a closed and convex function. Then f is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of "\Rightarrow"**: Recall, if g is strongly convex with minimizer x, then

$$g(y) \ge g(x) + \frac{\mu}{2} \|y - x\|^2, \quad \text{for all } y$$



Assume that f is a closed and convex function. Then f is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of "\Rightarrow"**: Recall, if g is strongly convex with minimizer x, then

$$g(y) \ge g(x) + \frac{\mu}{2} ||y - x||^2$$
, for all y

Hence, defining  $x_u = \nabla f^*(u)$  and  $x_v = \nabla f^*(v)$ ,

$$f(x_v) - u^T x_v \geq f(x_u) - u^T x_u + \frac{\mu}{2} \|x_u - x_v\|^2$$

$$f(x_u) - v^T x_u \ge f(x_v) - v^T x_v + \frac{\mu}{2} \|x_u - x_v\|^2$$

Assume that f is a closed and convex function. Then f is strongly convex with parameter  $\mu\Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of "\Rightarrow"**: Recall, if g is strongly convex with minimizer x, then

$$g(y) \ge g(x) + \frac{\mu}{2} \|y - x\|^2, \quad \text{for all } y$$

Hence, defining  $x_u = \nabla f^*(u)$  and  $x_v = \nabla f^*(v)$ ,

$$f(x_v) - u^T x_v \ge f(x_u) - u^T x_u + \frac{\mu}{2} \|x_u - x_v\|^2$$

$$f(x_u) - v^T x_u \ge f(x_v) - v^T x_v + \frac{\mu}{2} \|x_u - x_v\|^2$$

Adding these together, using the Cauchy-Schwarz inequality, and rearranging shows that

$$||x_u - x_v||^2 \le \frac{1}{u} ||u - v||^2$$

**Proof of "\Leftarrow"**: for simplicity, call  $g=f^*$  and  $L=\frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant L, so is  $g_x(z)=g(z)-\nabla g(x)^Tz$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T(z-y) + \frac{L}{2}\|z-y\|_2^2$$

**Proof of "** $\Leftarrow$ ": for simplicity, call  $g=f^*$  and  $L=\frac{1}{n}$ . As  $\nabla g$  is Lipschitz with constant L, so is  $q_x(z) = q(z) - \nabla q(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T(z-y) + \frac{L}{2}\|z-y\|_2^2$$

Minimizing each side over z, and rearranging, gives

$$\frac{1}{2I} \|\nabla g(x) - \nabla g(y)\|^2 \le g(y) - g(x) + \nabla g(x)^T (x - y)$$

**Proof of "** $\Leftarrow$ ": for simplicity, call  $g=f^*$  and  $L=\frac{1}{n}$ . As  $\nabla g$  is Lipschitz with constant L, so is  $q_x(z) = q(z) - \nabla q(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T (z - y) + \frac{L}{2} \|z - y\|_2^2$$

Minimizing each side over z, and rearranging, gives

$$\frac{1}{2L}\|\nabla g(x) - \nabla g(y)\|^2 \leq g(y) - g(x) + \nabla g(x)^T(x-y)$$

Exchanging roles of x, y, and adding together, gives

$$\frac{1}{I}\|\nabla g(x) - \nabla g(y)\|^2 \leq (\nabla g(x) - \nabla g(y))^T(x-y)$$

**Proof of "\Leftarrow"**: for simplicity, call  $g=f^*$  and  $L=\frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant L, so is  $g_x(z)=g(z)-\nabla g(x)^Tz$ , hence

$$g_x(z) \le g_x(y) + \nabla g_x(y)^T (z-y) + \frac{L}{2} ||z-y||_2^2$$

Minimizing each side over z, and rearranging, gives

$$\frac{1}{2L}\|\nabla g(x) - \nabla g(y)\|^2 \leq g(y) - g(x) + \nabla g(x)^T(x-y)$$

Exchanging roles of x, y, and adding together, gives

$$\frac{1}{L}\|\nabla g(x) - \nabla g(y)\|^2 \leq (\nabla g(x) - \nabla g(y))^T(x-y)$$

Let  $u = \nabla f(x)$ ,  $v = \nabla g(y)$ ; then  $x \in \partial g^*(u)$ ,  $y \in \partial g^*(v)$ , and the above reads  $(x-y)^T(u-v) \geq \frac{\|u-v\|^2}{L}$ , implying the result.

### Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent: (This is ignoring the role of A, and thus reflects the case when the singular values of A are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\sigma_{\min}(A)^2}}$  (second case).)

• If f is strongly convex with parameter  $\mu_i$  then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .

### Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent: (This is ignoring the role of A, and thus reflects the case when the singular values of A are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\sigma_{\min}(A)^2}}$  (second case).)

- If f is strongly convex with parameter  $\mu_i$  then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .
- ullet If f is strongly convex with parameter  $\mu$  and abla f is Lipschitz with parameter L, then dual gradient ascent with step sizes  $\alpha_k = \frac{2}{\frac{1}{2} + \frac{1}{\epsilon}}$  converges at linear rate  $O(\log(\frac{1}{\epsilon}))$ .



### Convergence guarantees


The following results hold from combining the last fact with what we already know about gradient descent: (This is ignoring the role of A, and thus reflects the case when the singular values of A are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\sigma_{\min}(A)^2}}$  (second case).)

- If f is strongly convex with parameter  $\mu_i$  then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{2})$ .
- ullet If f is strongly convex with parameter  $\mu$  and abla f is Lipschitz with parameter L, then dual gradient ascent with step sizes  $\alpha_k = \frac{2}{\frac{1}{2} + \frac{1}{2}}$  converges at linear rate  $O(\log(\frac{1}{\epsilon}))$ .
- Note that this describes convergence in the dual. Convergence in the primal requires more assumptions

## Example: equality constrained quadratic minimization.

$$f(x) = \frac{1}{2} x^T A x - b^T x \to \min_{x \in \mathbb{R}^n} \qquad \text{subject to} \quad Cx = d, \qquad A \in \mathbb{S}^n_+, C \in \mathbb{R}^{m \times n}, m < n.$$

Quadratic constrained optimization. n=10, m=5,  $\mu$ =1, L=10.




We need to find a minimum of a quadratic function in some linear subspace, defined by the solution of linear equation Cx = d. This is a conditional optimization problem, we start from strongly convex setting.

### Example: equality constrained quadratic minimization.

$$f(x) = \frac{1}{2} x^T A x - b^T x \to \min_{x \in \mathbb{R}^n} \qquad \text{subject to} \quad Cx = d, \qquad A \in \mathbb{S}^n_+, C \in \mathbb{R}^{m \times n}, m < n.$$

Quadratic constrained optimization. n=10, m=5,  $\mu$ =0.001, L=10.



Situation is getting worse as soon as we loose strong convexity, the dual convergence will still be linear, but the rate is very low.



Consider

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad Ax = b$$



Consider

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x=(x_1,\ldots,x_B)\in\mathbb{R}^n$  divides into B blocks of variables, with each  $x_i\in\mathbb{R}^{n_i}$ . We can also partition Aaccordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Consider

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x=(x_1,\ldots,x_B)\in\mathbb{R}^n$  divides into B blocks of variables, with each  $x_i\in\mathbb{R}^{n_i}$ . We can also partition A accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into B separate problems:

$$\begin{split} x^{\text{new}} &\in \arg\min_{x} \left( \sum_{i=1}^{B} f_i(x_i) + u^T A x \right) \\ \Rightarrow x_i^{\text{new}} &\in \arg\min_{x} \left( f_i(x_i) + u^T A_i x_i \right), \quad i = 1, \dots, B \end{split}$$

$$\begin{aligned} & \xrightarrow{} x_i & \in \arg\min_{x_i} \left( f_i(x_i) + u \ A_i x_i \right), \quad i=1,\dots \\ x_i^k & \in \arg\min_{x_i} \left( f_i(x_i) + (u^{k-1})^T A_i x_i \right), \quad i=1,\dots,B \end{aligned}$$

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

Consider

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x=(x_1,\ldots,x_B)\in\mathbb{R}^n$  divides into B blocks of variables, with each  $x_i\in\mathbb{R}^{n_i}$ . We can also partition A accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into B separate problems:

$$\begin{split} x^{\mathsf{new}} &\in \arg\min_{x} \left( \sum_{i=1}^{B} f_i(x_i) + u^T A x \right) \\ \Rightarrow x_i^{\mathsf{new}} &\in \arg\min_{x} \left( f_i(x_i) + u^T A_i x_i \right), \quad i = 1, \dots, B \end{split}$$

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

 $x_i^k \in \arg\min\left(f_i(x_i) + (u^{k-1})^T A_i x_i\right), \quad i = 1, \dots, B$ 

Can think of these steps as:

 $\bullet$   $\mbox{\bf Broadcast:}$  Send u to each of the B processors, each optimizes in parallel to find  $x_i$  .

Consider

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x = (x_1, \dots, x_R) \in \mathbb{R}^n$  divides into B blocks of variables, with each  $x_i \in \mathbb{R}^{n_i}$ . We can also partition A accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into B separate problems:

$$\begin{split} x^{\mathsf{new}} &\in \arg\min_{x} \left( \sum_{i=1}^{B} f_i(x_i) + u^T A x \right) \\ \Rightarrow x_i^{\mathsf{new}} &\in \arg\min_{x} \left( f_i(x_i) + u^T A_i x_i \right), \quad i = 1, \dots, B \end{split}$$

$$(x_i), \quad i=1,\ldots,D$$

Can think of these steps as:

• **Broadcast:** Send u to each of the B processors. each optimizes in parallel to find  $x_i$ .

• **Gather:** Collect  $A_i x_i$  from each processor, update the global dual variable u.

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

 $x_i^k \in \arg\min_{\boldsymbol{x}} \left( f_i(\boldsymbol{x}_i) + (\boldsymbol{u}^{k-1})^T A_i \boldsymbol{x}_i \right), \quad i = 1, \dots, B$ 

#### **Inequality constraints**

#### Consider the optimization problem:

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^{B} A_i x_i \leq b$$



#### **Inequality constraints**

Consider the optimization problem:

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^{B} A_i x_i \leq b$$

Using dual decomposition, specifically the projected subgradient method, the iterative steps can be expressed as:

The primal update step:

$$x_i^k \in \arg\min_{x} \left[ f_i(x_i) + \left( u^{k-1} \right)^T A_i x_i \right], \quad i = 1, \dots, B$$

#### Inequality constraints

Consider the optimization problem:

$$\min_{x} \sum_{i=1}^{B} f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^{B} A_i x_i \leq b$$

Using dual decomposition, specifically the projected subgradient method, the iterative steps can be expressed as:

• The primal update step:

$$x_i^k \in \arg\min_{x} \left[ f_i(x_i) + \left( u^{k-1} \right)^T A_i x_i \right], \quad i = 1, \dots, B$$

The dual update step:

$$u^k = \left(u^{k-1} + \alpha_k \left(\sum_{i=1}^B A_i x_i^k - b\right)\right)_+$$

where  $(u)_+$  denotes the positive part of u, i.e.,  $(u_+)_i = \max\{0,u_i\}$ , for  $i=1,\ldots,m$ .

• System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.

Dual ascent

- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- Resource Constraints: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.



- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints**: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.
- **Dual Update Rule:**

$$u_j^{\mathrm{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$



- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints**: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.
- Dual Update Rule:

$$u_j^{\mathrm{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^{B} A_i x_i$  represents the slacks.

Price Adjustments:



- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints**: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.
- **Dual Update Rule:**

$$u_j^{\mathrm{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

- Price Adjustments:
  - Increase price  $u_i$  if resource j is over-utilized  $(s_i < 0)$ .



- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints**: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.
- Dual Update Rule:

$$u_j^{\mathrm{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

- Price Adjustments:
  - Increase price  $u_i$  if resource j is over-utilized  $(s_i < 0)$ .
  - **Decrease price**  $u_i$  if resource j is under-utilized  $(s_i > 0)$ .

- System Overview: Consider a system with B units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints**: These are limits on shared resources, represented by the rows of A. Each component of the dual variable  $u_i$  represents the price of resource j.
- Dual Update Rule:

$$u_j^{\mathrm{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

- Price Adjustments:
  - Increase price  $u_i$  if resource j is over-utilized  $(s_i < 0)$ .
  - **Decrease price**  $u_i$  if resource j is under-utilized  $(s_i > 0)$ .
  - Never let prices get negative: hence the use of the positive part notation (.).



**Augmented Lagrangian method** 





Augmented Lagrangian method

**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\min_{x} f(x) + \frac{\rho}{2} \|Ax - b\|^{2}$$
 s.t.  $Ax = b$ 



**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\min_{x} f(x) + \frac{\rho}{2} \|Ax - b\|^{2}$$
 s.t.  $Ax = b$ 

where  $\rho>0$  is a parameter. This formulation is clearly equivalent to the original problem. The problem is strongly convex if matrix A has full column rank.



**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\min_{x} f(x) + \frac{\rho}{2} \|Ax - b\|^{2}$$
 s.t.  $Ax = b$ 

where  $\rho > 0$  is a parameter. This formulation is clearly equivalent to the original problem. The problem is strongly convex if matrix A has full column rank.

Dual gradient ascent: The iterative updates are given by:

$$\begin{split} x_k &= \arg\min_{x} \left[ f(x) + (u_{k-1})^T A x + \frac{\rho}{2} \|Ax - b\|^2 \right] \\ u_k &= u_{k-1} + \rho (Ax_k - b) \end{split}$$



#### Notice step size choice $\alpha_k = \rho$ in dual algorithm. Why?

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T A x + \frac{\rho}{2} ||Ax - b||^2$$

over x, we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T \left( u_{k-1} + \rho (Ax_k - b) \right)$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$



Notice step size choice  $\alpha_k = \rho$  in dual algorithm. Why?

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T A x + \frac{\rho}{2} \|Ax - b\|^2$$

over x, we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T \left( u_{k-1} + \rho (Ax_k - b) \right)$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$

This represents the stationarity condition for the original primal problem; under mild conditions,  $Ax_k-b\to 0$  as  $k\to\infty$ , so the KKT conditions are satisfied in the limit and  $x_k$ ,  $u_k$  converge to the solutions.

• Advantage: The augmented Lagrangian gives better convergence.



Notice step size choice  $\alpha_k = \rho$  in dual algorithm. Why?

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T A x + \frac{\rho}{2} \|Ax - b\|^2$$

over x, we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T \left( u_{k-1} + \rho (Ax_k - b) \right)$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$

This represents the stationarity condition for the original primal problem; under mild conditions,  $Ax_k-b\to 0$  as  $k\to\infty$ , so the KKT conditions are satisfied in the limit and  $x_k$ ,  $u_k$  converge to the solutions.


- Advantage: The augmented Lagrangian gives better convergence.
- **Disadvantage:** We lose decomposability! (Separability is ruined)



### Example: equality constrained quadratic minimization.

$$f(x) = \frac{1}{2} x^T A x - b^T x \to \min_{x \in \mathbb{R}^n} \qquad \text{subject to} \quad Cx = d, \qquad A \in \mathbb{S}^n_+, C \in \mathbb{R}^{m \times n}, m < n.$$

Quadratic constrained optimization. n=10, m=5,  $\mu$ =1, L=10.




One can see, clear numerical superiority of the Augmented Lagrangian method both in convex and strongly convex case.



### Example: equality constrained quadratic minimization.

$$f(x) = \frac{1}{2} x^T A x - b^T x \to \min_{x \in \mathbb{R}^n} \qquad \text{subject to} \quad Cx = d, \qquad A \in \mathbb{S}^n_+, C \in \mathbb{R}^{m \times n}, m < n.$$

Quadratic constrained optimization. n=10, m=5,  $\mu$ =0.001, L=10.



One can see, clear numerical superiority of the Augmented Lagrangian method both in convex and strongly convex case.



### Introduction to ADMM





**Alternating direction method of multipliers** or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\min_{x,z} f(x) + g(z)$$

$$\mathrm{s.t.}\ Ax+Bz=c$$



**Alternating direction method of multipliers** or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\min_{x,z} f(x) + g(z)$$
  
s.t.  $Ax + Bz = c$ 

 $\mathbf{S.t.}\ \mathbf{Mw} + \mathbf{Dz} =$ 

$$\min_{x,z} f(x) + g(z) + \frac{\rho}{2} \|Ax + Bz - c\|^2$$
 s.t.  $Ax + Bz = c$ 



Alternating direction method of multipliers or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\min_{x,z} f(x) + g(z)$$
s.t.  $Ax + Bz = c$ 

We augment the objective to include a penalty term for constraint violation:

$$\min_{x,z} f(x) + g(z) + \frac{\rho}{2} \|Ax + Bz - c\|^2$$
 s.t.  $Ax + Bz = c$ 

where  $\rho > 0$  is a parameter. The augmented Lagrangian for this problem is defined as:

$$L_{\rho}(x,z,u) = f(x) + g(z) + u^T(Ax + Bz - c) + \frac{\rho}{2}\|Ax + Bz - c\|^2$$



#### ADMM repeats the following steps, for k = 1, 2, 3, ...:

**1**. Update *x*:

$$x_k = \arg\min_x L_\rho(x, z_{k-1}, u_{k-1})$$



#### ADMM repeats the following steps, for k = 1, 2, 3, ...:

**1**. Update *x*:

$$x_k = \arg\min_x L_\rho(x, z_{k-1}, u_{k-1})$$

**2**. Update *z*:

$$z_k = \arg\min_z L_\rho(x_k, z, u_{k-1})$$



#### ADMM repeats the following steps, for k = 1, 2, 3, ...:

**1**. Update *x*:

$$x_k = \arg\min_{x} L_{\rho}(x, z_{k-1}, u_{k-1})$$

2. Update z:

$$z_k = \arg\min_z L_\rho(x_k, z, u_{k-1})$$

3. Update u:

$$u_k = u_{k-1} + \rho (Ax_k + Bz_k - c)$$

**⊕** ດ **ø** 

#### ADMM repeats the following steps, for k = 1, 2, 3, ...:

**1**. Update *x*:

$$x_k = \arg\min_{x} L_{\rho}(x, z_{k-1}, u_{k-1})$$

2. Update z:

$$z_k = \arg\min_z L_\rho(x_k, z, u_{k-1})$$

3. Update u:

$$u_k = u_{k-1} + \rho (Ax_k + Bz_k - c)$$

**⊕** ດ **ø** 

#### ADMM repeats the following steps, for k = 1, 2, 3, ...:

1. Update x:

$$x_k = \arg\min_{x} L_\rho(x, z_{k-1}, u_{k-1})$$

2. Update z:

$$z_k = \arg\min_z L_\rho(x_k, z, u_{k-1})$$

3. Update u:


$$u_k = u_{k-1} + \rho(Ax_k + Bz_k - c)$$

**Note:** The usual method of multipliers would replace the first two steps by a joint minimization:

$$(x^{(k)}, z^{(k)}) = \arg\min_{x} L_{\rho}(x, z, u^{(k-1)})$$

**⊕** ດ **ø** 

### **Example: Alternating Projections**



Consider finding a point in the intersection of convex sets  $U, V \subseteq \mathbb{R}^n$ :

$$\min_{x} I_{U}(x) + I_{V}(x)$$

To transform this problem into ADMM form, we express it as:

$$\min_{x,z} I_U(x) + I_V(z) \quad \text{subject to} \quad x-z = 0$$

Each ADMM cycle involves two projections:

$$\begin{split} x_k &= \arg\min_x P_U \left( z_{k-1} - w_{k-1} \right) \\ z_k &= \arg\min_z P_V \left( x_k + w_{k-1} \right) \\ w_k &= w_{k-1} + x_k - z_k \end{split}$$



#### **Sources**

• Ryan Tibshirani. Convex Optimization 10-725



Introduction to ADMM

