

### **Conditional methods**





### Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

♥ 0 @

### Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

♥ 0 @

### Unconstrained optimization

### Constrained optimization

$$\min_{x \in \mathbb{R}^n} f(x) \qquad \qquad \min_{x \in S} f(x)$$

- Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution. Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.



### Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

- $\min_{x \in S} f(x)$
- Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.
- The solution has to be inside the set S.



#### Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

### Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.
- The solution has to be inside the set S.
- Example:

$$\frac{1}{2} ||Ax - b||_2^2 \to \min_{\|x\|_2^2 \le 1}$$

റമ

#### Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

### Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.
- The solution has to be inside the set S.
- Example:

$$\frac{1}{2} ||Ax - b||_2^2 \to \min_{\|x\|_2^2 \le 1}$$

റമ

### Unconstrained optimization

### Constrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

$$\min_{x \in S} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

- Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.
- The solution has to be inside the set S.
- Example:

$$\frac{1}{2} \|Ax - b\|_2^2 \to \min_{\|x\|_2^2 \le 1}$$

Gradient Descent is a great way to solve unconstrained problem

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k) \tag{GD}$$

Is it possible to tune GD to fit constrained problem?





### Unconstrained optimization

### Constrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

• Any point  $x_0 \in \mathbb{R}^n$  is feasible and could be a solution.

- $\min_{x \in S} f(x)$
- Not all  $x \in \mathbb{R}^n$  are feasible and could be a solution.
- The solution has to be inside the set S.
- Example:

$$\frac{1}{2}\|Ax - b\|_2^2 \to \min_{\|x\|_2^2 \le 1}$$

Gradient Descent is a great way to solve unconstrained problem


$$x_{k+1} = x_k - \alpha_k \nabla f(x_k) \tag{GD}$$

Is it possible to tune GD to fit constrained problem?

**Yes**. We need to use projections to ensure feasibility on every iteration.



### **Example: White-box Adversarial Attacks**



• Mathematically, a neural network is a function  $f(\boldsymbol{w};\boldsymbol{x})$ 

 $\mathsf{P}\mathsf{u}\mathsf{c}\mathsf{y}\mathsf{h}\mathsf{o}\mathsf{k}$  1: Source

⊕ o a

### **Example: White-box Adversarial Attacks**




Рисунок 1: Source

- Mathematically, a neural network is a function f(w;x)
- $\bullet$  Typically, input x is given and network weights w optimized

⊕ n ø

### **Example: White-box Adversarial Attacks**




Рисунок 1: Source

- Mathematically, a neural network is a function f(w;x)
- $\begin{tabular}{ll} {\bf Typically, input} $x$ is given and network weights \\ $w$ optimized \\ \end{tabular}$
- Could also freeze weights w and optimize x, adversarially!

$$\min_{\delta} \operatorname{size}(\delta) \quad \text{s.t.} \quad \operatorname{pred}[f(w; x + \delta)] \neq y$$
 or

 $\max_{\delta} l(w; x+\delta, y) \text{ s.t. } \operatorname{size}(\delta) \leq \epsilon, \ 0 \leq x+\delta \leq 1$ 

 $f \to \min_{T, T}$ 

Conditional methods




Рисунок 2: Suppose, we start from a point  $x_k$ .

∌ດ Ø




Рисунок 3: And go in the direction of  $-\nabla f(x_k)$ .

♥ റ ⊘




Рисунок 4: Occasionally, we can end up outside the feasible set.

⊕ ი დ





Рисунок 5: Solve this little problem with projection!

**₹ 6 0** 

$$x_{k+1} = \operatorname{proj}_S\left(x_k - \alpha_k \nabla f(x_k)\right) \qquad \Leftrightarrow \qquad \begin{aligned} y_k &= x_k - \alpha_k \nabla f(x_k) \\ x_{k+1} &= \operatorname{proj}_S\left(y_k\right) \end{aligned}$$





Projection



The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y},S,\|\cdot\|)=\inf\{\|x-y\|\mid x\in S\}$$

The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x-y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point  $\mathbf{y} \in \mathbb{R}^n$  on set  $S \subseteq \mathbb{R}^n$  is a point  $\operatorname{proj}_S(\mathbf{y}) \in S$ :

$$\operatorname{proj}_{S}(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_{2}^{2}$$

Projection

⊕ ი ⊘

The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x-y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point  $\mathbf{y} \in \mathbb{R}^n$  on set  $S \subseteq \mathbb{R}^n$  is a point  $\operatorname{proj}_S(\mathbf{y}) \in S$ :

$$\operatorname{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

• Sufficient conditions of existence of a projection. If  $S \subseteq \mathbb{R}^n$  - closed set, then the projection on set S exists for any point.



The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y},S,\|\cdot\|)=\inf\{\|x-y\|\mid x\in S\}$$

We will focus on Euclidean projection (other options are possible) of a point  $\mathbf{y} \in \mathbb{R}^n$  on set  $S \subseteq \mathbb{R}^n$  is a point  $\operatorname{proj}_S(\mathbf{y}) \in S$ :

$$\operatorname{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- Sufficient conditions of existence of a projection. If  $S \subseteq \mathbb{R}^n$  closed set, then the projection on set S exists for any point.
- Sufficient conditions of uniqueness of a projection. If  $S \subseteq \mathbb{R}^n$  closed convex set, then the projection on set S is unique for any point.

Projection

The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x-y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point  $y \in \mathbb{R}^n$  on set  $S \subseteq \mathbb{R}^n$  is a point  $\operatorname{\mathsf{proj}}_S(\mathbf{y}) \in S$ :

$$\operatorname{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- Sufficient conditions of existence of a projection. If  $S \subseteq \mathbb{R}^n$  closed set, then the projection on set S exists for any point.
- Sufficient conditions of uniqueness of a projection. If  $S \subseteq \mathbb{R}^n$  closed convex set, then the projection on set S is unique for any point.
- If a set is open, and a point is beyond this set, then its projection on this set may not exist.

The distance d from point  $\mathbf{y} \in \mathbb{R}^n$  to closed set  $S \subset \mathbb{R}^n$ :

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x-y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point  $y \in \mathbb{R}^n$  on set  $S \subseteq \mathbb{R}^n$  is a point  $\operatorname{\mathsf{proj}}_S(\mathbf{y}) \in S$ :

$$\operatorname{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- Sufficient conditions of existence of a projection. If  $S \subseteq \mathbb{R}^n$  closed set, then the projection on set S exists for any point.
- Sufficient conditions of uniqueness of a projection. If  $S \subseteq \mathbb{R}^n$  closed convex set, then the projection on set S is unique for any point.
- If a set is open, and a point is beyond this set, then its projection on this set may not exist.
- If a point is in set, then its projection is the point itself.

#### **i** Theorem

Let  $S\subseteq\mathbb{R}^n$  be closed and convex,  $\forall x\in S,y\in\mathbb{R}^n.$  Then

$$\langle y - \operatorname{proj}_S(y), \mathbf{x} - \operatorname{proj}_S(y) \rangle \leq 0 \tag{1}$$

$$\|x - \mathrm{proj}_S(y)\|^2 + \|y - \mathrm{proj}_S(y)\|^2 \leq \|x - y\|^2 \tag{2}$$

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|)=\|x-y\|^2$  over S. By first-order characterization of optimality.




Рисунок 7: Obtuse or straight angle should be for any point  $x\in S$ 

Projection

#### i Theorem


Let  $S\subseteq\mathbb{R}^n$  be closed and convex,  $\forall x\in S,y\in\mathbb{R}^n.$  Then

$$\langle y - \operatorname{proj}_S(y), \mathbf{x} - \operatorname{proj}_S(y) \rangle \leq 0 \tag{1}$$

$$\|x - \mathrm{proj}_S(y)\|^2 + \|y - \mathrm{proj}_S(y)\|^2 \le \|x - y\|^2 \tag{2}$$

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|) = \|x-y\|^2$  over S. By first-order characterization of optimality.

$$\nabla d(\operatorname{proj}_{S}(y))^{T}(x-\operatorname{proj}_{S}(y))\geq 0$$





#### i Theorem

Let  $S\subseteq\mathbb{R}^n$  be closed and convex,  $\forall x\in S,y\in\mathbb{R}^n.$  Then

$$\langle y - \mathrm{proj}_S(y), \mathbf{x} - \mathrm{proj}_S(y) \rangle \leq 0 \tag{1}$$

$$\|x - \mathrm{proj}_S(y)\|^2 + \|y - \mathrm{proj}_S(y)\|^2 \leq \|x - y\|^2 \tag{2}$$

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|) = \|x-y\|^2$  over S. By first-order characterization of optimality.

$$\nabla d(\operatorname{proj}_S(y))^T(x-\operatorname{proj}_S(y)) \geq 0$$

$$2\left(\operatorname{proj}_{\mathcal{S}}(y)-y\right)^T(x-\operatorname{proj}_{\mathcal{S}}(y))\geq 0$$




Рисунок 7: Obtuse or straight angle should be for any point  $x \in S$ 

Projection

#### i Theorem

Let  $S \subseteq \mathbb{R}^n$  be closed and convex,  $\forall x \in S, y \in \mathbb{R}^n$ . Then


$$\langle y - \mathsf{proj}_S(y), \mathbf{x} - \mathsf{proj}_S(y) \rangle \leq 0$$

(1)

$$\|x - \mathrm{proj}_S(y)\|^2 + \|y - \mathrm{proj}_S(y)\|^2 \leq \|x - y\|^2 \tag{2}$$

1.  $proj_{S}(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|) = \|x-y\|^2$  over S. By first-order characterization of optimality.

$$\begin{split} \nabla d(\operatorname{proj}_S(y))^T(x - \operatorname{proj}_S(y)) &\geq 0 \\ 2\left(\operatorname{proj}_S(y) - y\right)^T(x - \operatorname{proj}_S(y)) &\geq 0 \\ \left(y - \operatorname{proj}_S(y)\right)^T(x - \operatorname{proj}_S(y)) &\leq 0 \end{split}$$



S

Рисунок 7: Obtuse or straight angle should be for any point  $x \in S$ 

 $\operatorname{proj}_S(y)$ 

#### **i** Theorem

Let  $S\subseteq\mathbb{R}^n$  be closed and convex,  $\forall x\in S,y\in\mathbb{R}^n.$  Then


$$\langle y - \operatorname{proj}_S(y), \mathbf{x} - \operatorname{proj}_S(y) \rangle \leq 0 \tag{1}$$

$$\|x - \mathrm{proj}_S(y)\|^2 + \|y - \mathrm{proj}_S(y)\|^2 \leq \|x - y\|^2 \tag{2}$$

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|)=\|x-y\|^2$  over S. By first-order characterization of optimality.

$$\begin{split} \nabla d(\operatorname{proj}_S(y))^T(x - \operatorname{proj}_S(y)) &\geq 0 \\ 2\left(\operatorname{proj}_S(y) - y\right)^T(x - \operatorname{proj}_S(y)) &\geq 0 \\ (y - \operatorname{proj}_S(y))^T(x - \operatorname{proj}_S(y)) &\leq 0 \end{split}$$

2. Use cosine rule  $2x^Ty=\|x\|^2+\|y\|^2-\|x-y\|^2$  with  $x=x-\operatorname{proj}_S(y)$  and  $y=y-\operatorname{proj}_S(y)$ . By the first property of the theorem:





### i Theorem

Let  $S \subseteq \mathbb{R}^n$  be closed and convex,  $\forall x \in S, y \in \mathbb{R}^n$ . Then


$$\langle y - \operatorname{proj}_{S}(y), \mathbf{x} - \operatorname{proj}_{S}(y) \rangle \le 0 \tag{1}$$
$$\|x - \operatorname{proj}_{S}(y)\|^{2} + \|y - \operatorname{proj}_{S}(y)\|^{2} \le \|x - y\|^{2} \tag{2}$$

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|)=\|x-y\|^2$  over S. By first-order characterization of optimality.

$$\nabla d(\operatorname{proj}_S(y))^T(x-\operatorname{proj}_S(y)) \geq 0$$
 
$$2\left(\operatorname{proj}_S(y)-y\right)^T(x-\operatorname{proj}_S(y)) \geq 0$$
 
$$\left(y-\operatorname{proj}_S(y)\right)^T(x-\operatorname{proj}_S(y)) \leq 0$$

2. Use cosine rule  $2x^Ty=\|x\|^2+\|y\|^2-\|x-y\|^2$  with  $x=x-\operatorname{proj}_S(y)$  and  $y=y-\operatorname{proj}_S(y)$ . By the first property of the theorem:

$$0 \ge 2x^T y = \|x - \operatorname{proj}_{S}(y)\|^2 + \|y + \operatorname{proj}_{S}(y)\|^2 - \|x - y\|^2$$



### i Theorem

Let  $S\subseteq\mathbb{R}^n$  be closed and convex,  $\forall x\in S,y\in\mathbb{R}^n$ . Then  $\langle y-\operatorname{proj}_S(y),\mathbf{x}-\operatorname{proj}_S(y)\rangle\leq 0 \tag{1}$   $\|x-\operatorname{proj}_S(y)\|^2+\|y-\operatorname{proj}_S(y)\|^2\leq \|x-y\|^2 \tag{2}$ 

1.  $\operatorname{proj}_S(y)$  is minimizer of differentiable convex function  $d(y,S,\|\cdot\|) = \|x-y\|^2$  over S. By first-order characterization of optimality.  $\nabla d(xx; \langle x \rangle) T(x, x, y; \langle x \rangle) \geq 0$ 

$$\nabla d(\mathsf{proj}_S(y))^T(x-\mathsf{proj}_S(y)) \geq 0$$
 
$$2\left(\mathsf{proj}_S(y)-y\right)^T(x-\mathsf{proj}_S(y)) \geq 0$$
 
$$\left(y-\mathsf{proj}_S(y)\right)^T(x-\mathsf{proj}_S(y)) \leq 0$$

2. Use cosine rule  $2x^Ty=\|x\|^2+\|y\|^2-\|x-y\|^2$  with  $x=x-\operatorname{proj}_S(y)$  and  $y=y-\operatorname{proj}_S(y)$ . By the first property of the theorem:

$$0 \geq 2x^Ty = \|x - \mathrm{proj}_S(y)\|^2 + \|y + \mathrm{proj}_S(y)\|^2 - \|x - y\|^2$$

• A function f is called non-expansive if f is L-Lipschitz with  $L \leq 1$   $^1$ . That is, for any two points  $x,y \in \text{dom} f$ ,

$$\|f(x)-f(y)\|\leq L\|x-y\|, \text{ where } L\leq 1.$$

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

 $<sup>^{1}</sup>$ Non-expansive becomes contractive if L < 1.

• A function f is called non-expansive if f is L-Lipschitz with  $L \leq 1$   $^1$ . That is, for any two points  $x,y \in \text{dom} f$ ,

$$||f(x) - f(y)|| \le L||x - y||$$
, where  $L \le 1$ .

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

• Projection operator is non-expansive:

$$\|\text{proj}(x) - \text{proj}(y)\|_2 \le \|x - y\|_2.$$

 $<sup>^{1}</sup>$ Non-expansive becomes contractive if L < 1.

• A function f is called non-expansive if f is L-Lipschitz with  $L \leq 1$   $^1$ . That is, for any two points  $x,y \in \text{dom} f$ ,

$$||f(x) - f(y)|| \le L||x - y||$$
, where  $L \le 1$ .

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

Projection operator is non-expansive:

$$\|\mathrm{proj}(x) - \mathrm{proj}(y)\|_2 \leq \|x - y\|_2.$$

• Next: variational characterization implies non-expansiveness. i.e.,

$$\langle y - \operatorname{proj}(y), x - \operatorname{proj}(y) \rangle \leq 0 \quad \forall x \in S \qquad \Rightarrow \qquad \|\operatorname{proj}(x) - \operatorname{proj}(y)\|_2 \leq \|x - y\|_2.$$

 $<sup>^{1}</sup>$ Non-expansive becomes contractive if L < 1.

Shorthand notation: let  $\pi=\operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x).$ 

Shorthand notation: let  $\pi=\operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization  $\slash$  obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \le 0 \quad \forall x \in S.$$

(3)

Shorthand notation: let  $\pi = \operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle < 0 \quad \forall x \in S.$$

Replace x by  $\pi(x)$  in Уравнение 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \le 0.$$
 (4)

(3)

Replace x by  $\pi(x)$  in Уравнение 3

Shorthand notation: let  $\pi = \operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization / obtuse angle inequality

 $\langle y - \pi(y), \pi(x) - \pi(y) \rangle < 0.$ 

$$\in \mathcal{S}$$
.

$$|y\rangle \leq 0 \quad \forall$$

$$|y\rangle \leq 0 \quad \forall$$

$$\langle y \rangle \leq 0 \quad \forall$$

$$|y\rangle \leq 0 \quad \forall y$$

$$\langle y \rangle \leq 0$$

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S.$$

(4)

$$r$$
 and

$$\mathsf{and}\ x$$

 $\langle x - \pi(x), \pi(y) - \pi(x) \rangle < 0.$ 

(3)

(5)

$$x$$
 and

$$x$$
 and

Replace 
$$y$$
 by  $x$  and  $x$  by  $\pi(y)$  in Уравнение 3

e 
$$y$$
 by

$$\mathsf{e}\ y$$
 l

ace 
$$y$$

Shorthand notation: let  $\pi = \operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S.$$

$$\langle y | n(y), w | n(y) \rangle \equiv 0$$

Replace 
$$x$$
 by  $\pi(x)$  in Уравнение 3 
$$\langle y-\pi(y),\pi(x)-\pi(y)\rangle \leq 0. \tag{4}$$

Replace 
$$y$$
 by  $x$  and  $x$  by  $\pi(y)$  in Уравнение 3

$$x$$
 and

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle < 0.$$

(Уравнение 4)+(Уравнение 5) will cancel 
$$\pi(y)-\pi(x)$$
, not good. So flip the sign of (Уравнение 5) gives

 $\langle \pi(x) - x, \pi(x) - \pi(y) \rangle < 0.$ 

(3)

(5)

(6)

 $\langle y - \pi(y) + \pi(x) - x, \pi(x) - \pi(y) \rangle < 0$ 

Shorthand notation: let  $\pi = \operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \le 0 \quad \forall x \in S.$$

Replace 
$$x$$
 by  $\pi(x)$  in Уравнение 3 Replace  $y$  by  $x$  and  $x$  by  $\pi(y)$  in Уравнение 3

 $\langle y-x,\pi(x)-\pi(y)\rangle < -\langle \pi(x)-\pi(y),\pi(x)-\pi(y)\rangle$ 

 $\langle y - x, \pi(y) - \pi(x) \rangle > \|\pi(x) - \pi(y)\|_2^2$  $\|(y-x)^{\top}(\pi(y)-\pi(x))\|_{2} > \|\pi(x)-\pi(y)\|_{2}^{2}$ 

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \le 0.$$
 (4)

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle < 0.$$

(Уравнение 4)+(Уравнение 5) will cancel 
$$\pi(y)-\pi(x)$$
, not good. So flip the sign of (Уравнение 5) gives

$$\langle \pi(x) - x, \pi(x) - \pi(y) \rangle \le 0.$$

(3)

(5)

(6)

Shorthand notation: let  $\pi = \operatorname{proj}$  and  $\pi(x)$  denotes  $\operatorname{proj}(x)$ .

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle < 0 \quad \forall x \in S.$$

 $\langle y - x, \pi(y) - \pi(x) \rangle > \|\pi(x) - \pi(y)\|_2^2$ 

 $\|(y-x)^{\top}(\pi(y)-\pi(x))\|_{2} > \|\pi(x)-\pi(y)\|_{2}^{2}$ 

Replace x by  $\pi(x)$  in Уравнение 3  $\langle u - \pi(u), \pi(x) - \pi(u) \rangle < 0.$ 

 $\langle y - \pi(y) + \pi(x) - x, \pi(x) - \pi(y) \rangle < 0$ 

(4)

$$x, \pi(x)$$

$$\langle \pi(x) - x, \pi(x) - \pi(y) \rangle \le 0.$$

 $\langle y-x,\pi(x)-\pi(y)\rangle \leq -\langle \pi(x)-\pi(y),\pi(x)-\pi(y)\rangle \quad \|y-x\|_2\|\pi(y)-\pi(x)\|_2, \text{ we get }$ 

Replace y by x and x by  $\pi(y)$  in Уравнение 3

(Уравнение 4)+(Уравнение 5) will cancel 
$$\pi(y)-\pi(x)$$
, not good. So flip the sign of (Уравнение 5) gives

 $\|y-x\|_2 \|\pi(y)-\pi(x)\|_2 \ge \|\pi(x)-\pi(y)\|_2^2$ .

Cancels  $\|\pi(x) - \pi(y)\|_2$  finishes the proof.

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle \le 0.$$

(3)

(5)



Find  $\pi_S(y)=\pi$ , if  $S=\{x\in\mathbb{R}^n\mid \|x-x_0\|\leq R\}$ ,  $y\notin S$ 

Find  $\pi_S(y)=\pi$  , if  $S=\{x\in\mathbb{R}^n\mid \|x-x_0\|\leq R\}$  ,  $y\notin S$ 

Build a hypothesis from the figure:  $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$ 

Find  $\pi_S(y) = \pi$ , if  $S = \{x \in \mathbb{R}^n \mid ||x - x_0|| \le R\}$ ,  $y \notin S$ 

Build a hypothesis from the figure:  $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$ Check the inequality for a convex closed set:  $(\pi - y)^T(x - \pi) \ge 0$ 

Find  $\pi_S(y) = \pi$ , if  $S = \{x \in \mathbb{R}^n \mid ||x - x_0|| \le R\}$ ,  $y \notin S$ 

Build a hypothesis from the figure:  $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$ 

Check the inequality for a convex closed set:  $(\pi-y)^T(x-\pi)>0$ 

$$\begin{split} \left(x_{0} - y + R\frac{y - x_{0}}{\|y - x_{0}\|}\right)^{T} \left(x - x_{0} - R\frac{y - x_{0}}{\|y - x_{0}\|}\right) &= \\ \left(\frac{(y - x_{0})(R - \|y - x_{0}\|)}{\|y - x_{0}\|}\right)^{T} \left(\frac{(x - x_{0})\|y - x_{0}\| - R(y - x_{0})}{\|y - x_{0}\|}\right) &= \\ \frac{R - \|y - x_{0}\|}{\|y - x_{0}\|^{2}} \left(y - x_{0}\right)^{T} \left((x - x_{0})\|y - x_{0}\| - R\left(y - x_{0}\right)\right) &= \\ \frac{R - \|y - x_{0}\|}{\|y - x_{0}\|} \left(\left(y - x_{0}\right)^{T} \left(x - x_{0}\right) - R\|y - x_{0}\|\right) &= \\ \left(R - \|y - x_{0}\|\right) \left(\frac{(y - x_{0})^{T} (x - x_{0})}{\|y - x_{0}\|} - R\right) \end{split}$$

Find  $\pi_S(y) = \pi$ , if  $S = \{x \in \mathbb{R}^n \mid ||x - x_0|| < R\}, \ u \notin S$ 

Build a hypothesis from the figure:  $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$ Check the inequality for a convex closed set:  $(\pi-y)^T(x-\pi)\geq 0$ 

$$\left( x_0 - y + R \frac{y - x_0}{\|y - x_0\|} \right)^T \left( x - x_0 - R \frac{y - x_0}{\|y - x_0\|} \right) = \\ \left( \frac{(y - x_0)(R - \|y - x_0\|)}{\|y - x_0\|} \right)^T \left( \frac{(x - x_0)\|y - x_0\| - R(y - x_0)}{\|y - x_0\|} \right) =$$

$$\frac{R-\left\|y-x_{0}\right\|}{\left\|y-x_{0}\right\|^{2}}\left(y-x_{0}\right)^{T}\left(\left(x-x_{0}\right)\left\|y-x_{0}\right\|-R\left(y-x_{0}\right)\right)=$$

$$\frac{R - \|y - x_0\|}{\|y - x_0\|} \left( (y - x_0)^T (x - x_0) - R\|y - x_0\| \right) = \frac{1}{\|y - x_0\|} \left( (y - x_0)^T (x - x_0) - R\|y - x_0\| \right) = \frac{1}{\|y - x_0\|} \left( (y - x_0)^T (x - x_0) - R\|y - x_0\| \right)$$

$$(R - \|y - x_0\|) \left( \frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R \right)$$

The first factor is negative for point selection y. The second factor is also negative, which follows from the Cauchy-Bunyakovsky

Find  $\pi_S(y) = \pi$ , if  $S = \{x \in \mathbb{R}^n \mid ||x - x_0|| < R\}, \ u \notin S$ 

Build a hypothesis from the figure:  $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$ 

Check the inequality for a convex closed set: 
$$(\pi-y)^T(x-\pi) \geq 0$$

$$\left(x_0-y+R\frac{y-x_0}{\|y-x_0\|}\right)^T\left(x-x_0-R\frac{y-x_0}{\|y-x_0\|}\right)=$$

$$\left( \frac{(y-x_0)(R-\|y-x_0\|)}{\|y-x_0\|} \right)^T \left( \frac{(x-x_0)\|y-x_0\|-R(y-x_0)}{\|y-x_0\|} \right) = \\ \frac{(y-x_0)^T(x-x_0) \leq \|y-x_0\|\|x-x_0\|}{\|y-x_0\|} \\ \frac{R-\|y-x_0\|}{\|y-x_0\|} \left( y-x_0 \right)^T \left( (x-x_0)\|y-x_0\|-R(y-x_0) \right) = \\ \frac{(y-x_0)^T(x-x_0) \leq \|y-x_0\|\|x-x_0\|}{\|y-x_0\|} - R \leq \frac{\|y-x_0\|\|x-x_0\|}{\|y-x_0\|}$$

$$\frac{R-\left\|y-x_{0}\right\|}{\left\|y-x_{0}\right\|^{2}}\left(y-x_{0}\right)^{T}\left(\left(x-x_{0}\right)\left\|y-x_{0}\right\|-R\left(y-x_{0}\right)\right)=$$


$$\frac{R-\left\|y-x_{0}\right\|^{2}}{\left\|y-x_{0}\right\|}\left(\left(y-x_{0}\right)^{T}\left(x-x_{0}\right)-R\|y-x_{0}\|\right)=$$

$$(R - \|y - x_0\|) \left( \frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R \right)$$

The first factor is negative for point selection y. The second factor is also negative, which

follows from the Cauchy-Bunyakovsky

$$\frac{(y-x_0)^T(x-x_0)}{|y-x_0||} - R \le \frac{\|y-x_0\|\|x-x_0\|}{\|y-x_0\|}$$



#### Example: projection on the halfspace

Find  $\pi_S(y)=\pi$ , if  $S=\{x\in\mathbb{R}^n\mid c^Tx=b\}$ ,  $y\notin S$ . Build a hypothesis from the figure:  $\pi=y+\alpha c$ . Coefficient  $\alpha$  is chosen so that  $\pi\in S$ :  $c^T\pi=b$ , so:

### **Example:** projection on the halfspace

Find  $\pi_S(y)=\pi$ , if  $S=\{x\in\mathbb{R}^n\mid c^Tx=b\}$ ,  $y\notin S$ . Build a hypothesis from the figure:  $\pi=y+\alpha c$ . Coefficient  $\alpha$  is chosen so that  $\pi\in S$ :  $c^T\pi=b$ , so:




Рисунок 9: Hyperplane

### Example: projection on the halfspace

Find  $\pi_S(y)=\pi$ , if  $S=\{x\in\mathbb{R}^n\mid c^Tx=b\}$ ,  $y\notin S$ . Build a hypothesis from the figure:  $\pi=y+\alpha c$ . Coefficient  $\alpha$  is chosen so that  $\pi\in S$ :  $c^T\pi=b$ . so:




Рисунок 9: Hyperplane

$$c^{T}(y + \alpha c) = b$$

$$c^{T}y + \alpha c^{T}c = b$$

$$c^{T}y = b - \alpha c^{T}c$$

Check the inequality for a convex closed set:  $(\pi - u)^T (x - \pi) > 0$ 


$$\begin{split} T(x-\pi) &\geq 0 \\ (y+\alpha c-y)^T(x-y-\alpha c) &= \\ \alpha c^T(x-y-\alpha c) &= \\ \alpha (c^Tx) - \alpha (c^Ty) - \alpha^2 (c^Tc) &= \\ \alpha b - \alpha (b - \alpha c^Tc) - \alpha^2 c^Tc &= \\ \alpha b - \alpha b + \alpha^2 c^Tc - \alpha^2 c^Tc &= 0 \geq 0 \end{split}$$

### **Projected Gradient Descent (PGD)**



#### Idea

$$x_{k+1} = \operatorname{proj}_S\left(x_k - \alpha_k \nabla f(x_k)\right) \qquad \Leftrightarrow \qquad \begin{aligned} y_k &= x_k - \alpha_k \nabla f(x_k) \\ x_{k+1} &= \operatorname{proj}_S\left(y_k\right) \end{aligned}$$





#### i Theorem

Let  $f: \mathbb{R}^n \to \mathbb{R}$  be an L-smooth convex function. Then, for any  $x,y \in \mathbb{R}^n$ , the following inequality holds:

$$\begin{split} f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 & \leq f(y) \text{ or, equivalently,} \\ \| \nabla f(y) - \nabla f(x) \|_2^2 = & \| \nabla f(x) - \nabla f(y) \|_2^2 \leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

#### Proof

1. To prove this, we'll consider another function  $\varphi(y) = f(y) - \langle \nabla f(x), y \rangle$ . It is obviously a convex function (as a sum of convex functions). And it is easy to verify, that it is an L-smooth function by definition, since  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x)$  and  $\|\nabla \varphi(y_1) - \nabla \varphi(y_2)\| = \|\nabla f(y_1) - \nabla f(y_2)\| \le L\|y_1 - y_2\|$ .

Projected Gradient Descent (PGD)

#### i Theorem

Let  $f: \mathbb{R}^n \to \mathbb{R}$  be an L-smooth convex function. Then, for any  $x,y \in \mathbb{R}^n$ , the following inequality holds:

$$\begin{split} f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 & \leq f(y) \text{ or, equivalently,} \\ \| \nabla f(y) - \nabla f(x) \|_2^2 = & \| \nabla f(x) - \nabla f(y) \|_2^2 \leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

#### Proof

- 1. To prove this, we'll consider another function  $\varphi(y) = f(y) \langle \nabla f(x), y \rangle$ . It is obviously a convex function (as a sum of convex functions). And it is easy to verify, that it is an L-smooth function by definition, since  $\nabla \varphi(y) = \nabla f(y) \nabla f(x)$  and  $\|\nabla \varphi(y_1) \nabla \varphi(y_2)\| = \|\nabla f(y_1) \nabla f(y_2)\| \le L\|y_1 y_2\|$ .
- 2. Now let's consider the smoothness parabolic property for the  $\varphi(y)$  function:

⊕ ი

#### i Theorem

Let  $f: \mathbb{R}^n \to \mathbb{R}$  be an L-smooth convex function. Then, for any  $x,y \in \mathbb{R}^n$ , the following inequality holds:

$$\begin{split} f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 & \leq f(y) \text{ or, equivalently,} \\ \| \nabla f(y) - \nabla f(x) \|_2^2 = & \| \nabla f(x) - \nabla f(y) \|_2^2 \leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

#### Proof

- 1. To prove this, we'll consider another function  $\varphi(y) = f(y) \langle \nabla f(x), y \rangle$ . It is obviously a convex function (as a sum of convex functions). And it is easy to verify, that it is an L-smooth function by definition, since  $\nabla \varphi(y) = \nabla f(y) \nabla f(x)$  and  $\|\nabla \varphi(y_1) \nabla \varphi(y_2)\| = \|\nabla f(y_1) \nabla f(y_2)\| \le L\|y_1 y_2\|$ .
- 2. Now let's consider the smoothness parabolic property for the  $\varphi(y)$  function:

$$\varphi(y) \le \varphi(x) + \langle \nabla \varphi(x), y - x \rangle + \frac{L}{2} \|y - x\|_2^2$$

Projected Gradient Descent (PGD)

#### i Theorem

Let  $f: \mathbb{R}^n \to \mathbb{R}$  be an L-smooth convex function. Then, for any  $x, y \in \mathbb{R}^n$ , the following inequality holds:

$$\begin{split} f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 & \leq f(y) \text{ or, equivalently,} \\ \| \nabla f(y) - \nabla f(x) \|_2^2 = \| \nabla f(x) - \nabla f(y) \|_2^2 & \leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

#### **Proof**

- 1. To prove this, we'll consider another function  $\varphi(y) = f(y) \langle \nabla f(x), y \rangle$ . It is obviously a convex function (as a sum of convex functions). And it is easy to verify, that it is an L-smooth function by definition, since  $\nabla \varphi(y) = \nabla f(y) \nabla f(x)$  and  $\|\nabla \varphi(y_1) \nabla \varphi(y_2)\| = \|\nabla f(y_1) \nabla f(y_2)\| \le L\|y_1 y_2\|$ .
- 2. Now let's consider the smoothness parabolic property for the  $\varphi(y)$  function:

$$\begin{split} \varphi(y) & \leq \varphi(x) + \langle \nabla \varphi(x), y - x \rangle + \frac{L}{2} \|y - x\|_2^2 \\ & \stackrel{x:=y,y:=y-\frac{1}{L}\nabla \varphi(y)}{\varphi\left(y - \frac{1}{L}\nabla \varphi(y)\right)} \leq \varphi(y) + \left\langle \nabla \varphi(y), -\frac{1}{L}\nabla \varphi(y) \right\rangle + \frac{1}{2L} \|\nabla \varphi(y)\|_2^2 \end{split}$$



#### i Theorem

Let  $f:\mathbb{R}^n\to\mathbb{R}$  be an L-smooth convex function. Then, for any  $x,y\in\mathbb{R}^n$ , the following inequality holds:

$$f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \|\nabla f(x) - \nabla f(y)\|_2^2 \leq f(y) \text{ or, equivalently,}$$
 
$$\|\nabla f(y) - \nabla f(x)\|_2^2 = \|\nabla f(x) - \nabla f(y)\|_2^2 \leq 2L\left(f(x) - f(y) - \langle \nabla f(y), x - y \rangle\right)$$

#### Proof

- 1. To prove this, we'll consider another function  $\varphi(y) = f(y) \langle \nabla f(x), y \rangle$ . It is obviously a convex function (as a sum of convex functions). And it is easy to verify, that it is an L-smooth function by definition, since  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x)$  and  $\|\nabla \varphi(y_1) - \nabla \varphi(y_2)\| = \|\nabla f(y_1) - \nabla f(y_2)\| \le L\|y_1 - y_2\|$ .
  - 2. Now let's consider the smoothness parabolic property for the  $\varphi(y)$  function:

$$\varphi(y) \leq \varphi(x) + \langle \nabla \varphi(x), y - x \rangle + \frac{L}{2} \|y - x\|_2^2$$
 
$$x := y, y := y - \frac{1}{L} \nabla \varphi(y) \quad \varphi\left(y - \frac{1}{L} \nabla \varphi(y)\right) \leq \varphi(y) + \left\langle \nabla \varphi(y), -\frac{1}{L} \nabla \varphi(y) \right\rangle + \frac{1}{2L} \|\nabla \varphi(y)\|_2^2$$
 
$$\varphi\left(y - \frac{1}{L} \nabla \varphi(y)\right) \leq \varphi(y) - \frac{1}{2L} \|\nabla \varphi(y)\|_2^2$$
 
$$\varphi\left(y - \frac{1}{L} \nabla \varphi(y)\right) \leq \varphi(y) - \frac{1}{2L} \|\nabla \varphi(y)\|_2^2$$

3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y=x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$



3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y=x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$



3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y=x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$


$$f(x) - \langle \nabla f(x), x \rangle \leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2I} \|\nabla f(y) - \nabla f(x)\|_2^2$$



3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y=x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$


$$\begin{split} f(x) - \langle \nabla f(x), x \rangle &\leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2L} \| \nabla f(y) - \nabla f(x) \|_2^2 \\ f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 &\leq f(y) \end{split}$$



3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y=x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$

$$\begin{split} &f(x) - \langle \nabla f(x), x \rangle \leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2L} \| \nabla f(y) - \nabla f(x) \|_2^2 \\ &f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 \leq f(y) \\ &\| \nabla f(y) - \nabla f(x) \|_2^2 \leq 2L \left( f(y) - f(x) - \langle \nabla f(x), y - x \rangle \right) \end{split}$$



3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y = x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$

4. Now, substitute  $\varphi(y) = f(y) - \langle \nabla f(x), y \rangle$ :

$$\begin{split} f(x) - \langle \nabla f(x), x \rangle &\leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2L} \| \nabla f(y) - \nabla f(x) \|_2^2 \\ f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 &\leq f(y) \\ \| \nabla f(y) - \nabla f(x) \|_2^2 &\leq 2L \left( f(y) - f(x) - \langle \nabla f(x), y - x \rangle \right) \end{split}$$
 switch x and y 
$$\| \nabla f(x) - \nabla f(y) \|_2^2 &\leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

Projected Gradient Descent (PGD)

3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y = x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$

4. Now, substitute  $\varphi(y) = f(y) - \langle \nabla f(x), y \rangle$ :

$$\begin{split} f(x) - \langle \nabla f(x), x \rangle &\leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2L} \| \nabla f(y) - \nabla f(x) \|_2^2 \\ f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 &\leq f(y) \\ \| \nabla f(y) - \nabla f(x) \|_2^2 &\leq 2L \left( f(y) - f(x) - \langle \nabla f(x), y - x \rangle \right) \end{split}$$
 switch x and y 
$$\| \nabla f(x) - \nabla f(y) \|_2^2 &\leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

Projected Gradient Descent (PGD)

3. From the first order optimality conditions for the convex function  $\nabla \varphi(y) = \nabla f(y) - \nabla f(x) = 0$ . We can conclude, that for any x, the minimum of the function  $\varphi(y)$  is at the point y = x. Therefore:

$$\varphi(x) \leq \varphi\left(y - \frac{1}{L}\nabla\varphi(y)\right) \leq \varphi(y) - \frac{1}{2L}\|\nabla\varphi(y)\|_2^2$$

4. Now, substitute  $\varphi(y) = f(y) - \langle \nabla f(x), y \rangle$ :

$$\begin{split} f(x) - \langle \nabla f(x), x \rangle &\leq f(y) - \langle \nabla f(x), y \rangle - \frac{1}{2L} \| \nabla f(y) - \nabla f(x) \|_2^2 \\ f(x) + \langle \nabla f(x), y - x \rangle + \frac{1}{2L} \| \nabla f(x) - \nabla f(y) \|_2^2 &\leq f(y) \\ \| \nabla f(y) - \nabla f(x) \|_2^2 &\leq 2L \left( f(y) - f(x) - \langle \nabla f(x), y - x \rangle \right) \\ \text{switch x and y} \quad \| \nabla f(x) - \nabla f(y) \|_2^2 &\leq 2L \left( f(x) - f(y) - \langle \nabla f(y), x - y \rangle \right) \end{split}$$

The lemma has been proved. From the first view it does not make a lot of geometrical sense, but we will use it as a convenient tool to bound the difference between gradients.

⊕ ი

#### i Theorem

Let  $f: \mathbb{R}^n \to \mathbb{R}$  be continuously differentiable on  $\mathbb{R}^n$ . Then, the function f is  $\mu$ -strongly convex if and only if for any  $x,y \in \mathbb{R}^d$  the following holds:

$$\begin{split} \text{Strongly convex case } \mu > 0 & \left\langle \nabla f(x) - \nabla f(y), x - y \right\rangle \geq \mu \|x - y\|^2 \\ \text{Convex case } \mu = 0 & \left\langle \nabla f(x) - \nabla f(y), x - y \right\rangle \geq 0 \end{split}$$

#### Proof

1. We will only give the proof for the strongly convex case, the convex one follows from it with setting  $\mu=0$ . We start from necessity. For the strongly convex function

$$\begin{split} f(y) &\geq f(x) + \langle \nabla f(x), y - x \rangle + \frac{\mu}{2} \|x - y\|_2^2 \\ f(x) &\geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{\mu}{2} \|x - y\|_2^2 \\ \text{sum } &\langle \nabla f(x) - \nabla f(y), x - y \rangle \geq \mu \|x - y\|^2 \end{split}$$

Projected Gradient Descent (PGD)

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x-y \rangle \geq \mu \|x-y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x-y \rangle \geq \mu \|x-y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$f(x) - f(y) - \langle \nabla f(y), x - y \rangle = \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$\begin{split} f(x) - f(y) - \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle \\ \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y), x - y \rangle dt \\ &= \int_0^1 \langle \nabla f(y + t(x - y)) - \nabla f(y), (x - y) \rangle dt \end{split}$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$\begin{split} f(x) - f(y) - \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle \\ \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y), x - y \rangle dt \\ &= \int_0^1 \langle \nabla f(y + t(x - y)) - \nabla f(y), (x - y) \rangle dt \\ &= \int_0^1 t^{-1} \langle \nabla f(y + t(x - y)) - \nabla f(y), t(x - y) \rangle dt \end{split}$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$\begin{split} f(x) - f(y) - \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle \\ \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y), x - y \rangle dt \\ &= \int_0^1 \langle \nabla f(y + t(x - y)) - \nabla f(y), (x - y) \rangle dt \\ &= \int_0^1 t^{-1} \langle \nabla f(y + t(x - y)) - \nabla f(y), t(x - y) \rangle dt \\ &\geq \int_0^1 t^{-1} \mu \|t(x - y)\|^2 dt \end{split}$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$\begin{split} f(x) - f(y) - \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle \\ \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y), x - y \rangle dt \\ &= \int_0^1 \langle \nabla f(y + t(x - y)) - \nabla f(y), (x - y) \rangle dt \\ &= \int_0^1 t^{-1} \langle \nabla f(y + t(x - y)) - \nabla f(y), t(x - y) \rangle dt \\ &\geq \int_0^1 t^{-1} \mu \|t(x - y)\|^2 dt = \mu \|x - y\|^2 \int_0^1 t dt \end{split}$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

Thus, we have a strong convexity criterion satisfied

$$f(x) \geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{\mu}{2} \|x - y\|_2^2$$

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

Thus, we have a strong convexity criterion satisfied

$$f(x) \geq f(y) + \langle \nabla f(y), x - y \rangle + \frac{\mu}{2} \|x - y\|_2^2$$
 or, equivivalently:

# 

2. For the sufficiency we assume, that  $\langle \nabla f(x) - \nabla f(y), x - y \rangle \ge \mu \|x - y\|^2$ . Using Newton-Leibniz theorem  $f(x) = f(y) + \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt$ :

$$\begin{split} f(x) - f(y) - \langle \nabla f(y), x - y \rangle &= \int_0^1 \langle \nabla f(y + t(x - y)), x - y \rangle dt - \langle \nabla f(y), x - y \rangle \\ &\langle \nabla f(y), x - y \rangle = \int_0^1 \langle \nabla f(y), x - y \rangle dt \\ &= \int_0^1 \langle \nabla f(y + t(x - y)) - \nabla f(y), (x - y) \rangle dt \\ &= \int_0^1 t^{-1} \langle \nabla f(y + t(x - y)) - \nabla f(y), t(x - y) \rangle dt \\ &\geq \int_0^1 t^{-1} \mu \|t(x - y)\|^2 dt = \mu \|x - y\|^2 \int_0^1 t dt = \frac{\mu}{2} \|x - y\|_2^2 \end{split}$$

Thus, we have a strong convexity criterion satisfied

$$f(x) \ge f(y) + \langle \nabla f(y), x - y \rangle + \frac{\mu}{2} \|x - y\|_2^2$$
 or, equivivalently:

switch x and y 
$$-\langle \nabla f(x), x-y \rangle \leq -\left(f(x)-f(y)+\frac{\mu}{2}\|x-y\|_2^2\right)$$



#### **i** Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$



#### **i** Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that  $y_k = x_k - \frac{1}{L}\nabla f(x_k)$  and cosine rule  $2x^T u = \|x\|^2 + \|u\|^2 - \|x - u\|^2$ :

#### i Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that  $y_k = x_k - \frac{1}{L}\nabla f(x_k)$  and cosine rule  $2x^T u = \|x\|^2 + \|u\|^2 - \|x - u\|^2$ :

$$\text{Smoothness:} \quad f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

#### **i** Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that  $y_k = x_k - \frac{1}{L}\nabla f(x_k)$  and cosine rule  $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$ :

Smoothness: 
$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

Method: 
$$= f(x_k) - L\langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

(7)

#### i Theorem

Let  $f:\mathbb{R}^n\to\mathbb{R}$  be convex and differentiable. Let  $S\subseteq\mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k > 0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that  $y_k = x_k - \frac{1}{L}\nabla f(x_k)$  and cosine rule  $2x^T y = ||x||^2 + ||y||^2 - ||x - y||^2$ :

Smoothness: 
$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

Method: 
$$= f(x_k) - L\langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

Cosine rule: 
$$= f(x_k) - \frac{L}{2} \left( \|y_k - x_k\|^2 + \|x_{k+1} - x_k\|^2 - \|y_k - x_{k+1}\|^2 \right) + \frac{L}{2} \|x_{k+1} - x_k\|^2$$

#### Convergence rate for smooth and convex case $\heartsuit$ $\heartsuit$

#### i Theorem

Let  $f:\mathbb{R}^n\to\mathbb{R}$  be convex and differentiable. Let  $S\subseteq\mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\frac{1}{L}$  achieves the following convergence after iteration k > 0:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that  $y_k = x_k - \frac{1}{L}\nabla f(x_k)$  and cosine rule  $2x^Ty = ||x||^2 + ||y||^2 - ||x - y||^2$ :

Smoothness: 
$$f(x, y) < f(x, y) + |\nabla f(x, y)| |x, y - x_x| + \frac{L}{2} ||x_x - x_y||^2$$

 $\text{Smoothness:} \quad f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2$ 

$$\begin{aligned} & \text{Method:} & = f(x_k) - L \langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2 \\ & \text{Cosine rule:} & = f(x_k) - \frac{L}{2} \left( \|y_k - x_k\|^2 + \|x_{k+1} - x_k\|^2 - \|y_k - x_{k+1}\|^2 \right) + \frac{L}{2} \|x_{k+1} - x_k\|^2 \end{aligned}$$

$$= f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2$$

(7)

Cosine rule:

Projected Gradient Descent (PGD)

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

3. We will use now projection property:  $\|x - \operatorname{proj}_S(y)\|^2 + \|y - \operatorname{proj}_S(y)\|^2 \le \|x - y\|^2$  with  $x = x^*, y = y_k$ :

$$\begin{split} \|x^* - \mathrm{proj}_S(y_k)\|^2 + \|y_k - \mathrm{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2 \end{split}$$

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

3. We will use now projection property:  $\|x - \operatorname{proj}_S(y)\|^2 + \|y - \operatorname{proj}_S(y)\|^2 \le \|x - y\|^2$  with  $x = x^*, y = y_k$ :

$$\begin{split} \|x^* - \mathrm{proj}_S(y_k)\|^2 + \|y_k - \mathrm{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2 \end{split}$$

4. Now, using convexity and previous part:

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

3. We will use now projection property:  $\|x - \operatorname{proj}_S(y)\|^2 + \|y - \operatorname{proj}_S(y)\|^2 \le \|x - y\|^2$  with  $x = x^*, y = y_k$ :

$$\begin{split} \|x^* - \mathrm{proj}_S(y_k)\|^2 + \|y_k - \mathrm{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2 \end{split}$$

4. Now, using convexity and previous part:

Convexity: 
$$f(x_k) - f^* \leq \langle \nabla f(x_k), x_k - x^* \rangle$$

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

3. We will use now projection property:  $\|x - \operatorname{proj}_S(y)\|^2 + \|y - \operatorname{proj}_S(y)\|^2 \le \|x - y\|^2$  with  $x = x^*, y = y_k$ :

$$\begin{split} \|x^* - \mathrm{proj}_S(y_k)\|^2 + \|y_k - \mathrm{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2 \end{split}$$

4. Now, using convexity and previous part:

$$\begin{split} \text{Convexity:} & f(x_k) - f^* \leq \langle \nabla f(x_k), x_k - x^* \rangle \\ & \leq \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_{k+1} - x^* \|^2 - \| y_k - x_{k+1} \|^2 \right) \end{split}$$

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{split} \left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_k - x^* - \frac{1}{L} \nabla f(x_k) \|^2 \right) \\ \left\langle \nabla f(x_k), x_k - x^* \right\rangle &= \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| y_k - x^* \|^2 \right) \end{split}$$

3. We will use now projection property:  $\|x - \operatorname{proj}_S(y)\|^2 + \|y - \operatorname{proj}_S(y)\|^2 \le \|x - y\|^2$  with  $x = x^*, y = y_k$ :

$$\begin{split} \|x^* - \mathrm{proj}_S(y_k)\|^2 + \|y_k - \mathrm{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2 \end{split}$$

4. Now, using convexity and previous part:

$$\begin{split} \text{Convexity:} & f(x_k) - f^* \leq \langle \nabla f(x_k), x_k - x^* \rangle \\ & \leq \frac{L}{2} \left( \frac{1}{L^2} \| \nabla f(x_k) \|^2 + \| x_k - x^* \|^2 - \| x_{k+1} - x^* \|^2 - \| y_k - x_{k+1} \|^2 \right) \end{split}$$

 $\text{Sum for } i = 0, k - 1 \quad \sum_{i = 0}^{k - 1} \left[ f(x_i) - f^* \right] \leq \sum_{i = 0}^{k - 1} \frac{1}{2L} \|\nabla f(x_i)\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i = 0}^{i - 1} \|y_i - x_{i + 1}\|^2$ 







#### 

$$\sum_{i=0}^{k-1} \left[ f(x_i) - f^* \right] \leq \sum_{i=0}^{k-1} \left[ f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2$$

### 

$$\begin{split} \sum_{i=0}^{k-1} \left[ f(x_i) - f^* \right] & \leq \sum_{i=0}^{k-1} \left[ f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \end{split}$$



$$\begin{split} \sum_{i=0}^{k-1} \left[ f(x_i) - f^* \right] & \leq \sum_{i=0}^{k-1} \left[ f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \end{split}$$



$$\begin{split} \sum_{i=0}^{k-1} \left[ f(x_i) - f^* \right] & \leq \sum_{i=0}^{k-1} \left[ f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ & \sum_{i=0}^{k-1} f(x_i) - kf^* \leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \end{split}$$

$$\begin{split} \sum_{i=0}^{k-1} \left[ f(x_i) - f^* \right] & \leq \sum_{i=0}^{k-1} \left[ f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ & \leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ & \leq \sum_{i=0}^{k-1} f(x_i) - kf^* \leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ & \sum_{i=0}^{k} \left[ f(x_i) - f^* \right] \leq \frac{L}{2} \|x_0 - x^*\|^2 \end{split}$$



6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$



6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$



6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that  $x_{k+1} = \operatorname{proj}_S(y_k)$ . By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$



6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that  $x_{k+1} = \operatorname{proj}_{S}(y_{k})$ . By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$

and recall that  $y_k = x_k - \frac{1}{L} \nabla f(x_k)$  implies  $||y_k - x_k|| = \frac{1}{L} ||\nabla f(x_k)||$ . Hence

$$\frac{L}{2} \, \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \, \|y_k - x_k\|^2 = \frac{L}{2} \, \frac{1}{L^2} \, \|\nabla f(x_k)\|^2 = \frac{1}{2L} \, \|\nabla f(x_k)\|^2.$$

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that  $x_{k+1} = \operatorname{proj}_{S}(y_{k})$ . By definition of projection,


$$||y_k - x_{k+1}|| \le ||y_k - x_k||,$$

and recall that  $y_k = x_k - \frac{1}{L} \nabla f(x_k)$  implies  $||y_k - x_k|| = \frac{1}{L} ||\nabla f(x_k)||$ . Hence

$$\frac{L}{2} \, \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \, \|y_k - x_k\|^2 = \frac{L}{2} \, \frac{1}{L^2} \, \|\nabla f(x_k)\|^2 = \frac{1}{2L} \, \|\nabla f(x_k)\|^2.$$

Substitute back into (\*):

$$f(x_{k+1}) \le f(x_k) - \frac{1}{2I} \|\nabla f(x_k)\|^2 + \frac{1}{2I} \|\nabla f(x_k)\|^2 = f(x_k).$$



6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that  $x_{k+1} = \operatorname{proj}_S(y_k)$ . By definition of projection,

$$||y_k - x_{k+1}|| \le ||y_k - x_k||,$$

and recall that  $y_k = x_k - \frac{1}{L} \nabla f(x_k)$  implies  $||y_k - x_k|| = \frac{1}{L} ||\nabla f(x_k)||$ . Hence

$$\frac{L}{2} \, \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \, \|y_k - x_k\|^2 = \frac{L}{2} \, \frac{1}{L^2} \, \|\nabla f(x_k)\|^2 = \frac{1}{2L} \, \|\nabla f(x_k)\|^2.$$

Substitute back into (\*):

$$f(x_{k+1}) \le f(x_k) - \frac{1}{2I} \|\nabla f(x_k)\|^2 + \frac{1}{2I} \|\nabla f(x_k)\|^2 = f(x_k).$$

Hence

$$f(x_{k+1}) \le f(x_k)$$
 for each  $k$ ,

so  $\{f(x_k)\}$  is a monotonically nonincreasing sequence.



7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} \big[ f(x_i) - f^* \big] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$



7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} \big[ f(x_i) - f^* \big] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$



7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Since  $f(x_i)$  decreases in i, in particular  $f(x_k) \leq f(x_i)$  for all  $i \leq k$ . Therefore

$$k\left[f(x_k) - f^*\right] \leq \sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2,$$



7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Since  $f(x_i)$  decreases in i, in particular  $f(x_k) \leq f(x_i)$  for all  $i \leq k$ . Therefore

$$k\left[f(x_k) - f^*\right] \leq \sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2,$$

which immediately gives

$$f(x_k) - f^* \le \frac{L\|x_0 - x^*\|_2^2}{2k}.$$

This completes the proof of the  $\mathcal{O}(\frac{1}{k})$  convergence rate for convex and L-smooth f under projection constraints.

#### i Theorem

Let  $f:\mathbb{R}^n\to\mathbb{R}$  be  $\mu$ -strongly convex. Let  $S\subseteq\mathbb{R}^n$ d be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Projected Gradient Descent algorithm with stepsize  $\alpha \leq \frac{1}{L}$  achieves the following convergence after iteration k > 0:

$$\|x_k - x^*\|_2^2 \leq \left(1 - \alpha \mu\right)^k \|x_0 - x^*\|_2^2$$

#### Proof

1. We first prove the stationary point property:  $\operatorname{proj}_{S}(x^* - \alpha \nabla f(x^*)) = x^*$ .

This follows from the projection criterion and the first-order optimality condition for  $x^*$ . Let  $y = x^* - \alpha \nabla f(x^*)$ . We need to show  $\langle y - x^*, x - x^* \rangle < 0$  for all  $x \in S$ .

$$\langle (x^* - \alpha \nabla f(x^*)) - x^*, x - x^* \rangle = -\alpha \langle \nabla f(x^*), x - x^* \rangle < 0$$

The inequality holds because  $\alpha > 0$  and  $\langle \nabla f(x^*), x - x^* \rangle \geq 0$  is the optimality condition for  $x^*$ .

#### Convergence rate for smooth strongly convex case 🔷 🎔

 $1. \ \,$  Considering the distance to the solution and using the stationary point property:



$$\|x_{k+1} - x^*\|_2^2 = \|\mathrm{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2$$



$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\text{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\text{proj}_S(x_k - \alpha \nabla f(x_k)) - \text{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \end{split}$$



$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\mathrm{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\mathrm{proj}_S(x_k - \alpha \nabla f(x_k)) - \mathrm{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \\ \text{nonexpansiveness} &\leq \|x_k - \alpha \nabla f(x_k) - (x^* - \alpha \nabla f(x^*))\|_2^2 \end{split}$$



$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - \operatorname{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \\ &\quad \operatorname{nonexpansiveness} \leq \|x_k - \alpha \nabla f(x_k) - (x^* - \alpha \nabla f(x^*))\|_2^2 \\ &= \|x_k - x^*\|^2 - 2\alpha \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle + \alpha^2 \|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \end{split}$$



1. Considering the distance to the solution and using the stationary point property:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - \operatorname{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \\ \text{nonexpansiveness} &\leq \|x_k - \alpha \nabla f(x_k) - (x^* - \alpha \nabla f(x^*))\|_2^2 \\ &= \|x_k - x^*\|^2 - 2\alpha \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle + \alpha^2 \|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \end{split}$$

2. Now we use smoothness from the convergence tools and strong convexity:



1. Considering the distance to the solution and using the stationary point property:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - \operatorname{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \\ & \operatorname{nonexpansiveness} \leq \|x_k - \alpha \nabla f(x_k) - (x^* - \alpha \nabla f(x^*))\|_2^2 \\ &= \|x_k - x^*\|^2 - 2\alpha \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle + \alpha^2 \|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \end{split}$$

2. Now we use smoothness from the convergence tools and strong convexity:

smoothness 
$$\|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \leq 2L\left(f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle\right)$$



1. Considering the distance to the solution and using the stationary point property:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - x^*\|_2^2 \\ \text{stationary point property} &= \|\operatorname{proj}_S(x_k - \alpha \nabla f(x_k)) - \operatorname{proj}_S(x^* - \alpha \nabla f(x^*))\|_2^2 \\ &\quad \operatorname{nonexpansiveness} \leq \|x_k - \alpha \nabla f(x_k) - (x^* - \alpha \nabla f(x^*))\|_2^2 \\ &= \|x_k - x^*\|^2 - 2\alpha \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle + \alpha^2 \|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \end{split}$$

2. Now we use smoothness from the convergence tools and strong convexity:

$$\begin{split} &\text{smoothness} \ \|\nabla f(x_k) - \nabla f(x^*)\|_2^2 \leq 2L\left(f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle\right) \\ &\text{strong convexity} \ - \langle \nabla f(x_k) - \nabla f(x^*), x_k - x^* \rangle \leq -\left(f(x_k) - f(x^*) + \frac{\mu}{2}\|x_k - x^*\|_2^2\right) - \langle \nabla f(x^*), x_k - x^* \rangle \end{split}$$



3. Substitute it:



3. Substitute it:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 & \leq \|x_k - x^*\|^2 - 2\alpha \left( f(x_k) - f(x^*) + \frac{\mu}{2} \|x_k - x^*\|_2^2 \right) - 2\alpha \langle \nabla f(x^*), x_k - x^* \rangle + \\ & + \alpha^2 2L \left( f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \right) \end{split}$$



#### 3. Substitute it:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 & \leq \|x_k - x^*\|^2 - 2\alpha \left( f(x_k) - f(x^*) + \frac{\mu}{2} \|x_k - x^*\|_2^2 \right) - 2\alpha \langle \nabla f(x^*), x_k - x^* \rangle + \\ & + \alpha^2 2L \left( f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \right) \\ & \leq (1 - \alpha \mu) \|x_k - x^*\|^2 + 2\alpha (\alpha L - 1) \left( f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \right) \end{split}$$



3. Substitute it:

$$\begin{split} \|x_{k+1} - x^*\|_2^2 & \leq \|x_k - x^*\|^2 - 2\alpha \left( f(x_k) - f(x^*) + \frac{\mu}{2} \|x_k - x^*\|_2^2 \right) - 2\alpha \langle \nabla f(x^*), x_k - x^* \rangle + \\ & + \alpha^2 2L \left( f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \right) \\ & \leq (1 - \alpha \mu) \|x_k - x^*\|^2 + 2\alpha (\alpha L - 1) \left( f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \right) \end{split}$$

4. Due to convexity of  $f: f(x_k) - f(x^*) - \langle \nabla f(x^*), x_k - x^* \rangle \ge 0$ . Therefore, if we use  $\alpha \le \frac{1}{L}$ :

$$\|x_{k+1}-x^*\|_2^2 \leq (1-\alpha\mu)\|x_k-x^*\|^2,$$

which is exactly linear convergence of the method with up to  $1-\frac{\mu}{L}$  convergence rate.

 $f \to \min_{x,y,z}$  Projected Gradient Descent (PGD)

#### Frank-Wolfe Method







Рисунок 11: Marguerite Straus Frank (1927-2024)



Рисунок 12: Philip Wolfe (1927-2016)








Рисунок 13: Illustration of Frank-Wolfe (conditional gradient) algorithm

**♥೧**0




Рисунок 14: Illustration of Frank-Wolfe (conditional gradient) algorithm

**♥ ೧** 0




Рисунок 15: Illustration of Frank-Wolfe (conditional gradient) algorithm

⊕ ∩ ∅




Рисунок 16: Illustration of Frank-Wolfe (conditional gradient) algorithm

**⊕** ೧ **0** 




Рисунок 17: Illustration of Frank-Wolfe (conditional gradient) algorithm

**⊕** ೧ **ø** 




Рисунок 18: Illustration of Frank-Wolfe (conditional gradient) algorithm

**♥೧**0





Рисунок 19: Illustration of Frank-Wolfe (conditional gradient) algorithm

**⊕ດ** Ø

$$\begin{split} y_k &= \arg\min_{x \in S} f^I_{x_k}(x) = \arg\min_{x \in S} \langle \nabla f(x_k), x \rangle \\ x_{k+1} &= \gamma_k x_k + (1 - \gamma_k) y_k \end{split}$$





# Convergence rate for smooth and convex case $\heartsuit$ $\heartsuit$

#### i Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$  be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Frank-Wolfe algorithm with step size  $\gamma_k = \frac{k-1}{k+1}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \leq \frac{2LR^2}{k+1}$$

where  $R = \max_{x,y \in S} \|x-y\|$  is the diameter of the set S.

Frank-Wolfe Method

#### **i** Theorem

Let  $f:\mathbb{R}^n \to \mathbb{R}$  be convex and differentiable. Let  $S\subseteq \mathbb{R}^n$  be a closed convex set, and assume that there is a minimizer  $x^*$  of f over S; furthermore, suppose that f is smooth over S with parameter L. The Frank-Wolfe algorithm with step size  $\gamma_k = \frac{k-1}{k+1}$  achieves the following convergence after iteration k>0:

$$f(x_k) - f^* \le \frac{2LR^2}{k+1}$$

where  $R = \max_{x,y \in S} \|x - y\|$  is the diameter of the set S.

1. By L-smoothness of f, we have:

$$\begin{split} f\left(x_{k+1}\right) - f\left(x_{k}\right) &\leq \left\langle \nabla f\left(x_{k}\right), x_{k+1} - x_{k}\right\rangle + \frac{L}{2}\left\|x_{k+1} - x_{k}\right\|^{2} \\ &= \left(1 - \gamma_{k}\right) \left\langle \nabla f\left(x_{k}\right), y_{k} - x_{k}\right\rangle + \frac{L(1 - \gamma_{k})^{2}}{2}\left\|y_{k} - x_{k}\right\|^{2} \end{split}$$



2. By convexity of f, for any  $x \in S$ , including  $x^*$ :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for  $x = x^*$ :

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

 $f \to \min_{x,y,z}$  Frank-Wolfe Method

2. By convexity of f, for any  $x \in S$ , including  $x^*$ :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for  $x = x^*$ :

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of  $y_k$ , we have  $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$ , thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \leq \langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

2. By convexity of f, for any  $x \in S$ , including  $x^*$ :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for  $x = x^*$ :

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of  $y_k$ , we have  $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$ , thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \leq \langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

4. Combining the above inequalities:

$$\begin{split} f\left(x_{k+1}\right) - f\left(x_{k}\right) &\leq \left(1 - \gamma_{k}\right) \left\langle \nabla f\left(x_{k}\right), y_{k} - x_{k}\right\rangle + \frac{L(1 - \gamma_{k})^{2}}{2} \left\|y_{k} - x_{k}\right\|^{2} \\ &\leq \left(1 - \gamma_{k}\right) \left(f(x^{*}) - f(x_{k})\right) + \frac{L(1 - \gamma_{k})^{2}}{2} R^{2} \end{split}$$

2. By convexity of f, for any  $x \in S$ , including  $x^*$ :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for  $x = x^*$ :

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of  $y_k$ , we have  $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$ , thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \le \langle \nabla f(x_k), x^* - x_k \rangle \le f(x^*) - f(x_k)$$

4. Combining the above inequalities:

$$\begin{split} f\left(x_{k+1}\right) - f\left(x_{k}\right) &\leq \left(1 - \gamma_{k}\right) \left\langle \nabla f\left(x_{k}\right), y_{k} - x_{k}\right\rangle + \frac{L(1 - \gamma_{k})^{2}}{2} \left\|y_{k} - x_{k}\right\|^{2} \\ &\leq \left(1 - \gamma_{k}\right) \left(f(x^{*}) - f(x_{k})\right) + \frac{L(1 - \gamma_{k})^{2}}{2} R^{2} \end{split}$$

5. Rearranging terms:

$$f\left(x_{k+1}\right) - f(x^*) \leq \gamma_k \left(f(x_k) - f(x^*)\right) + (1 - \gamma_k)^2 \frac{LR^2}{2}$$

6. Denoting 
$$\delta_k = \frac{f(x_k) - f(x^*)}{IP^2}$$
, we get:

$$\delta_{k+1} \le \gamma_k \delta_k + \frac{(1 - \gamma_k)^2}{2} = \frac{k - 1}{k + 1} \delta_k + \frac{2}{(k + 1)^2}$$



 $f \to \min_{x,y,z}$  Frank-Wolfe Method



6. Denoting  $\delta_k = \frac{f(x_k) - f(x^*)}{I \cdot D^2}$ , we get:

$$\delta_{k+1} \le \gamma_k \delta_k + \frac{(1 - \gamma_k)^2}{2} = \frac{k - 1}{k + 1} \delta_k + \frac{2}{(k + 1)^2}$$

7. We will prove that  $\delta_k \leq \frac{2}{k+1}$  by induction.

which gives us the desired result:

$$f(x_k) - f^* \le \frac{2LR^2}{k+1}$$

 $f \to \min_{x,y,z}$  Frank-Wolfe Method



6. Denoting  $\delta_k = \frac{f(x_k) - f(x^*)}{I \cdot D^2}$ , we get:

$$\delta_{k+1} \le \gamma_k \delta_k + \frac{(1 - \gamma_k)^2}{2} = \frac{k - 1}{k + 1} \delta_k + \frac{2}{(k + 1)^2}$$

- 7. We will prove that  $\delta_k \leq \frac{2}{k+1}$  by induction.
  - Base:  $\delta_2 \le \frac{1}{2} < \frac{2}{2}$

which gives us the desired result:

$$f(x_k) - f^* \le \frac{2LR^2}{k+1}$$



6. Denoting  $\delta_k = \frac{f(x_k) - f(x^*)}{I \cdot D^2}$ , we get:

$$\delta_{k+1} \leq \gamma_k \delta_k + \frac{(1-\gamma_k)^2}{2} = \frac{k-1}{k+1} \delta_k + \frac{2}{(k+1)^2}$$

- 7. We will prove that  $\delta_k \leq \frac{2}{k+1}$  by induction.
  - Base:  $\delta_2 \le \frac{1}{2} < \frac{2}{3}$
  - Assume  $\delta_k \leq \frac{2}{k+1}$

which gives us the desired result:

$$f(x_k) - f^* \le \frac{2LR^2}{k+1}$$



6. Denoting  $\delta_k = \frac{f(x_k) - f(x^*)}{I R^2}$ , we get:

$$\delta_{k+1} \le \gamma_k \delta_k + \frac{(1-\gamma_k)^2}{2} = \frac{k-1}{k+1} \delta_k + \frac{2}{(k+1)^2}$$

- 7. We will prove that  $\delta_k \leq \frac{2}{k+1}$  by induction.
  - Base:  $\delta_2 \le \frac{1}{2} < \frac{2}{3}$
  - Assume  $\delta_k \leq \frac{2}{k+1}$
  - Then  $\delta_{k+1} \leq \frac{k-1}{k+1} \cdot \frac{2}{k+1} + \frac{2}{(k+1)^2} = \frac{2k}{k^2+2k+1} < \frac{2}{k+2}$

which gives us the desired result:

$$f(x_k) - f^* \le \frac{2LR^2}{k+1}$$

 $f \to \min_{x,y,z}$  Frank-Wolfe Method

### Lower bound for Frank-Wolfe method <sup>2</sup>

#### **i** Theorem

Consider any algorithm that accesses the feasible set  $S \subseteq \mathbb{R}^n$  only via a linear minimization oracle (LMO). Let the diameter of the set S be R. There exists an L-smooth strongly convex function  $f: \mathbb{R}^n \to \mathbb{R}$  such that this algorithm requires at least

$$\min\left(\frac{n}{2}, \frac{LR^2}{16\varepsilon}\right)$$

iterations (i.e., calls to the LMO) to construct a point  $\hat{x} \in S$  with  $f(\hat{x}) - \min_{x \in S} f(x) \le \varepsilon$ . The lower bound applies both for convex and strongly convex functions.

## Lower bound for Frank-Wolfe method <sup>2</sup>

#### i Theorem

Consider any algorithm that accesses the feasible set  $S\subseteq\mathbb{R}^n$  only via a linear minimization oracle (LMO). Let the diameter of the set S be R. There exists an L-smooth strongly convex function  $f:\mathbb{R}^n\to\mathbb{R}$  such that this algorithm requires at least

$$\min\left(\frac{n}{2}, \frac{LR^2}{16\varepsilon}\right)$$

iterations (i.e., calls to the LMO) to construct a point  $\hat{x} \in S$  with  $f(\hat{x}) - \min_{x \in S} f(x) \le \varepsilon$ . The lower bound applies both for convex and strongly convex functions.

**Sketch of the proof.** Consider the following optimization problem:

$$\min_{x \in S} f(x) = \min_{x \in S} \frac{1}{2} ||x||_2^2$$

Frank-Wolfe Method

$$S = \left\{ x \in \mathbb{R}^n \mid x \ge 0, \ \sum_{i=1}^n x_i = 1 \right\}$$

Note, that:

- *f* is 1-smooth;
- the diameter of S is R=2:
- f is strongly convex.

<sup>2</sup> The Complexity of Large-scale Convex Programming under a Linear Optimization Oracle

### 

1. The optimal solution is

$$x^* = \frac{1}{n} \mathbf{1} = \frac{1}{n} \sum_{i=1}^n e_i, \quad \text{and} \quad f(x^*) = \frac{1}{2n},$$

where  $e_i = (0,\dots,0,\underbrace{1}_{\text{position }i},0,\dots,0)^{\top}$  is the i-th standard basis vector.

### Lower bound for Frank-Wolfe method <sup>3</sup> \*

1. The optimal solution is

$$x^* = \frac{1}{n} \mathbf{1} = \frac{1}{n} \sum_{i=1}^n e_i, \quad \text{and} \quad f(x^*) = \frac{1}{2n},$$

where  $e_i = (0, \dots, 0, \frac{1}{\text{nosition } i}, 0, \dots, 0)^{\top}$  is the *i*-th standard basis vector.

2. A linear minimization oracle (LMO) over S returns a vertex  $e_i$ . After k iterations, the method will have discovered at most k different basis vectors  $e_{i_1}, \dots, e_{i_k}$ . The best convex combination one can form is

$$\hat{x} = \frac{1}{k} \sum_{i=1}^{k} e_{i_j}.$$

## Lower bound for Frank-Wolfe method <sup>3</sup> 🔷

1. The optimal solution is

$$x^* = \frac{1}{n}\mathbf{1} = \frac{1}{n}\sum_{i=1}^n e_i, \quad \text{and} \quad f(x^*) = \frac{1}{2n},$$

where  $e_i = (0,\dots,0,\underbrace{1}_{\text{position }i},0,\dots,0)^{\top}$  is the i-th standard basis vector.

2. A linear minimization oracle (LMO) over S returns a vertex  $e_i$ . After k iterations, the method will have discovered at most k different basis vectors  $e_{i_1}, \dots, e_{i_k}$ . The best convex combination one can form is

$$\hat{x} = \frac{1}{k} \sum_{i=1}^{k} e_{i_j}.$$

3. Evaluating the function at  $\hat{x}$ , we obtain:

$$f(\hat{x}) - f(x^*) \ge \frac{1}{2} \left( \frac{1}{\min\{k, n\}} - \frac{1}{n} \right).$$

Frank-Wolfe Method

# Lower bound for Frank-Wolfe method <sup>3</sup>

1. The optimal solution is

$$x^* = \frac{1}{n} \mathbf{1} = \frac{1}{n} \sum_{i=1}^n e_i, \quad \text{and} \quad f(x^*) = \frac{1}{2n},$$

where  $e_i = (0, \dots, 0, \underbrace{1}_{\text{position } i}, 0, \dots, 0)^{\top}$  is the i-th standard basis vector.

2. A linear minimization oracle (LMO) over S returns a vertex  $e_i$ . After k iterations, the method will have discovered at most k different basis vectors  $e_{i_1}, \dots, e_{i_k}$ . The best convex combination one can form is

$$\hat{x} = \frac{1}{k} \sum_{i=1}^{k} e_{i_j}.$$

Evaluating the function at  $\hat{x}$ , we obtain:

$$f(\hat{x}) - f(x^*) \ge \frac{1}{2} \left( \frac{1}{\min\{k, n\}} - \frac{1}{n} \right).$$

4. To ensure that  $f(\hat{x}) - f(x^*) \le \varepsilon$ , it is necessary that (full proof is in the paper):

$$k \ge \min\left\{\frac{n}{2}, \frac{1}{4\varepsilon}\right\} = \min\left\{\frac{n}{2}, \frac{LR^2}{16\varepsilon}\right\}.$$

<sup>&</sup>lt;sup>3</sup> The Complexity of Large-scale Convex Programming under a Linear Optimization Oracle

• Method does not require projections, in some special cases allows to compute iterations in closed form

Frank-Wolfe Method

⊕ ი

- Method does not require projections, in some special cases allows to compute iterations in closed form
- Global convergence rate is  $O(\frac{1}{k})$  for smooth and convex functions. Strong convexity does not improve the rate. This is the lower bound for LMO

- Method does not require projections, in some special cases allows to compute iterations in closed form
- Global convergence rate is  $O\left(\frac{1}{k}\right)$  for smooth and convex functions. Strong convexity does not improve the rate. This is the lower bound for LMO
- In comparison with projected gradient descent, the rate is worse, but iteration could be cheaper and more sparse

⊕ ი

- Method does not require projections, in some special cases allows to compute iterations in closed form
- Global convergence rate is  $O\left(\frac{1}{L}\right)$  for smooth and convex functions. Strong convexity does not improve the rate. This is the lower bound for LMO
- In comparison with projected gradient descent, the rate is worse, but iteration could be cheaper and more sparse
- Recently, it was shown that for strongly convex sets, the rate can be improved to  $O(\frac{1}{12})$  ( $\mathbb{Z}$  paper)



- Method does not require projections, in some special cases allows to compute iterations in closed form
- Global convergence rate is  $O\left(\frac{1}{k}\right)$  for smooth and convex functions. Strong convexity does not improve the rate. This is the lower bound for LMO
- In comparison with projected gradient descent, the rate is worse, but iteration could be cheaper and more sparse
- Recently, it was shown that for strongly convex sets, the rate can be improved to  $O\left(\frac{1}{k^2}\right)$  (**b** paper)
- If we allow away steps, the convergence becomes linear ( paper) in strongly convex case





- Method does not require projections, in some special cases allows to compute iterations in closed form
- Global convergence rate is  $O(\frac{1}{k})$  for smooth and convex functions. Strong convexity does not improve the rate. This is the lower bound for LMO
- In comparison with projected gradient descent, the rate is worse, but iteration could be cheaper and more sparse
- Recently, it was shown that for strongly convex sets, the rate can be improved to  $O\left(\frac{1}{L^2}\right)$  ( paper)
- If we allow away steps, the convergence becomes linear ( paper ) in strongly convex case
- Recent work showed the extension to non-smooth case ( paper) with convergence rate  $O\left(\frac{1}{\sqrt{k}}\right)$


## **Numerical experiments**





























## 2d example. Projected gradient descent



#### 2d example. Projected gradient descent



#### 2d example. Projected gradient descent



#### Quadratic function. Box constraints

$$\min_{\substack{x \in \mathbb{R}^n \\ -1 \le x \le 1}} \frac{1}{2} x^{\top} A x - b^{\top} x,$$

$$A \in \mathbb{R}^{n \times n}, \quad \lambda(A) \in [\mu; L].$$

The projection is simple:

$$\pi_S(x) = \mathsf{clip}(x, -\mathbf{1}, \mathbf{1}).$$

or


$$\pi_S(x) = \max\left(-\mathbf{1}, \min(\mathbf{1}, x)\right).$$

The linear minimization oracle (LMO) for a given gradient g is given by  $y = \underset{z \in S}{\operatorname{argmin}} \langle g, z \rangle.$ 

Since the feasible set is separable across coordinates, the solution is computed coordinate—wise as

$$y_i = \begin{cases} -1, & \text{if } g_i > 0, \\ 1, & \text{if } g_i \leq 0. \end{cases}$$

Constrained convex quadratic problem: n=80,  $\mu$ =0, L=10



1.0

Projected Gradient Descent

Time (seconds)

1e-3

0.0 0.2 0.4 0.6 0.8 1.0

0.0 0.2 0.4 0.6

 $f \to \min_{x,y,z}$  Num

Numerical experiments

Time (seconds)

Frank-Wolfe

1e-3

#### Quadratic function. Box constraints

$$\min_{\substack{x \in \mathbb{R}^n \\ -1 \le x \le 1}} \frac{1}{2} x^{\top} A x - b^{\top} x,$$

$$A\in\mathbb{R}^{n\times n},\quad\lambda\left(A\right)\in\left[\mu;L\right].$$

The projection is simple:

$$\pi_S(x) = \mathsf{clip}(x, -\mathbf{1}, \mathbf{1}).$$

or


$$\pi_S(x) = \max\left(-\mathbf{1}, \min(\mathbf{1}, x)\right).$$


The linear minimization oracle (LMO) for a given gradient g is given by  $y = \operatorname{argmin} \langle g, z \rangle$ . Since the feasible set is separable across

coordinates, the solution is computed

$$y_i = \begin{cases} -1, & \text{if } g_i > 0, \\ 1, & \text{if } g_i \leq 0. \end{cases}$$

Constrained strongly Convex quadratic problem: n=80,  $\mu=1$ , L=10





1.00

1e-3

0.50 0.75

Time (seconds)

coordinate-wise as

## Quadratic function. Simplex constraints (Lucky problem with diagonal matrix)

$$\min_{\substack{x \in \mathbb{R}^n \\ x \ge 0, \mathbf{1}^T x = 1}} \frac{1}{2} x^T A x,$$


$$^{\times n}, \quad \lambda(A) \in [0; 100].$$


The projection onto the unit simplex  $\pi_S(x)$  can be done in  $\mathcal{O}(n\log n)$  or expected  $\mathcal{O}(n)$  time. <sup>4</sup> The LMO for a given gradient g is given by  $y = \operatorname{argmin}\langle g, z \rangle$ . The solution corresponds to

 $z \in S$ 

 $A \in \mathbb{R}^{n \times n}$ ,

 $y = e_j$  where  $j = \underset{i}{\operatorname{argmin}} g_i$ .










#### Quadratic function. Simplex constraints

$$\min_{\substack{x \in \mathbb{R}^n \\ x \ge 0, \mathbf{1}^T x = 1}} \frac{1}{2} x^T A x,$$

$$A \in \mathbb{R}^{n \times n}, \quad \lambda\left(A\right) \in [0; 100].$$











#### Quadratic function. Simplex constraints

$$\min_{\substack{x \in \mathbb{R}^n \\ x \ge 0, \mathbf{1}^T x = 1}} \frac{1}{2} x^T A x,$$

$$A \in \mathbb{R}^{n \times n}, \quad \lambda\left(A\right) \in [0; 100].$$

# $\min_{1^T x = 1, x \ge 0} 1/2x^T A x, n = 300$









#### Quadratic function. Simplex constraints

$$\min_{\substack{x \in \mathbb{R}^n \\ x \ge 0, \mathbf{1}^T x = 1}} \frac{1}{2} x^T A x,$$

 $A \in \mathbb{R}^{n \times n}, \quad \lambda(A) \in [1; 100].$ 





Projected Gradient Descent

--- Frank-Wolfe



## PGD vs Frank-Wolfe

The key difference between PGD and FW is that PGD requires projection, while FW needs only linear minimization oracle (LMO).

In a recent book authors presented the following comparison table with complexities of linear minimizations and projections on some convex sets up to an additive error  $\epsilon$  in the Euclidean norm.

Set	Linear minimization	Projection
	$\mathcal{O}(n)$	$\tilde{\mathcal{O}}\!\!\left(rac{n}{\epsilon^2} ight)$
Nuclear norm ball of $n \times m$ matrices	$\mathcal{O}\left(\nu \ln(m+n) \frac{\sqrt{\sigma_1}}{\sqrt{\epsilon}}\right)$ $\mathcal{O}\left((n\log m)(n+m\log m)\right)$	$\mathcal{O}\!\!\left(mn\min\{m,n\}\right)$
Flow polytope on a graph with $m$ vertices and $n$ edges (capacity bound on edges)	$\mathcal{O}\!\!\left((n\log m)\big(n+m\log m\big)\right)$	$\tilde{\mathcal{O}}\!\!\left(\frac{n}{\epsilon^2}\right)$ or $\mathcal{O}(n^4\log n)$
Birkhoff polytope ( $n \times n$ doubly stochastic	$\mathcal{O}(n^3)$	$ ilde{\mathcal{O}}\!\!\left(rac{n^2}{\epsilon^2} ight)$
matrices)		-

When  $\epsilon$  is missing, there is no additive error. The  $\widetilde{\mathcal{O}}$  hides polylogarithmic factors in the dimensions and polynomial factors in constants related to the distance to the optimum. For the nuclear norm ball, i.e., the spectrahedron,  $\nu$  denotes the number of non-zero entries and  $\sigma_1$  denotes the top singular value of the projected matrix.