




Gradient methods for conditional problems.
Projected Gradient Descent. Frank-Wolfe
method. Idea of Mirror Descent algorithm

Даня Меркулов

Методы Оптимизации в Машинном Обучении. ФКН ВШЭ

Conditional methods

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.
- The solution has to be inside the set S .

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.
- The solution has to be inside the set S .
- Example:

$$\frac{1}{2} \|Ax - b\|_2^2 \rightarrow \min_{\|x\|_2^2 \leq 1}$$

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.
- The solution has to be inside the set S .
- Example:

$$\frac{1}{2} \|Ax - b\|_2^2 \rightarrow \min_{\|x\|_2^2 \leq 1}$$

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.
- The solution has to be inside the set S .
- Example:

$$\frac{1}{2} \|Ax - b\|_2^2 \rightarrow \min_{\|x\|_2^2 \leq 1}$$

Gradient Descent is a great way to solve unconstrained problem

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k) \quad (\text{GD})$$

Is it possible to tune GD to fit constrained problem?

Constrained optimization

Unconstrained optimization

$$\min_{x \in \mathbb{R}^n} f(x)$$

- Any point $x_0 \in \mathbb{R}^n$ is feasible and could be a solution.

Constrained optimization

$$\min_{x \in S} f(x)$$

- Not all $x \in \mathbb{R}^n$ are feasible and could be a solution.
- The solution has to be inside the set S .
- Example:

$$\frac{1}{2} \|Ax - b\|_2^2 \rightarrow \min_{\|x\|_2^2 \leq 1}$$

Gradient Descent is a great way to solve unconstrained problem

$$x_{k+1} = x_k - \alpha_k \nabla f(x_k) \quad (\text{GD})$$

Is it possible to tune GD to fit constrained problem?

Yes. We need to use projections to ensure feasibility on every iteration.

Example: White-box Adversarial Attacks




Figure 1: Source

- Mathematically, a neural network is a function $f(w; x)$

Example: White-box Adversarial Attacks





Figure 1: Source

- Mathematically, a neural network is a function $f(w; x)$
- Typically, input x is given and network weights w optimized

Example: White-box Adversarial Attacks



- Mathematically, a neural network is a function $f(w; x)$
- Typically, input x is given and network weights w optimized
- Could also freeze weights w and optimize x , adversarially!

$$\min_{\delta} \text{size}(\delta) \quad \text{s.t.} \quad \text{pred}[f(w; x + \delta)] \neq y$$

or

$$\max_{\delta} l(w; x + \delta, y) \quad \text{s.t.} \quad \text{size}(\delta) \leq \epsilon, \quad 0 \leq x + \delta \leq 1$$

Idea of Projected Gradient Descent




Figure 2: Suppose, we start from a point x_k .

Idea of Projected Gradient Descent




Figure 3: And go in the direction of $-\nabla f(x_k)$.

Idea of Projected Gradient Descent




Figure 4: Occasionally, we can end up outside the feasible set.

Idea of Projected Gradient Descent




Figure 5: Solve this little problem with projection!

Idea of Projected Gradient Descent

$$x_{k+1} = \text{proj}_S(x_k - \alpha_k \nabla f(x_k)) \quad \Leftrightarrow \quad \begin{aligned} y_k &= x_k - \alpha_k \nabla f(x_k) \\ x_{k+1} &= \text{proj}_S(y_k) \end{aligned}$$




Figure 6: Illustration of Projected Gradient Descent algorithm

Projection

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point $\mathbf{y} \in \mathbb{R}^n$ on set $S \subseteq \mathbb{R}^n$ is a point $\text{proj}_S(\mathbf{y}) \in S$:

$$\text{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|_2^2$$

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point $\mathbf{y} \in \mathbb{R}^n$ on set $S \subseteq \mathbb{R}^n$ is a point $\text{proj}_S(\mathbf{y}) \in S$:

$$\text{proj}_S(\mathbf{y}) = \underset{\mathbf{x} \in S}{\operatorname{argmin}} \frac{1}{2} \|\mathbf{x} - \mathbf{y}\|_2^2$$

- **Sufficient conditions of existence of a projection.** If $S \subseteq \mathbb{R}^n$ - closed set, then the projection on set S exists for any point.

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point $\mathbf{y} \in \mathbb{R}^n$ on set $S \subseteq \mathbb{R}^n$ is a point $\text{proj}_S(\mathbf{y}) \in S$:

$$\text{proj}_S(\mathbf{y}) = \underset{x \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- **Sufficient conditions of existence of a projection.** If $S \subseteq \mathbb{R}^n$ - closed set, then the projection on set S exists for any point.
- **Sufficient conditions of uniqueness of a projection.** If $S \subseteq \mathbb{R}^n$ - closed convex set, then the projection on set S is unique for any point.

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point $\mathbf{y} \in \mathbb{R}^n$ on set $S \subseteq \mathbb{R}^n$ is a point $\text{proj}_S(\mathbf{y}) \in S$:

$$\text{proj}_S(\mathbf{y}) = \underset{x \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- **Sufficient conditions of existence of a projection.** If $S \subseteq \mathbb{R}^n$ - closed set, then the projection on set S exists for any point.
- **Sufficient conditions of uniqueness of a projection.** If $S \subseteq \mathbb{R}^n$ - closed convex set, then the projection on set S is unique for any point.
- If a set is open, and a point is beyond this set, then its projection on this set may not exist.

Projection

The distance d from point $\mathbf{y} \in \mathbb{R}^n$ to closed set $S \subset \mathbb{R}^n$:

$$d(\mathbf{y}, S, \|\cdot\|) = \inf\{\|x - y\| \mid x \in S\}$$

We will focus on Euclidean projection (other options are possible) of a point $\mathbf{y} \in \mathbb{R}^n$ on set $S \subseteq \mathbb{R}^n$ is a point $\text{proj}_S(\mathbf{y}) \in S$:

$$\text{proj}_S(\mathbf{y}) = \underset{x \in S}{\operatorname{argmin}} \frac{1}{2} \|x - y\|_2^2$$

- **Sufficient conditions of existence of a projection.** If $S \subseteq \mathbb{R}^n$ - closed set, then the projection on set S exists for any point.
- **Sufficient conditions of uniqueness of a projection.** If $S \subseteq \mathbb{R}^n$ - closed convex set, then the projection on set S is unique for any point.
- If a set is open, and a point is beyond this set, then its projection on this set may not exist.
- If a point is in set, then its projection is the point itself.

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$

$$2 (\text{proj}_S(y) - y)^T (x - \text{proj}_S(y)) \geq 0$$




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$

$$2 (\text{proj}_S(y) - y)^T (x - \text{proj}_S(y)) \geq 0$$

$$(y - \text{proj}_S(y))^T (x - \text{proj}_S(y)) \leq 0$$




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$

$$2 (\text{proj}_S(y) - y)^T (x - \text{proj}_S(y)) \geq 0$$

$$(y - \text{proj}_S(y))^T (x - \text{proj}_S(y)) \leq 0$$

2. Use cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$ with $x = x - \text{proj}_S(y)$ and $y = y - \text{proj}_S(y)$. By the first property of the theorem:




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$

$$2 (\text{proj}_S(y) - y)^T (x - \text{proj}_S(y)) \geq 0$$

$$(y - \text{proj}_S(y))^T (x - \text{proj}_S(y)) \leq 0$$

2. Use cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$ with $x = x - \text{proj}_S(y)$ and $y = y - \text{proj}_S(y)$. By the first property of the theorem:

$$0 \geq 2x^T y = \|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 - \|x - y\|^2$$




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection criterion (Bourbaki-Cheney-Goldstein inequality)

i Theorem

Let $S \subseteq \mathbb{R}^n$ be closed and convex, $\forall x \in S, y \in \mathbb{R}^n$. Then

$$\langle y - \text{proj}_S(y), x - \text{proj}_S(y) \rangle \leq 0 \quad (1)$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2 \quad (2)$$

1. $\text{proj}_S(y)$ is minimizer of differentiable convex function $d(y, S, \|\cdot\|) = \|x - y\|^2$ over S . By first-order characterization of optimality.

$$\nabla d(\text{proj}_S(y))^T (x - \text{proj}_S(y)) \geq 0$$

$$2 (\text{proj}_S(y) - y)^T (x - \text{proj}_S(y)) \geq 0$$

$$(y - \text{proj}_S(y))^T (x - \text{proj}_S(y)) \leq 0$$

2. Use cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$ with $x = x - \text{proj}_S(y)$ and $y = y - \text{proj}_S(y)$. By the first property of the theorem:

$$0 \geq 2x^T y = \|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 - \|x - y\|^2$$

$$\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$$




Figure 7: Obtuse or straight angle should be for any point $x \in S$

Projection operator is non-expansive

- A function f is called non-expansive if f is L -Lipschitz with $L \leq 1$ ¹. That is, for any two points $x, y \in \text{dom} f$,

$$\|f(x) - f(y)\| \leq L\|x - y\|, \text{ where } L \leq 1.$$

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

¹Non-expansive becomes contractive if $L < 1$.

Projection operator is non-expansive

- A function f is called non-expansive if f is L -Lipschitz with $L \leq 1$ ¹. That is, for any two points $x, y \in \text{dom} f$,

$$\|f(x) - f(y)\| \leq L\|x - y\|, \text{ where } L \leq 1.$$

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

- Projection operator is non-expansive:

$$\|\text{proj}(x) - \text{proj}(y)\|_2 \leq \|x - y\|_2.$$

¹Non-expansive becomes contractive if $L < 1$.

Projection operator is non-expansive

- A function f is called non-expansive if f is L -Lipschitz with $L \leq 1$ ¹. That is, for any two points $x, y \in \text{dom} f$,

$$\|f(x) - f(y)\| \leq L\|x - y\|, \text{ where } L \leq 1.$$

It means the distance between the mapped points is possibly smaller than that of the unmapped points.

- Projection operator is non-expansive:

$$\|\text{proj}(x) - \text{proj}(y)\|_2 \leq \|x - y\|_2.$$

- Next: variational characterization implies non-expansiveness. i.e.,

$$\langle y - \text{proj}(y), x - \text{proj}(y) \rangle \leq 0 \quad \forall x \in S \quad \Rightarrow \quad \|\text{proj}(x) - \text{proj}(y)\|_2 \leq \|x - y\|_2.$$

¹Non-expansive becomes contractive if $L < 1$.

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Replace x by $\pi(x)$ in Equation 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \leq 0. \quad (4)$$

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Replace x by $\pi(x)$ in Equation 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \leq 0. \quad (4)$$

Replace y by x and x by $\pi(y)$ in Equation 3

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle \leq 0. \quad (5)$$

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Replace x by $\pi(x)$ in Equation 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \leq 0. \quad (4)$$

Replace y by x and x by $\pi(y)$ in Equation 3

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle \leq 0. \quad (5)$$

(Equation 4)+(Equation 5) will cancel $\pi(y) - \pi(x)$, not good. So flip the sign of (Equation 5) gives

$$\langle \pi(x) - x, \pi(x) - \pi(y) \rangle \leq 0. \quad (6)$$

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Replace x by $\pi(x)$ in Equation 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \leq 0. \quad (4)$$

Replace y by x and x by $\pi(y)$ in Equation 3

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle \leq 0. \quad (5)$$

(Equation 4)+(Equation 5) will cancel $\pi(y) - \pi(x)$, not good. So flip the sign of (Equation 5) gives

$$\langle \pi(x) - x, \pi(x) - \pi(y) \rangle \leq 0. \quad (6)$$

$$\langle y - \pi(y) + \pi(x) - x, \pi(x) - \pi(y) \rangle \leq 0$$

$$\langle y - x, \pi(x) - \pi(y) \rangle \leq -\langle \pi(x) - \pi(y), \pi(x) - \pi(y) \rangle$$

$$\langle y - x, \pi(y) - \pi(x) \rangle \geq \|\pi(x) - \pi(y)\|_2^2$$

$$\|(y - x)^\top (\pi(y) - \pi(x))\|_2 \geq \|\pi(x) - \pi(y)\|_2^2$$

Projection operator is non-expansive

Shorthand notation: let $\pi = \text{proj}$ and $\pi(x)$ denotes $\text{proj}(x)$.

Begins with the variational characterization / obtuse angle inequality

$$\langle y - \pi(y), x - \pi(y) \rangle \leq 0 \quad \forall x \in S. \quad (3)$$

Replace x by $\pi(x)$ in Equation 3

$$\langle y - \pi(y), \pi(x) - \pi(y) \rangle \leq 0. \quad (4)$$

Replace y by x and x by $\pi(y)$ in Equation 3

$$\langle x - \pi(x), \pi(y) - \pi(x) \rangle \leq 0. \quad (5)$$

(Equation 4)+(Equation 5) will cancel $\pi(y) - \pi(x)$, not good. So flip the sign of (Equation 5) gives

$$\langle \pi(x) - x, \pi(x) - \pi(y) \rangle \leq 0. \quad (6)$$

$$\langle y - \pi(y) + \pi(x) - x, \pi(x) - \pi(y) \rangle \leq 0$$

$$\langle y - x, \pi(x) - \pi(y) \rangle \leq -\langle \pi(x) - \pi(y), \pi(x) - \pi(y) \rangle$$

$$\langle y - x, \pi(y) - \pi(x) \rangle \geq \|\pi(x) - \pi(y)\|_2^2$$

$$\|(y - x)^\top (\pi(y) - \pi(x))\|_2 \geq \|\pi(x) - \pi(y)\|_2^2$$

By Cauchy-Schwarz inequality, the left-hand-side is upper bounded by

$\|y - x\|_2 \|\pi(y) - \pi(x)\|_2$, we get
 $\|y - x\|_2 \|\pi(y) - \pi(x)\|_2 \geq \|\pi(x) - \pi(y)\|_2^2$.
Cancels $\|\pi(x) - \pi(y)\|_2$ finishes the proof.

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$

Build a hypothesis from the figure: $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$

Build a hypothesis from the figure: $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$

Check the inequality for a convex closed set: $(\pi - y)^T(x - \pi) \geq 0$

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$

Build a hypothesis from the figure: $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$

Check the inequality for a convex closed set: $(\pi - y)^T(x - \pi) \geq 0$

$$\begin{aligned} & \left(x_0 - y + R \frac{y - x_0}{\|y - x_0\|} \right)^T \left(x - x_0 - R \frac{y - x_0}{\|y - x_0\|} \right) = \\ & \left(\frac{(y - x_0)(R - \|y - x_0\|)}{\|y - x_0\|} \right)^T \left(\frac{(x - x_0)\|y - x_0\| - R(y - x_0)}{\|y - x_0\|} \right) = \\ & \frac{R - \|y - x_0\|}{\|y - x_0\|^2} (y - x_0)^T ((x - x_0)\|y - x_0\| - R(y - x_0)) = \\ & \frac{R - \|y - x_0\|}{\|y - x_0\|} \left((y - x_0)^T (x - x_0) - R\|y - x_0\| \right) = \\ & (R - \|y - x_0\|) \left(\frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R \right) \end{aligned}$$

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$

Build a hypothesis from the figure: $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$

Check the inequality for a convex closed set: $(\pi - y)^T(x - \pi) \geq 0$

The first factor is negative for point selection y .
The second factor is also negative, which follows from the Cauchy-Bunyakovsky inequality:

$$\begin{aligned} \left(x_0 - y + R \frac{y - x_0}{\|y - x_0\|} \right)^T \left(x - x_0 - R \frac{y - x_0}{\|y - x_0\|} \right) &= \\ \left(\frac{(y - x_0)(R - \|y - x_0\|)}{\|y - x_0\|} \right)^T \left(\frac{(x - x_0)\|y - x_0\| - R(y - x_0)}{\|y - x_0\|} \right) &= \\ \frac{R - \|y - x_0\|}{\|y - x_0\|^2} (y - x_0)^T ((x - x_0)\|y - x_0\| - R(y - x_0)) &= \\ \frac{R - \|y - x_0\|}{\|y - x_0\|} \left((y - x_0)^T (x - x_0) - R\|y - x_0\| \right) &= \\ (R - \|y - x_0\|) \left(\frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R \right) \end{aligned}$$

Example: projection on the ball

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid \|x - x_0\| \leq R\}$, $y \notin S$


Build a hypothesis from the figure: $\pi = x_0 + R \cdot \frac{y - x_0}{\|y - x_0\|}$

Check the inequality for a convex closed set: $(\pi - y)^T(x - \pi) \geq 0$

$$\begin{aligned} \left(x_0 - y + R \frac{y - x_0}{\|y - x_0\|} \right)^T \left(x - x_0 - R \frac{y - x_0}{\|y - x_0\|} \right) &= \\ \left(\frac{(y - x_0)(R - \|y - x_0\|)}{\|y - x_0\|} \right)^T \left(\frac{(x - x_0)\|y - x_0\| - R(y - x_0)}{\|y - x_0\|} \right) &= \\ \frac{R - \|y - x_0\|}{\|y - x_0\|^2} (y - x_0)^T ((x - x_0)\|y - x_0\| - R(y - x_0)) &= \\ \frac{R - \|y - x_0\|}{\|y - x_0\|} \left((y - x_0)^T (x - x_0) - R\|y - x_0\| \right) &= \\ (R - \|y - x_0\|) \left(\frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R \right) \end{aligned}$$

The first factor is negative for point selection y .
The second factor is also negative, which follows from the Cauchy-Bunyakovsky inequality:

$$\begin{aligned} (y - x_0)^T (x - x_0) &\leq \|y - x_0\| \|x - x_0\| \\ \frac{(y - x_0)^T (x - x_0)}{\|y - x_0\|} - R &\leq \frac{\|y - x_0\| \|x - x_0\|}{\|y - x_0\|} - R \end{aligned}$$



Example: projection on the halfspace

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid c^T x = b\}$, $y \notin S$. Build a hypothesis from the figure: $\pi = y + \alpha c$. Coefficient α is chosen so that $\pi \in S$: $c^T \pi = b$, so:

Example: projection on the halfspace

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid c^T x = b\}$, $y \notin S$. Build a hypothesis from the figure: $\pi = y + \alpha c$. Coefficient α is chosen so that $\pi \in S$: $c^T \pi = b$, so:




Figure 9: Hyperplane

Example: projection on the halfspace

Find $\pi_S(y) = \pi$, if $S = \{x \in \mathbb{R}^n \mid c^T x = b\}$, $y \notin S$. Build a hypothesis from the figure: $\pi = y + \alpha c$. Coefficient α is chosen so that $\pi \in S$: $c^T \pi = b$, so:




Figure 9: Hyperplane

$$c^T(y + \alpha c) = b$$

$$c^T y + \alpha c^T c = b$$

$$c^T y = b - \alpha c^T c$$

Check the inequality for a convex closed set:

$$(\pi - y)^T(x - \pi) \geq 0$$

$$(y + \alpha c - y)^T(x - y - \alpha c) =$$

$$\alpha c^T(x - y - \alpha c) =$$

$$\alpha(c^T x) - \alpha(c^T y) - \alpha^2(c^T c) =$$

$$\alpha b - \alpha(b - \alpha c^T c) - \alpha^2 c^T c =$$

$$\alpha b - \alpha b + \alpha^2 c^T c - \alpha^2 c^T c = 0 \geq 0$$

Projected Gradient Descent (PGD)

Idea

$$x_{k+1} = \text{proj}_S(x_k - \alpha_k \nabla f(x_k)) \quad \Leftrightarrow \quad \begin{aligned} y_k &= x_k - \alpha_k \nabla f(x_k) \\ x_{k+1} &= \text{proj}_S(y_k) \end{aligned}$$




Figure 10: Illustration of Projected Gradient Descent algorithm

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L \|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that $y_k = x_k - \frac{1}{L} \nabla f(x_k)$ and cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$:

(7)

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that $y_k = x_k - \frac{1}{L}\nabla f(x_k)$ and cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$:

$$\text{Smoothness: } f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

(7)

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that $y_k = x_k - \frac{1}{L}\nabla f(x_k)$ and cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$:

Smoothness:
$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

Method:
$$= f(x_k) - L\langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

(7)

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that $y_k = x_k - \frac{1}{L}\nabla f(x_k)$ and cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$:

Smoothness:
$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

Method:
$$= f(x_k) - L\langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

Cosine rule:
$$= f(x_k) - \frac{L}{2}(\|y_k - x_k\|^2 + \|x_{k+1} - x_k\|^2 - \|y_k - x_{k+1}\|^2) + \frac{L}{2}\|x_{k+1} - x_k\|^2 \quad (7)$$

Convergence rate for smooth and convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{L\|x_0 - x^*\|_2^2}{2k}$$

1. Let's prove sufficient decrease lemma, assuming, that $y_k = x_k - \frac{1}{L}\nabla f(x_k)$ and cosine rule $2x^T y = \|x\|^2 + \|y\|^2 - \|x - y\|^2$:

Smoothness:
$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

Method:
$$= f(x_k) - L\langle y_k - x_k, x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2$$

Cosine rule:
$$\begin{aligned} &= f(x_k) - \frac{L}{2}(\|y_k - x_k\|^2 + \|x_{k+1} - x_k\|^2 - \|y_k - x_{k+1}\|^2) + \frac{L}{2}\|x_{k+1} - x_k\|^2 \\ &= f(x_k) - \frac{1}{2L}\|\nabla f(x_k)\|^2 + \frac{L}{2}\|y_k - x_{k+1}\|^2 \end{aligned} \quad (7)$$

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle = \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right)$$
$$\langle \nabla f(x_k), x_k - x^* \rangle = \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)$$

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle = \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right)$$
$$\langle \nabla f(x_k), x_k - x^* \rangle = \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)$$

3. We will use now projection property: $\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$ with $x = x^*, y = y_k$:

$$\|x^* - \text{proj}_S(y_k)\|^2 + \|y_k - \text{proj}_S(y_k)\|^2 \leq \|x^* - y_k\|^2$$
$$\|y_k - x^*\|^2 \geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2$$

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle = \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right)$$
$$\langle \nabla f(x_k), x_k - x^* \rangle = \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)$$

3. We will use now projection property: $\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$ with $x = x^*, y = y_k$:

$$\|x^* - \text{proj}_S(y_k)\|^2 + \|y_k - \text{proj}_S(y_k)\|^2 \leq \|x^* - y_k\|^2$$
$$\|y_k - x^*\|^2 \geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2$$

4. Now, using convexity and previous part:

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle = \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right)$$
$$\langle \nabla f(x_k), x_k - x^* \rangle = \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)$$

3. We will use now projection property: $\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$ with $x = x^*, y = y_k$:

$$\|x^* - \text{proj}_S(y_k)\|^2 + \|y_k - \text{proj}_S(y_k)\|^2 \leq \|x^* - y_k\|^2$$
$$\|y_k - x^*\|^2 \geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2$$

4. Now, using convexity and previous part:

Convexity:
$$f(x_k) - f^* \leq \langle \nabla f(x_k), x_k - x^* \rangle$$

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{aligned}\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right) \\ \langle \nabla f(x_k), x_k - x^* \rangle &= \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)\end{aligned}$$

3. We will use now projection property: $\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$ with $x = x^*, y = y_k$:

$$\begin{aligned}\|x^* - \text{proj}_S(y_k)\|^2 + \|y_k - \text{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2\end{aligned}$$

4. Now, using convexity and previous part:

$$\begin{aligned}\text{Convexity:} \quad f(x_k) - f^* &\leq \langle \nabla f(x_k), x_k - x^* \rangle \\ &\leq \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 - \|y_k - x_{k+1}\|^2 \right)\end{aligned}$$

Convergence rate for smooth and convex case

2. Now we do not immediately have progress at each step. Let's use again cosine rule:

$$\begin{aligned}\left\langle \frac{1}{L} \nabla f(x_k), x_k - x^* \right\rangle &= \frac{1}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_k - x^* - \frac{1}{L} \nabla f(x_k)\|^2 \right) \\ \langle \nabla f(x_k), x_k - x^* \rangle &= \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|y_k - x^*\|^2 \right)\end{aligned}$$

3. We will use now projection property: $\|x - \text{proj}_S(y)\|^2 + \|y - \text{proj}_S(y)\|^2 \leq \|x - y\|^2$ with $x = x^*, y = y_k$:

$$\begin{aligned}\|x^* - \text{proj}_S(y_k)\|^2 + \|y_k - \text{proj}_S(y_k)\|^2 &\leq \|x^* - y_k\|^2 \\ \|y_k - x^*\|^2 &\geq \|x^* - x_{k+1}\|^2 + \|y_k - x_{k+1}\|^2\end{aligned}$$

4. Now, using convexity and previous part:

$$\begin{aligned}\text{Convexity:} \quad f(x_k) - f^* &\leq \langle \nabla f(x_k), x_k - x^* \rangle \\ &\leq \frac{L}{2} \left(\frac{1}{L^2} \|\nabla f(x_k)\|^2 + \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 - \|y_k - x_{k+1}\|^2 \right)\end{aligned}$$

$$\text{Sum for } i = 0, k-1 \quad \sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \sum_{i=0}^{k-1} \frac{1}{2L} \|\nabla f(x_i)\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2$$

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \sum_{i=0}^{k-1} \left[f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2$$

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

$$\begin{aligned}\sum_{i=0}^{k-1} [f(x_i) - f^*] &\leq \sum_{i=0}^{k-1} \left[f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2\end{aligned}$$

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

$$\begin{aligned}\sum_{i=0}^{k-1} [f(x_i) - f^*] &\leq \sum_{i=0}^{k-1} \left[f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2\end{aligned}$$

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

$$\begin{aligned}\sum_{i=0}^{k-1} [f(x_i) - f^*] &\leq \sum_{i=0}^{k-1} \left[f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ \sum_{i=0}^{k-1} f(x_i) - kf^* &\leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2\end{aligned}$$

Convergence rate for smooth and convex case

5. Bound gradients with sufficient decrease lemma 7:

$$\begin{aligned}\sum_{i=0}^{k-1} [f(x_i) - f^*] &\leq \sum_{i=0}^{k-1} \left[f(x_i) - f(x_{i+1}) + \frac{L}{2} \|y_i - x_{i+1}\|^2 \right] + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 + \frac{L}{2} \|x_0 - x^*\|^2 - \frac{L}{2} \sum_{i=0}^{i-1} \|y_i - x_{i+1}\|^2 \\ &\leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ \sum_{i=0}^{k-1} f(x_i) - kf^* &\leq f(x_0) - f(x_k) + \frac{L}{2} \|x_0 - x^*\|^2 \\ \sum_{i=1}^k [f(x_i) - f^*] &\leq \frac{L}{2} \|x_0 - x^*\|^2\end{aligned}$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that $x_{k+1} = \text{proj}_S(y_k)$. By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that $x_{k+1} = \text{proj}_S(y_k)$. By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$

and recall that $y_k = x_k - \frac{1}{L} \nabla f(x_k)$ implies $\|y_k - x_k\| = \frac{1}{L} \|\nabla f(x_k)\|$. Hence

$$\frac{L}{2} \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \|y_k - x_k\|^2 = \frac{L}{2} \frac{1}{L^2} \|\nabla f(x_k)\|^2 = \frac{1}{2L} \|\nabla f(x_k)\|^2.$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that $x_{k+1} = \text{proj}_S(y_k)$. By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$

and recall that $y_k = x_k - \frac{1}{L} \nabla f(x_k)$ implies $\|y_k - x_k\| = \frac{1}{L} \|\nabla f(x_k)\|$. Hence

$$\frac{L}{2} \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \|y_k - x_k\|^2 = \frac{L}{2} \frac{1}{L^2} \|\nabla f(x_k)\|^2 = \frac{1}{2L} \|\nabla f(x_k)\|^2.$$

Substitute back into (*):

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{1}{2L} \|\nabla f(x_k)\|^2 = f(x_k).$$

Convergence rate for smooth and convex case

6. From the sufficient decrease inequality

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{L}{2} \|y_k - x_{k+1}\|^2,$$

we use the fact that $x_{k+1} = \text{proj}_S(y_k)$. By definition of projection,

$$\|y_k - x_{k+1}\| \leq \|y_k - x_k\|,$$

and recall that $y_k = x_k - \frac{1}{L} \nabla f(x_k)$ implies $\|y_k - x_k\| = \frac{1}{L} \|\nabla f(x_k)\|$. Hence

$$\frac{L}{2} \|y_k - x_{k+1}\|^2 \leq \frac{L}{2} \|y_k - x_k\|^2 = \frac{L}{2} \frac{1}{L^2} \|\nabla f(x_k)\|^2 = \frac{1}{2L} \|\nabla f(x_k)\|^2.$$

Substitute back into (*):

$$f(x_{k+1}) \leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2 + \frac{1}{2L} \|\nabla f(x_k)\|^2 = f(x_k).$$

Hence

$$f(x_{k+1}) \leq f(x_k) \quad \text{for each } k,$$

so $\{f(x_k)\}$ is a monotonically nonincreasing sequence.

Convergence rate for smooth and convex case

7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Convergence rate for smooth and convex case

7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Convergence rate for smooth and convex case

7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Since $f(x_i)$ decreases in i , in particular $f(x_k) \leq f(x_i)$ for all $i \leq k$. Therefore

$$k [f(x_k) - f^*] \leq \sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2,$$

Convergence rate for smooth and convex case

7. Final convergence bound From step 5, we have already established

$$\sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2.$$

Since $f(x_i)$ decreases in i , in particular $f(x_k) \leq f(x_i)$ for all $i \leq k$. Therefore

$$k [f(x_k) - f^*] \leq \sum_{i=0}^{k-1} [f(x_i) - f^*] \leq \frac{L}{2} \|x_0 - x^*\|_2^2,$$

which immediately gives

$$f(x_k) - f^* \leq \frac{L \|x_0 - x^*\|_2^2}{2k}.$$

This completes the proof of the $\mathcal{O}(\frac{1}{k})$ convergence rate for convex and L -smooth f under projection constraints.

Convergence rate for smooth strongly convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be μ -strongly convex (or, PL-function with parameter μ) and L -smooth. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(x_0) - f^*)$$

1. We will use the notation $x_{k+1} = \text{proj}_S(x_k - \frac{1}{L}\nabla f(x_k)) = \pi(y_k)$. Since projection is nonexpansive, we have:

$$\|x_{k+1} - x_k\| = \|\pi(y_k) - \pi(x_k)\| \leq \|y_k - x_k\| = \frac{1}{L}\|\nabla f(x_k)\| \quad \Rightarrow \quad \|x_{k+1} - x_k\|^2 \leq \frac{1}{L^2}\|\nabla f(x_k)\|^2 \quad (8)$$

Convergence rate for smooth strongly convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be μ -strongly convex (or, PL-function with parameter μ) and L -smooth. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(x_0) - f^*)$$

1. We will use the notation $x_{k+1} = \text{proj}_S(x_k - \frac{1}{L}\nabla f(x_k)) = \pi(y_k)$. Since projection is nonexpansive, we have:

$$\|x_{k+1} - x_k\| = \|\pi(y_k) - \pi(x_k)\| \leq \|y_k - x_k\| = \frac{1}{L}\|\nabla f(x_k)\| \Rightarrow \|x_{k+1} - x_k\|^2 \leq \frac{1}{L^2}\|\nabla f(x_k)\|^2 \quad (8)$$

2. By L -smoothness of f , we have:

(9)

Convergence rate for smooth strongly convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be μ -strongly convex (or, PL-function with parameter μ) and L -smooth. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(x_0) - f^*)$$

1. We will use the notation $x_{k+1} = \text{proj}_S(x_k - \frac{1}{L}\nabla f(x_k)) = \pi(y_k)$. Since projection is nonexpansive, we have:

$$\|x_{k+1} - x_k\| = \|\pi(y_k) - \pi(x_k)\| \leq \|y_k - x_k\| = \frac{1}{L}\|\nabla f(x_k)\| \Rightarrow \|x_{k+1} - x_k\|^2 \leq \frac{1}{L^2}\|\nabla f(x_k)\|^2 \quad (8)$$

2. By L -smoothness of f , we have:

$$f(x_{k+1}) - f(x_k) \leq \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2 \quad (9)$$

Convergence rate for smooth strongly convex case

Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be μ -strongly convex (or, PL-function with parameter μ) and L -smooth. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; The Projected Gradient Descent algorithm with stepsize $\frac{1}{L}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(x_0) - f^*)$$

1. We will use the notation $x_{k+1} = \text{proj}_S(x_k - \frac{1}{L}\nabla f(x_k)) = \pi(y_k)$. Since projection is nonexpansive, we have:

$$\|x_{k+1} - x_k\| = \|\pi(y_k) - \pi(x_k)\| \leq \|y_k - x_k\| = \frac{1}{L}\|\nabla f(x_k)\| \Rightarrow \|x_{k+1} - x_k\|^2 \leq \frac{1}{L^2}\|\nabla f(x_k)\|^2 \quad (8)$$

2. By L -smoothness of f , we have:

$$\begin{aligned} f(x_{k+1}) - f(x_k) &\leq \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2}\|x_{k+1} - x_k\|^2 \\ &\leq \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{1}{2L}\|\nabla f(x_k)\|^2 \end{aligned} \quad (9)$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle \leq 0$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0\end{aligned}$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle \leq 0$$

$$\langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle \leq 0$$

$$\langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle \leq 0$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle \leq 0$$

$$\langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle \leq 0$$

$$\langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle \leq 0$$

$$\langle \nabla f(x_k), x_{k+1} - x_k \rangle \leq -L \|x_{k+1} - x_k\|^2$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle &\leq 0 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \|x_{k+1} - x_k\|^2 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \frac{1}{L^2} \|\nabla f(x_k)\|^2 = -\frac{1}{L} \|\nabla f(x_k)\|^2\end{aligned}$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle &\leq 0 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \|x_{k+1} - x_k\|^2 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \frac{1}{L^2} \|\nabla f(x_k)\|^2 = -\frac{1}{L} \|\nabla f(x_k)\|^2\end{aligned}$$

4. Returning back to the smoothness inequality (9) we obtain sufficient decrease property:

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle &\leq 0 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \|x_{k+1} - x_k\|^2 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \frac{1}{L^2} \|\nabla f(x_k)\|^2 = -\frac{1}{L} \|\nabla f(x_k)\|^2\end{aligned}$$

4. Returning back to the smoothness inequality (9) we obtain sufficient decrease property:

$$f(x_{k+1}) \leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{1}{2L} \|\nabla f(x_k)\|^2$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle &\leq 0 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \|x_{k+1} - x_k\|^2 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \frac{1}{L^2} \|\nabla f(x_k)\|^2 = -\frac{1}{L} \|\nabla f(x_k)\|^2\end{aligned}$$

4. Returning back to the smoothness inequality (9) we obtain sufficient decrease property:

$$\begin{aligned}f(x_{k+1}) &\leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{1}{2L} \|\nabla f(x_k)\|^2 \\ &\leq f(x_k) - \frac{1}{L} \|\nabla f(x_k)\|^2 + \frac{1}{2L} \|\nabla f(x_k)\|^2\end{aligned}$$

Convergence rate for smooth strongly convex case

3. By first-order optimality for that Euclidean projection problem, $\langle y_k - x_{k+1}, z - x_{k+1} \rangle \leq 0 \quad \forall z \in S$. In particular, for $z = x_k$,

$$\begin{aligned}\langle y_k - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - \frac{1}{L} \nabla f(x_k) - x_{k+1}, x_k - x_{k+1} \rangle &\leq 0 \\ \langle x_k - x_{k+1}, x_k - x_{k+1} \rangle - \frac{1}{L} \langle \nabla f(x_k), x_k - x_{k+1} \rangle &\leq 0 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \|x_{k+1} - x_k\|^2 \\ \langle \nabla f(x_k), x_{k+1} - x_k \rangle &\leq -L \frac{1}{L^2} \|\nabla f(x_k)\|^2 = -\frac{1}{L} \|\nabla f(x_k)\|^2\end{aligned}$$

4. Returning back to the smoothness inequality (9) we obtain sufficient decrease property:

$$\begin{aligned}f(x_{k+1}) &\leq f(x_k) + \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{1}{2L} \|\nabla f(x_k)\|^2 \\ &\leq f(x_k) - \frac{1}{L} \|\nabla f(x_k)\|^2 + \frac{1}{2L} \|\nabla f(x_k)\|^2 \\ &\leq f(x_k) - \frac{1}{2L} \|\nabla f(x_k)\|^2\end{aligned}$$

Convergence rate for smooth strongly convex case

5. If we have a PL-function, we can use the following inequality: $\|\nabla f(x_k)\|^2 \geq 2\mu (f(x_k) - f^*)$. Thus, we can write:

$$\begin{aligned} f(x_{k+1}) &\leq f(x_k) - \frac{\mu}{L} (f(x_k) - f^*) \\ f(x_{k+1}) - f^* &\leq f(x_k) - f^* - \frac{\mu}{L} (f(x_k) - f^*) \end{aligned}$$

Convergence rate for smooth strongly convex case

5. If we have a PL-function, we can use the following inequality: $\|\nabla f(x_k)\|^2 \geq 2\mu (f(x_k) - f^*)$. Thus, we can write:

$$\begin{aligned} f(x_{k+1}) &\leq f(x_k) - \frac{\mu}{L} (f(x_k) - f^*) \\ f(x_{k+1}) - f^* &\leq f(x_k) - f^* - \frac{\mu}{L} (f(x_k) - f^*) \end{aligned}$$

6. Now we can use induction to get the following bound:

$$f(x_k) - f^* \leq \left(1 - \frac{\mu}{L}\right)^k (f(x_0) - f^*)$$

Frank-Wolfe Method

Idea




Figure 11: Illustration of Frank-Wolfe (conditional gradient) algorithm

Idea




Figure 12: Illustration of Frank-Wolfe (conditional gradient) algorithm




Figure 13: Illustration of Frank-Wolfe (conditional gradient) algorithm

Idea




Figure 14: Illustration of Frank-Wolfe (conditional gradient) algorithm




Figure 15: Illustration of Frank-Wolfe (conditional gradient) algorithm




Figure 16: Illustration of Frank-Wolfe (conditional gradient) algorithm

Idea




Figure 17: Illustration of Frank-Wolfe (conditional gradient) algorithm

Idea

$$y_k = \arg \min_{x \in S} f_{x_k}^I(x) = \arg \min_{x \in S} \langle \nabla f(x_k), x \rangle$$

$$x_{k+1} = \gamma_k x_k + (1 - \gamma_k) y_k$$




Figure 18: Illustration of Frank-Wolfe (conditional gradient) algorithm

Convergence rate for smooth and convex case

i Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Frank-Wolfe algorithm with step size $\gamma_k = \frac{k-1}{k+1}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{2LR^2}{k+1}$$

where $R = \max_{x,y \in S} \|x - y\|$ is the diameter of the set S .

Convergence rate for smooth and convex case

i Theorem

Let $f : \mathbb{R}^n \rightarrow \mathbb{R}$ be convex and differentiable. Let $S \subseteq \mathbb{R}^n$ be a closed convex set, and assume that there is a minimizer x^* of f over S ; furthermore, suppose that f is smooth over S with parameter L . The Frank-Wolfe algorithm with step size $\gamma_k = \frac{k-1}{k+1}$ achieves the following convergence after iteration $k > 0$:

$$f(x_k) - f^* \leq \frac{2LR^2}{k+1}$$

where $R = \max_{x,y \in S} \|x - y\|$ is the diameter of the set S .

1. By L -smoothness of f , we have:

$$\begin{aligned} f(x_{k+1}) - f(x_k) &\leq \langle \nabla f(x_k), x_{k+1} - x_k \rangle + \frac{L}{2} \|x_{k+1} - x_k\|^2 \\ &= (1 - \gamma_k) \langle \nabla f(x_k), y_k - x_k \rangle + \frac{L(1 - \gamma_k)^2}{2} \|y_k - x_k\|^2 \end{aligned}$$

Convergence rate for smooth and convex case

2. By convexity of f , for any $x \in S$, including x^* :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for $x = x^*$:

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

Convergence rate for smooth and convex case

2. By convexity of f , for any $x \in S$, including x^* :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for $x = x^*$:

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of y_k , we have $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$, thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \leq \langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

Convergence rate for smooth and convex case

2. By convexity of f , for any $x \in S$, including x^* :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for $x = x^*$:

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of y_k , we have $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$, thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \leq \langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

4. Combining the above inequalities:

$$\begin{aligned} f(x_{k+1}) - f(x_k) &\leq (1 - \gamma_k) \langle \nabla f(x_k), y_k - x_k \rangle + \frac{L(1 - \gamma_k)^2}{2} \|y_k - x_k\|^2 \\ &\leq (1 - \gamma_k) (f(x^*) - f(x_k)) + \frac{L(1 - \gamma_k)^2}{2} R^2 \end{aligned}$$

Convergence rate for smooth and convex case

2. By convexity of f , for any $x \in S$, including x^* :

$$\langle \nabla f(x_k), x - x_k \rangle \leq f(x) - f(x_k)$$

In particular, for $x = x^*$:

$$\langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

3. By definition of y_k , we have $\langle \nabla f(x_k), y_k \rangle \leq \langle \nabla f(x_k), x^* \rangle$, thus:

$$\langle \nabla f(x_k), y_k - x_k \rangle \leq \langle \nabla f(x_k), x^* - x_k \rangle \leq f(x^*) - f(x_k)$$

4. Combining the above inequalities:

$$\begin{aligned} f(x_{k+1}) - f(x_k) &\leq (1 - \gamma_k) \langle \nabla f(x_k), y_k - x_k \rangle + \frac{L(1 - \gamma_k)^2}{2} \|y_k - x_k\|^2 \\ &\leq (1 - \gamma_k) (f(x^*) - f(x_k)) + \frac{L(1 - \gamma_k)^2}{2} R^2 \end{aligned}$$

5. Rearranging terms:

$$f(x_{k+1}) - f(x^*) \leq \gamma_k (f(x_k) - f(x^*)) + (1 - \gamma_k)^2 \frac{LR^2}{2}$$

Convergence rate for smooth and convex case

6. Denoting $\delta_k = \frac{f(x_k) - f(x^*)}{LR^2}$, we get:

$$\delta_{k+1} \leq \gamma_k \delta_k + \frac{(1 - \gamma_k)^2}{2} = \frac{k-1}{k+1} \delta_k + \frac{2}{(k+1)^2}$$

Convergence rate for smooth and convex case

6. Denoting $\delta_k = \frac{f(x_k) - f(x^*)}{LR^2}$, we get:

$$\delta_{k+1} \leq \gamma_k \delta_k + \frac{(1 - \gamma_k)^2}{2} = \frac{k-1}{k+1} \delta_k + \frac{2}{(k+1)^2}$$

7. Starting from $\delta_2 \leq \frac{1}{2}$ and applying induction on k , we can show that:

$$\delta_k \leq \frac{2}{k+1}$$

which gives us the desired result:

$$f(x_k) - f^* \leq \frac{2LR^2}{k+1}$$