


# Introduction to dual methods

Daniil Merkulov

Optimization methods. MIPT

## Definition




Recall that given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , the function defined by



$$f^*(y) = \max_x [y^T x - f(x)]$$

is called its conjugate.

## Geometrical intuition



## Geometrical intuition



# Conjugate function properties

Recall that given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x [y^T x - f(x)]$$

is called its conjugate.

- Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x [f(x) - y^T x]$$

# Conjugate function properties

Recall that given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x [y^T x - f(x)]$$

is called its conjugate.

- Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x [f(x) - y^T x]$$

- If  $f$  is closed and convex, then  $f^{**} = f$ . Also,

$$x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x) \Leftrightarrow x \in \arg \min_z [f(z) - y^T z]$$

# Conjugate function properties

Recall that given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$ , the function defined by

$$f^*(y) = \max_x [y^T x - f(x)]$$

is called its conjugate.

- Conjugates appear frequently in dual programs, since

$$-f^*(y) = \min_x [f(x) - y^T x]$$

- If  $f$  is closed and convex, then  $f^{**} = f$ . Also,

$$x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x) \Leftrightarrow x \in \arg \min_z [f(z) - y^T z]$$

- If  $f$  is strictly convex, then

$$\nabla f^*(y) = \arg \min_z [f(z) - y^T z]$$

## Conjugate function properties (proofs)

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that  $f$  is convex and closed.

- **Proof of  $\Leftarrow$ :** Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^T z - f(z)$  over  $z$ . But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{and} \quad \partial f^*(y) = \text{cl}(\text{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .



# Conjugate function properties (proofs)

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that  $f$  is convex and closed.

- **Proof of  $\Leftarrow$ :** Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^T z - f(z)$  over  $z$ . But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{and} \quad \partial f^*(y) = \text{cl}(\text{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

- **Proof of  $\Rightarrow$ :** From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .

# Conjugate function properties (proofs)

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that  $f$  is convex and closed.

- **Proof of  $\Leftarrow$ :** Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^T z - f(z)$  over  $z$ . But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{and} \quad \partial f^*(y) = \text{cl}(\text{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

- **Proof of  $\Rightarrow$ :** From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .

## Conjugate function properties (proofs)

We will show that  $x \in \partial f^*(y) \Leftrightarrow y \in \partial f(x)$ , assuming that  $f$  is convex and closed.

- **Proof of  $\Leftarrow$ :** Suppose  $y \in \partial f(x)$ . Then  $x \in M_y$ , the set of maximizers of  $y^T z - f(z)$  over  $z$ . But

$$f^*(y) = \max_z \{y^T z - f(z)\} \quad \text{and} \quad \partial f^*(y) = \text{cl}(\text{conv}(\bigcup_{z \in M_y} \{z\})).$$

Thus  $x \in \partial f^*(y)$ .

- **Proof of  $\Rightarrow$ :** From what we showed above, if  $x \in \partial f^*(y)$ , then  $y \in \partial f^*(x)$ , but  $f^{**} = f$ .

Clearly  $y \in \partial f(x) \Leftrightarrow x \in \arg \min_z \{f(z) - y^T z\}$

Lastly, if  $f$  is strictly convex, then we know that  $f(z) - y^T z$  has a unique minimizer over  $z$ , and this must be  $\nabla f^*(y)$ .

## Dual (sub)gradient method

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

## Dual (sub)gradient method

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_u -f^*(-A^T u) - b^T u$$

where  $f^*$  is the conjugate of  $f$ . Defining  $g(u) = -f^*(-A^T u) - b^T u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

## Dual (sub)gradient method

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_u -f^*(-A^T u) - b^T u$$

where  $f^*$  is the conjugate of  $f$ . Defining  $g(u) = -f^*(-A^T u) - b^T u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b \quad \text{where} \quad x \in \arg \min_z [f(z) + u^T Az]$$

## Dual (sub)gradient method

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_u -f^*(-A^T u) - b^T u$$

where  $f^*$  is the conjugate of  $f$ . Defining  $g(u) = -f^*(-A^T u) - b^T u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b \quad \text{where} \quad x \in \arg \min_z [f(z) + u^T Az]$$

Dual ascent method for maximizing dual objective:

- Step sizes  $\alpha_k$ ,  $k = 1, 2, 3, \dots$ , are chosen in standard ways.

i

$$x_k \in \arg \min_x [f(x) + (u_{k-1})^T Ax]$$

$$u_k = u_{k-1} + \alpha_k (Ax_k - b)$$

## Dual (sub)gradient method

Even if we can't derive dual (conjugate) in closed form, we can still use dual-based gradient or subgradient methods.

Consider the problem:

$$\min_x f(x) \quad \text{subject to} \quad Ax = b$$

Its dual problem is:

$$\max_u -f^*(-A^T u) - b^T u$$

where  $f^*$  is the conjugate of  $f$ . Defining  $g(u) = -f^*(-A^T u) - b^T u$ , note that:

$$\partial g(u) = A \partial f^*(-A^T u) - b$$

Therefore, using what we know about conjugates

$$\partial g(u) = Ax - b \quad \text{where} \quad x \in \arg \min_z [f(z) + u^T Az]$$

Dual ascent method for maximizing dual objective:

i



$$x_k \in \arg \min_x [f(x) + (u_{k-1})^T Ax]$$

$$u_k = u_{k-1} + \alpha_k (Ax_k - b)$$

- Step sizes  $\alpha_k$ ,  $k = 1, 2, 3, \dots$ , are chosen in standard ways.
- Proximal gradients and acceleration can be applied as they would usually.



## Slopes of $f$ and $f^*$



## Slopes of $f$ and $f^*$

Assume that  $f$  is a closed and convex function. Then  $f$  is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

## Slopes of $f$ and $f^*$

Assume that  $f$  is a closed and convex function. Then  $f$  is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of “ $\Rightarrow$ ”:** Recall, if  $g$  is strongly convex with minimizer  $x$ , then

$$g(y) \geq g(x) + \frac{\mu}{2} \|y - x\|^2, \quad \text{for all } y$$

## Slopes of $f$ and $f^*$

Assume that  $f$  is a closed and convex function. Then  $f$  is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of “ $\Rightarrow$ ”:** Recall, if  $g$  is strongly convex with minimizer  $x$ , then

$$g(y) \geq g(x) + \frac{\mu}{2} \|y - x\|^2, \quad \text{for all } y$$

Hence, defining  $x_u = \nabla f^*(u)$  and  $x_v = \nabla f^*(v)$ ,

$$f(x_v) - u^T x_v \geq f(x_u) - u^T x_u + \frac{\mu}{2} \|x_u - x_v\|^2$$

$$f(x_u) - v^T x_u \geq f(x_v) - v^T x_v + \frac{\mu}{2} \|x_u - x_v\|^2$$

## Slopes of $f$ and $f^*$

Assume that  $f$  is a closed and convex function. Then  $f$  is strongly convex with parameter  $\mu \Leftrightarrow \nabla f^*$  is Lipschitz with parameter  $1/\mu$ .

**Proof of “ $\Rightarrow$ ”:** Recall, if  $g$  is strongly convex with minimizer  $x$ , then

$$g(y) \geq g(x) + \frac{\mu}{2} \|y - x\|^2, \quad \text{for all } y$$

Hence, defining  $x_u = \nabla f^*(u)$  and  $x_v = \nabla f^*(v)$ ,

$$f(x_v) - u^T x_v \geq f(x_u) - u^T x_u + \frac{\mu}{2} \|x_u - x_v\|^2$$

$$f(x_u) - v^T x_u \geq f(x_v) - v^T x_v + \frac{\mu}{2} \|x_u - x_v\|^2$$

Adding these together, using the Cauchy-Schwarz inequality, and rearranging shows that

$$\|x_u - x_v\|^2 \leq \frac{1}{\mu} \|u - v\|^2$$

## Slopes of $f$ and $f^*$

**Proof of “ $\Leftarrow$ ”:** for simplicity, call  $g = f^*$  and  $L = \frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant  $L$ , so is  $g_x(z) = g(z) - \nabla g(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T (z - y) + \frac{L}{2} \|z - y\|_2^2$$

## Slopes of $f$ and $f^*$

**Proof of “ $\Leftarrow$ ”:** for simplicity, call  $g = f^*$  and  $L = \frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant  $L$ , so is  $g_x(z) = g(z) - \nabla g(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T (z - y) + \frac{L}{2} \|z - y\|_2^2$$

Minimizing each side over  $z$ , and rearranging, gives

$$\frac{1}{2L} \|\nabla g(x) - \nabla g(y)\|^2 \leq g(y) - g(x) + \nabla g(x)^T (x - y)$$

## Slopes of $f$ and $f^*$

**Proof of “ $\Leftarrow$ ”:** for simplicity, call  $g = f^*$  and  $L = \frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant  $L$ , so is  $g_x(z) = g(z) - \nabla g(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T (z - y) + \frac{L}{2} \|z - y\|_2^2$$

Minimizing each side over  $z$ , and rearranging, gives

$$\frac{1}{2L} \|\nabla g(x) - \nabla g(y)\|^2 \leq g(y) - g(x) + \nabla g(x)^T (x - y)$$

Exchanging roles of  $x$ ,  $y$ , and adding together, gives

$$\frac{1}{L} \|\nabla g(x) - \nabla g(y)\|^2 \leq (\nabla g(x) - \nabla g(y))^T (x - y)$$



## Slopes of $f$ and $f^*$

**Proof of “ $\Leftarrow$ ”:** for simplicity, call  $g = f^*$  and  $L = \frac{1}{\mu}$ . As  $\nabla g$  is Lipschitz with constant  $L$ , so is  $g_x(z) = g(z) - \nabla g(x)^T z$ , hence

$$g_x(z) \leq g_x(y) + \nabla g_x(y)^T (z - y) + \frac{L}{2} \|z - y\|_2^2$$

Minimizing each side over  $z$ , and rearranging, gives

$$\frac{1}{2L} \|\nabla g(x) - \nabla g(y)\|^2 \leq g(y) - g(x) + \nabla g(x)^T (x - y)$$

Exchanging roles of  $x$ ,  $y$ , and adding together, gives

$$\frac{1}{L} \|\nabla g(x) - \nabla g(y)\|^2 \leq (\nabla g(x) - \nabla g(y))^T (x - y)$$

Let  $u = \nabla f(x)$ ,  $v = \nabla g(y)$ ; then  $x \in \partial g^*(u)$ ,  $y \in \partial g^*(v)$ , and the above reads  $(x - y)^T (u - v) \geq \frac{\|u - v\|^2}{L}$ , implying the result.



# Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent:<sup>1</sup>

- If  $f$  is strongly convex with parameter  $\mu$ , then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .

---

<sup>1</sup>This is ignoring the role of  $A$ , and thus reflects the case when the singular values of  $A$  are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\mu} + \frac{\sigma_{\min}(A)^2}{L}}$  (second case).



# Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent:<sup>1</sup>

- If  $f$  is strongly convex with parameter  $\mu$ , then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .
- If  $f$  is strongly convex with parameter  $\mu$  and  $\nabla f$  is Lipschitz with parameter  $L$ , then dual gradient ascent with step sizes  $\alpha_k = \frac{2}{\frac{1}{\mu} + \frac{1}{L}}$  converges at linear rate  $O(\log(\frac{1}{\epsilon}))$ .

---

<sup>1</sup>This is ignoring the role of  $A$ , and thus reflects the case when the singular values of  $A$  are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\mu} + \frac{\sigma_{\min}(A)^2}{L}}$  (second case).



# Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent:<sup>1</sup>

- If  $f$  is strongly convex with parameter  $\mu$ , then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .
- If  $f$  is strongly convex with parameter  $\mu$  and  $\nabla f$  is Lipschitz with parameter  $L$ , then dual gradient ascent with step sizes  $\alpha_k = \frac{2}{\frac{1}{\mu} + \frac{1}{L}}$  converges at linear rate  $O(\log(\frac{1}{\epsilon}))$ .

---

<sup>1</sup>This is ignoring the role of  $A$ , and thus reflects the case when the singular values of  $A$  are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\mu} + \frac{\sigma_{\min}(A)^2}{L}}$  (second case).





# Convergence guarantees

The following results hold from combining the last fact with what we already know about gradient descent:<sup>1</sup>

- If  $f$  is strongly convex with parameter  $\mu$ , then dual gradient ascent with constant step sizes  $\alpha_k = \mu$  converges at sublinear rate  $O(\frac{1}{\epsilon})$ .
- If  $f$  is strongly convex with parameter  $\mu$  and  $\nabla f$  is Lipschitz with parameter  $L$ , then dual gradient ascent with step sizes  $\alpha_k = \frac{2}{\frac{1}{\mu} + \frac{1}{L}}$  converges at linear rate  $O(\log(\frac{1}{\epsilon}))$ .

Note that this describes convergence in the dual. (Convergence in the primal requires more assumptions)

---

<sup>1</sup>This is ignoring the role of  $A$ , and thus reflects the case when the singular values of  $A$  are all close to 1. To be more precise, the step sizes here should be:  $\frac{\mu}{\sigma_{\max}(A)^2}$  (first case) and  $\frac{2}{\frac{\sigma_{\max}(A)^2}{\mu} + \frac{\sigma_{\min}(A)^2}{L}}$  (second case).

# Dual decomposition

Consider

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad Ax = b$$

## Dual decomposition

Consider

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x = (x_1, \dots, x_B) \in \mathbb{R}^n$  divides into  $B$  blocks of variables, with each  $x_i \in \mathbb{R}^{n_i}$ . We can also partition  $A$  accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

## Dual decomposition

Consider

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x = (x_1, \dots, x_B) \in \mathbb{R}^n$  divides into  $B$  blocks of variables, with each  $x_i \in \mathbb{R}^{n_i}$ . We can also partition  $A$  accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into  $B$  separate problems:

$$\begin{aligned} x^{\text{new}} &\in \arg \min_x \left( \sum_{i=1}^B f_i(x_i) + u^T Ax \right) \\ \Rightarrow x_i^{\text{new}} &\in \arg \min_{x_i} \left( f_i(x_i) + u^T A_i x_i \right), \quad i = 1, \dots, B \end{aligned}$$

$$x_i^k \in \arg \min_{x_i} \left( f_i(x_i) + (u^{k-1})^T A_i x_i \right), \quad i = 1, \dots, B$$

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

## Dual decomposition

Consider

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x = (x_1, \dots, x_B) \in \mathbb{R}^n$  divides into  $B$  blocks of variables, with each  $x_i \in \mathbb{R}^{n_i}$ . We can also partition  $A$  accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into  $B$  separate problems:

$$\begin{aligned} x^{\text{new}} &\in \arg \min_x \left( \sum_{i=1}^B f_i(x_i) + u^T Ax \right) \\ \Rightarrow x_i^{\text{new}} &\in \arg \min_{x_i} (f_i(x_i) + u^T A_i x_i), \quad i = 1, \dots, B \end{aligned}$$

$$x_i^k \in \arg \min_{x_i} (f_i(x_i) + (u^{k-1})^T A_i x_i), \quad i = 1, \dots, B$$

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

Can think of these steps as:

- **Broadcast:** Send  $u$  to each of the  $B$  processors, each optimizes in parallel to find  $x_i$ .

## Dual decomposition

Consider

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad Ax = b$$

Here  $x = (x_1, \dots, x_B) \in \mathbb{R}^n$  divides into  $B$  blocks of variables, with each  $x_i \in \mathbb{R}^{n_i}$ . We can also partition  $A$  accordingly:

$$A = [A_1 \dots A_B], \text{ where } A_i \in \mathbb{R}^{m \times n_i}$$

Simple but powerful observation, in calculation of subgradient, is that the minimization decomposes into  $B$  separate problems:

$$\begin{aligned} x^{\text{new}} &\in \arg \min_x \left( \sum_{i=1}^B f_i(x_i) + u^T Ax \right) \\ \Rightarrow x_i^{\text{new}} &\in \arg \min_{x_i} (f_i(x_i) + u^T A_i x_i), \quad i = 1, \dots, B \end{aligned}$$

$$x_i^k \in \arg \min_{x_i} (f_i(x_i) + (u^{k-1})^T A_i x_i), \quad i = 1, \dots, B$$

$$u^k = u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right)$$

Can think of these steps as:

- **Broadcast:** Send  $u$  to each of the  $B$  processors, each optimizes in parallel to find  $x_i$ .
- **Gather:** Collect  $A_i x_i$  from each processor, update the global dual variable  $u$ .

# Inequality constraints

Consider the optimization problem:

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^B A_i x_i \leq b$$

# Inequality constraints

Consider the optimization problem:

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^B A_i x_i \leq b$$

Using **dual decomposition**, specifically the **projected subgradient method**, the iterative steps can be expressed as:

- The primal update step:

$$x_i^k \in \arg \min_{x_i} \left[ f_i(x_i) + (u^{k-1})^T A_i x_i \right], \quad i = 1, \dots, B$$



# Inequality constraints

Consider the optimization problem:

$$\min_x \sum_{i=1}^B f_i(x_i) \quad \text{subject to} \quad \sum_{i=1}^B A_i x_i \leq b$$

Using **dual decomposition**, specifically the **projected subgradient method**, the iterative steps can be expressed as:

- The primal update step:

$$x_i^k \in \arg \min_{x_i} \left[ f_i(x_i) + (u^{k-1})^T A_i x_i \right], \quad i = 1, \dots, B$$

- The dual update step:

$$u^k = \left( u^{k-1} + \alpha_k \left( \sum_{i=1}^B A_i x_i^k - b \right) \right)_+$$

where  $(u)_+$  denotes the positive part of  $u$ , i.e.,  $(u_+)_i = \max\{0, u_i\}$ , for  $i = 1, \dots, m$ .

# Price Coordination Interpretation (Vandenbergh)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.

# Price Coordination Interpretation (Vandenberghe)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .

# Price Coordination Interpretation (Vandenberghe)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .
- **Dual Update Rule:**

$$u_j^{\text{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^B A_i x_i$  represents the slacks.

# Price Coordination Interpretation (Vandenbergh)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .
- **Dual Update Rule:**

$$u_j^{\text{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^B A_i x_i$  represents the slacks.

- **Price Adjustments:**

# Price Coordination Interpretation (Vandenberghe)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .

- **Dual Update Rule:**

$$u_j^{\text{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^B A_i x_i$  represents the slacks.

- **Price Adjustments:**
  - **Increase price**  $u_j$  if resource  $j$  is over-utilized ( $s_j < 0$ ).

# Price Coordination Interpretation (Vandenberghe)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .

- **Dual Update Rule:**

$$u_j^{\text{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^B A_i x_i$  represents the slacks.

- **Price Adjustments:**
  - **Increase price**  $u_j$  if resource  $j$  is over-utilized ( $s_j < 0$ ).
  - **Decrease price**  $u_j$  if resource  $j$  is under-utilized ( $s_j > 0$ ).

# Price Coordination Interpretation (Vandenberghe)

- **System Overview:** Consider a system with  $B$  units, where each unit independently chooses its decision variable  $x_i$ , which determines how to allocate its goods.
- **Resource Constraints:** These are limits on shared resources, represented by the rows of  $A$ . Each component of the dual variable  $u_j$  represents the price of resource  $j$ .

- **Dual Update Rule:**

$$u_j^{\text{new}} = (u_j - ts_j)_+, \quad j = 1, \dots, m$$

where  $s = b - \sum_{i=1}^B A_i x_i$  represents the slacks.

- **Price Adjustments:**

- **Increase price**  $u_j$  if resource  $j$  is over-utilized ( $s_j < 0$ ).
- **Decrease price**  $u_j$  if resource  $j$  is under-utilized ( $s_j > 0$ ).
- **Never let prices get negative;** hence the use of the positive part notation  $(\cdot)_+$ .



# Augmented Lagrangian method aka method of multipliers

**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\begin{aligned} \min_x \quad & f(x) + \frac{\rho}{2} \|Ax - b\|^2 \\ \text{s.t.} \quad & Ax = b \end{aligned}$$

# Augmented Lagrangian method aka method of multipliers

**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\begin{aligned} \min_x \quad & f(x) + \frac{\rho}{2} \|Ax - b\|^2 \\ \text{s.t.} \quad & Ax = b \end{aligned}$$

where  $\rho > 0$  is a parameter. This formulation is clearly equivalent to the original problem. The problem is strongly convex if matrix  $A$  has full column rank.

# Augmented Lagrangian method aka method of multipliers

**Dual ascent disadvantage:** convergence requires strong conditions. Augmented Lagrangian method transforms the primal problem:

$$\begin{aligned} \min_x \quad & f(x) + \frac{\rho}{2} \|Ax - b\|^2 \\ \text{s.t.} \quad & Ax = b \end{aligned}$$

where  $\rho > 0$  is a parameter. This formulation is clearly equivalent to the original problem. The problem is strongly convex if matrix  $A$  has full column rank.

**Dual gradient ascent:** The iterative updates are given by:

$$\begin{aligned} x_k &= \arg \min_x \left[ f(x) + (u_{k-1})^T Ax + \frac{\rho}{2} \|Ax - b\|^2 \right] \\ u_k &= u_{k-1} + \rho(Ax_k - b) \end{aligned}$$

# Augmented Lagrangian method aka method of multipliers

**Notice step size choice  $\alpha_k = \rho$  in dual algorithm. Why?**

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T Ax + \frac{\rho}{2} \|Ax - b\|^2$$

over  $x$ , we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T (u_{k-1} + \rho(Ax_k - b))$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$

# Augmented Lagrangian method aka method of multipliers

**Notice step size choice  $\alpha_k = \rho$  in dual algorithm. Why?**

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T Ax + \frac{\rho}{2} \|Ax - b\|^2$$

over  $x$ , we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T (u_{k-1} + \rho(Ax_k - b))$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$

This represents the stationarity condition for the original primal problem; under mild conditions,  $Ax_k - b \rightarrow 0$  as  $k \rightarrow \infty$ , so the KKT conditions are satisfied in the limit and  $x_k, u_k$  converge to the solutions.

- **Advantage:** The augmented Lagrangian gives better convergence.

# Augmented Lagrangian method aka method of multipliers

**Notice step size choice  $\alpha_k = \rho$  in dual algorithm. Why?**

Since  $x_k$  minimizes the function:

$$f(x) + (u_{k-1})^T Ax + \frac{\rho}{2} \|Ax - b\|^2$$

over  $x$ , we have the stationarity condition:

$$0 \in \partial f(x_k) + A^T (u_{k-1} + \rho(Ax_k - b))$$

which simplifies to:

$$\partial f(x_k) + A^T u_k$$

This represents the stationarity condition for the original primal problem; under mild conditions,  $Ax_k - b \rightarrow 0$  as  $k \rightarrow \infty$ , so the KKT conditions are satisfied in the limit and  $x_k, u_k$  converge to the solutions.

- **Advantage:** The augmented Lagrangian gives better convergence.
- **Disadvantage:** We lose decomposability! (Separability is ruined)

# Alternating Direction Method of Multipliers (ADMM)

**Alternating direction method of multipliers** or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\min_{x,z} f(x) + g(z)$$

$$\text{s.t. } Ax + Bz = c$$

# Alternating Direction Method of Multipliers (ADMM)

**Alternating direction method of multipliers** or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\begin{aligned} \min_{x,z} \quad & f(x) + g(z) \\ \text{s.t.} \quad & Ax + Bz = c \end{aligned}$$

We augment the objective to include a penalty term for constraint violation:

$$\begin{aligned} \min_{x,z} \quad & f(x) + g(z) + \frac{\rho}{2} \|Ax + Bz - c\|^2 \\ \text{s.t.} \quad & Ax + Bz = c \end{aligned}$$



# Alternating Direction Method of Multipliers (ADMM)

**Alternating direction method of multipliers** or ADMM aims for the best of both worlds. Consider the following optimization problem:

Minimize the function:

$$\begin{aligned} \min_{x,z} \quad & f(x) + g(z) \\ \text{s.t.} \quad & Ax + Bz = c \end{aligned}$$

We augment the objective to include a penalty term for constraint violation:

$$\begin{aligned} \min_{x,z} \quad & f(x) + g(z) + \frac{\rho}{2} \|Ax + Bz - c\|^2 \\ \text{s.t.} \quad & Ax + Bz = c \end{aligned}$$

where  $\rho > 0$  is a parameter. The augmented Lagrangian for this problem is defined as:

$$L_\rho(x, z, u) = f(x) + g(z) + u^T (Ax + Bz - c) + \frac{\rho}{2} \|Ax + Bz - c\|^2$$

# Alternating Direction Method of Multipliers (ADMM)

ADMM repeats the following steps, for  $k = 1, 2, 3, \dots$ :

1. Update  $x$ :

$$x_k = \arg \min_x L_\rho(x, z_{k-1}, u_{k-1})$$

# Alternating Direction Method of Multipliers (ADMM)

ADMM repeats the following steps, for  $k = 1, 2, 3, \dots$ :

1. Update  $x$ :

$$x_k = \arg \min_x L_\rho(x, z_{k-1}, u_{k-1})$$

2. Update  $z$ :

$$z_k = \arg \min_z L_\rho(x_k, z, u_{k-1})$$

# Alternating Direction Method of Multipliers (ADMM)

ADMM repeats the following steps, for  $k = 1, 2, 3, \dots$ :

1. Update  $x$ :

$$x_k = \arg \min_x L_\rho(x, z_{k-1}, u_{k-1})$$

2. Update  $z$ :

$$z_k = \arg \min_z L_\rho(x_k, z, u_{k-1})$$

3. Update  $u$ :

$$u_k = u_{k-1} + \rho(Ax_k + Bz_k - c)$$

# Alternating Direction Method of Multipliers (ADMM)

ADMM repeats the following steps, for  $k = 1, 2, 3, \dots$ :

1. Update  $x$ :

$$x_k = \arg \min_x L_\rho(x, z_{k-1}, u_{k-1})$$

2. Update  $z$ :

$$z_k = \arg \min_z L_\rho(x_k, z, u_{k-1})$$

3. Update  $u$ :

$$u_k = u_{k-1} + \rho(Ax_k + Bz_k - c)$$

# Alternating Direction Method of Multipliers (ADMM)

ADMM repeats the following steps, for  $k = 1, 2, 3, \dots$ :

1. Update  $x$ :

$$x_k = \arg \min_x L_\rho(x, z_{k-1}, u_{k-1})$$

2. Update  $z$ :

$$z_k = \arg \min_z L_\rho(x_k, z, u_{k-1})$$


3. Update  $u$ :

$$u_k = u_{k-1} + \rho(Ax_k + Bz_k - c)$$

**Note:** The usual method of multipliers would replace the first two steps by a joint minimization:

$$(x^{(k)}, z^{(k)}) = \arg \min_{x, z} L_\rho(x, z, u^{(k-1)})$$

## Example: Alternating Projections



Consider finding a point in the intersection of convex sets  $U, V \subseteq \mathbb{R}^n$ :

$$\min_x I_U(x) + I_V(x)$$

To transform this problem into ADMM form, we express it as:

$$\min_{x,z} I_U(x) + I_V(z) \quad \text{subject to} \quad x - z = 0$$

Each ADMM cycle involves two projections:

$$x_k = \arg \min_x P_U(z_{k-1} - w_{k-1})$$

$$z_k = \arg \min_z P_V(x_k + w_{k-1})$$

$$w_k = w_{k-1} + x_k - z_k$$

# Sources

- Ryan Tibshirani. Convex Optimization 10-725