Негладкая выпуклая оптимизация. Нижние оценки. Субградиентный метод.

Даниил Меркулов

Методы оптимизации. МФТИ

Non-smooth problems

ℓ_1-regularized linear least squares

ℓ_1 induces sparsity



@fminxyz

Norms are not smooth

$$\min_{x \in \mathbb{R}^n} f(x),$$

A classical convex optimization problem is considered. We assume that $f(x)$ is a convex function, but now we do not require smoothness.



Рис. 1: Norm cones for different p - norms are non-smooth

Wolfe's example



Рис. 2: Wolfe's example. Open in Colab

Subgradient calculus

Convex function linear lower bound



Рис. 3: Taylor linear approximation serves as a global lower bound for a convex function

An important property of a continuous convex function $f(x)$ is that at any chosen point x_0 for all $x \in \text{dom } f$ the inequality holds:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

Convex function linear lower bound



Рис. 3: Taylor linear approximation serves as a global lower bound for a convex function

An important property of a continuous convex function $f(x)$ is that at any chosen point x_0 for all $x \in \text{dom } f$ the inequality holds:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

for some vector g, i.e., the tangent to the function's graph is the *global* estimate from below for the function.

- If $f(x)$ is differentiable, then $g = \nabla f(x_0)$

Convex function linear lower bound



Рис. 3: Taylor linear approximation serves as a global lower bound for a convex function

An important property of a continuous convex function $f(x)$ is that at any chosen point x_0 for all $x \in \text{dom } f$ the inequality holds:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

for some vector g, i.e., the tangent to the function's graph is the *global* estimate from below for the function.

- If $f(x)$ is differentiable, then $g = \nabla f(x_0)$
- Not all continuous convex functions are differentiable.

Convex function linear lower bound



Рис. 3: Taylor linear approximation serves as a global lower bound for a convex function

An important property of a continuous convex function $f(x)$ is that at any chosen point x_0 for all $x \in \text{dom } f$ the inequality holds:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

for some vector g, i.e., the tangent to the function's graph is the *global* estimate from below for the function.

- If $f(x)$ is differentiable, then $g = \nabla f(x_0)$
- Not all continuous convex functions are differentiable.

Convex function linear lower bound



Рис. 3: Taylor linear approximation serves as a global lower bound for a convex function

An important property of a continuous convex function $f(x)$ is that at any chosen point x_0 for all $x \in \text{dom } f$ the inequality holds:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

for some vector g, i.e., the tangent to the function's graph is the *global* estimate from below for the function.

- If $f(x)$ is differentiable, then $g = \nabla f(x_0)$
- Not all continuous convex functions are differentiable.

We do not want to lose such a lovely property.

Subgradient and subdifferential

A vector g is called the **subgradient** of a function $f(x) : S \to \mathbb{R}$ at a point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

Subgradient and subdifferential

A vector g is called the **subgradient** of a function $f(x) : S \to \mathbb{R}$ at a point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

The set of all subgradients of a function $f(x)$ at a point x_0 is called the **subdifferential** of f at x_0 and is denoted by $\partial f(x_0)$.

Subgradient and subdifferential

A vector g is called the **subgradient** of a function $f(x) : S \to \mathbb{R}$ at a point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

The set of all subgradients of a function $f(x)$ at a point x_0 is called the **subdifferential** of f at x_0 and is denoted by $\partial f(x_0)$.



Рис. 4: Subdifferential is a set of all possible subgradients

Subgradient and subdifferential

Find $\partial f(x)$, if $f(x) = |x|$

Subgradient and subdifferential

Find $\partial f(x)$, if $f(x) = |x|$





Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

Subdifferential of a differentiable function

Let $f : S \to \mathbb{R}$ be a function defined on the set S in a Euclidean space $\mathbb{R}^n$. If $x_0 \in \mathbf{ri}(S)$ and f is differentiable at x_0, then either $\partial f(x_0) = \emptyset$ or $\partial f(x_0) = \{\nabla f(x_0)\}$. Moreover, if the function f is convex, the first scenario is impossible.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

Subdifferential of a differentiable function

Let $f : S \to \mathbb{R}$ be a function defined on the set S in a Euclidean space $\mathbb{R}^n$. If $x_0 \in \mathbf{ri}(S)$ and f is differentiable at x_0, then either $\partial f(x_0) = \emptyset$ or $\partial f(x_0) = \{\nabla f(x_0)\}$. Moreover, if the function f is convex, the first scenario is impossible.

Proof

1. Assume, that $s \in \partial f(x_0)$ for some $s \in \mathbb{R}^n$ distinct from $\nabla f(x_0)$. Let $v \in \mathbb{R}^n$ be a unit vector. Because x_0 is an interior point of S, there exists $\delta > 0$ such that $x_0 + tv \in S$ for all $0 < t < \delta$. By the definition of the subgradient, we have

$$f(x_0 + tv) \geq f(x_0) + t\langle s, v\rangle$$

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

Subdifferential of a differentiable function

Let $f : S \to \mathbb{R}$ be a function defined on the set S in a Euclidean space $\mathbb{R}^n$. If $x_0 \in \mathbf{ri}(S)$ and f is differentiable at x_0, then either $\partial f(x_0) = \emptyset$ or $\partial f(x_0) = \{\nabla f(x_0)\}$. Moreover, if the function f is convex, the first scenario is impossible.

Proof

1. Assume, that $s \in \partial f(x_0)$ for some $s \in \mathbb{R}^n$ distinct from $\nabla f(x_0)$. Let $v \in \mathbb{R}^n$ be a unit vector. Because x_0 is an interior point of S, there exists $\delta > 0$ such that $x_0 + tv \in S$ for all $0 < t < \delta$. By the definition of the subgradient, we have

$$f(x_0 + tv) \geq f(x_0) + t\langle s, v\rangle$$

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

> **Subdifferential of a differentiable function**
>
> Let $f : S \to \mathbb{R}$ be a function defined on the set S in a Euclidean space $\mathbb{R}^n$. If $x_0 \in \mathbf{ri}(S)$ and f is differentiable at x_0, then either $\partial f(x_0) = \emptyset$ or $\partial f(x_0) = \{\nabla f(x_0)\}$. Moreover, if the function f is convex, the first scenario is impossible.

Proof

1. Assume, that $s \in \partial f(x_0)$ for some $s \in \mathbb{R}^n$ distinct from $\nabla f(x_0)$. Let $v \in \mathbb{R}^n$ be a unit vector. Because x_0 is an interior point of S, there exists $\delta > 0$ such that $x_0 + tv \in S$ for all $0 < t < \delta$. By the definition of the subgradient, we have

$$f(x_0 + tv) \geq f(x_0) + t\langle s, v\rangle$$

which implies:

$$\frac{f(x_0 + tv) - f(x_0)}{t} \geq \langle s, v\rangle$$

for all $0 < t < \delta$. Taking the limit as t approaches 0 and using the definition of the gradient, we get:

$$\langle \nabla f(x_0), v\rangle = \lim_{t \to 0; 0 < t < \delta} \frac{f(x_0 + tv) - f(x_0)}{t} \geq \langle s, v\rangle$$

2. From this, $\langle s - \nabla f(x_0), v\rangle \geq 0$. Due to the arbitrariness of v, one can set

$$v = -\frac{s - \nabla f(x_0)}{\|s - \nabla f(x_0)\|},$$

leading to $s = \nabla f(x_0)$.

Subdifferential properties

- If $x_0 \in \mathbf{ri}(S)$, then $\partial f(x_0)$ is a convex compact set.
- The convex function $f(x)$ is differentiable at the point $x_0 \Rightarrow \partial f(x_0) = \{\nabla f(x_0)\}$.
- If $\partial f(x_0) \neq \emptyset \quad \forall x_0 \in S$, then $f(x)$ is convex on S.

> **Subdifferential of a differentiable function**
>
> Let $f : S \to \mathbb{R}$ be a function defined on the set S in a Euclidean space $\mathbb{R}^n$. If $x_0 \in \mathbf{ri}(S)$ and f is differentiable at x_0, then either $\partial f(x_0) = \emptyset$ or $\partial f(x_0) = \{\nabla f(x_0)\}$. Moreover, if the function f is convex, the first scenario is impossible.

Proof

1. Assume, that $s \in \partial f(x_0)$ for some $s \in \mathbb{R}^n$ distinct from $\nabla f(x_0)$. Let $v \in \mathbb{R}^n$ be a unit vector. Because x_0 is an interior point of S, there exists $\delta > 0$ such that $x_0 + tv \in S$ for all $0 < t < \delta$. By the definition of the subgradient, we have

$$f(x_0 + tv) \geq f(x_0) + t\langle s, v\rangle$$

which implies:

$$\frac{f(x_0 + tv) - f(x_0)}{t} \geq \langle s, v\rangle$$

for all $0 < t < \delta$. Taking the limit as t approaches 0 and using the definition of the gradient, we get:

$$\langle \nabla f(x_0), v\rangle = \lim_{t \to 0; 0 < t < \delta} \frac{f(x_0 + tv) - f(x_0)}{t} \geq \langle s, v\rangle$$

2. From this, $\langle s - \nabla f(x_0), v\rangle \geq 0$. Due to the arbitrariness of v, one can set

$$v = -\frac{s - \nabla f(x_0)}{\|s - \nabla f(x_0)\|},$$

leading to $s = \nabla f(x_0)$.

3. Furthermore, if the function f is convex, then according to the differential condition of convexity $f(x) \geq f(x_0) + \langle \nabla f(x_0), x - x_0\rangle$ for all $x \in S$. But by definition, this means $\nabla f(x_0) \in \partial f(x_0)$.

Subdifferential calculus

Moreau - Rockafellar theorem (subdifferential of a linear combination)

Let $f_i(x)$ be convex functions on convex sets $S_i, \ i = \overline{1, n}$. Then if $\bigcap\limits_{i=1}^{n} \mathbf{ri}(S_i) \neq \emptyset$ then the function $f(x) = \sum\limits_{i=1}^{n} a_i f_i(x), \ a_i > 0$ has a subdifferential $\partial_S f(x)$ on the set $S = \bigcap\limits_{i=1}^{n} S_i$ and

$$\partial_S f(x) = \sum_{i=1}^{n} a_i \partial_{S_i} f_i(x)$$

Subdifferential calculus

Moreau - Rockafellar theorem (subdifferential of a linear combination)

Let $f_i(x)$ be convex functions on convex sets S_i, $i = \overline{1, n}$. Then if $\bigcap\limits_{i=1}^{n} \mathbf{ri}(S_i) \neq \emptyset$ then the function $f(x) = \sum\limits_{i=1}^{n} a_i f_i(x)$, $a_i > 0$ has a subdifferential $\partial_S f(x)$ on the set $S = \bigcap\limits_{i=1}^{n} S_i$ and

$$\partial_S f(x) = \sum_{i=1}^{n} a_i \partial_{S_i} f_i(x)$$

Dubovitsky - Milutin theorem (subdifferential of a point-wise maximum)

Let $f_i(x)$ be convex functions on the open convex set $S \subseteq \mathbb{R}^n$, $x_0 \in S$, and the pointwise maximum is defined as $f(x) = \max\limits_i f_i(x)$. Then:

$$\partial_S f(x_0) = \mathbf{conv} \left\{ \bigcup_{i \in I(x_0)} \partial_S f_i(x_0) \right\}, \quad I(x) = \{i \in [1$$

Subdifferential calculus

- $\partial(\alpha f)(x) = \alpha \partial f(x)$, for $\alpha \geq 0$

Subdifferential calculus

- $\partial(\alpha f)(x) = \alpha \partial f(x)$, for $\alpha \geq 0$
- $\partial(\sum f_i)(x) = \sum \partial f_i(x)$, f_i - convex functions

Subdifferential calculus

- $\partial(\alpha f)(x) = \alpha \partial f(x)$, for $\alpha \geq 0$
- $\partial(\sum f_i)(x) = \sum \partial f_i(x)$, f_i - convex functions
- $\partial(f(Ax+b))(x) = A^T \partial f(Ax+b)$, f - convex function

Subdifferential calculus

- $\partial(\alpha f)(x) = \alpha \partial f(x)$, for $\alpha \geq 0$
- $\partial(\sum f_i)(x) = \sum \partial f_i(x)$, f_i - convex functions
- $\partial(f(Ax+b))(x) = A^T \partial f(Ax+b)$, f - convex function
- $z \in \partial f(x)$ if and only if $x \in \partial f^*(z)$.

Subgradient Method

Algorithm

A vector g is called the **subgradient** of the function $f(x) : S \to \mathbb{R}$ at the point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

Algorithm

A vector g is called the **subgradient** of the function $f(x) : S \to \mathbb{R}$ at the point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

The idea is very simple: let's replace the gradient $\nabla f(x_k)$ in the gradient descent algorithm with a subgradient g_k at point x_k:

$$x_{k+1} = x_k - \alpha_k g_k,$$

where g_k is an arbitrary subgradient of the function $f(x)$ at the point x_k, $g_k \in \partial f(x_k)$

Algorithm

A vector g is called the **subgradient** of the function $f(x) : S \to \mathbb{R}$ at the point x_0 if $\forall x \in S$:

$$f(x) \geq f(x_0) + \langle g, x - x_0 \rangle$$

The idea is very simple: let's replace the gradient $\nabla f(x_k)$ in the gradient descent algorithm with a subgradient g_k at point x_k:

$$x_{k+1} = x_k - \alpha_k g_k,$$

where g_k is an arbitrary subgradient of the function $f(x)$ at the point x_k, $g_k \in \partial f(x_k)$

Note that the **subgradient method is not guaranteed to be a descent method**; the negative subgradient need not be a descent direction, or the step size may cause $f(x_{k+1}) > f(x_k)$.

That is why we usually track the best value of the objective function

$$f_k^{\text{best}} = \min_{i=1,\ldots,k} f(x_i).$$

Convergence bound

$$\|x_{k+1} - x^*\|^2 = \|x_k - x^* - \alpha_k g_k\|^2 =$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 = \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*))
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 = \|x_k - x^* - \alpha_k g_k\|^2 = \\
= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*))
\end{aligned}
$$

$$
2\alpha_k (f(x_k) - f(x^*)) \leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) \leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 = \|x_k - x^* - \alpha_k g_k\|^2 = \\
= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) \leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq R^2 + G^2 \sum_{k=0}^{T-1} \alpha_k^2
\end{aligned}
$$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq R^2 + G^2 \sum_{k=0}^{T-1} \alpha_k^2
\end{aligned}
$$

- Let's write down how close we came to the optimum $x^* = \arg\min_{x \in \mathbb{R}^n} f(x) = \arg f^*$ on the last iteration:

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq R^2 + G^2 \sum_{k=0}^{T-1} \alpha_k^2
\end{aligned}
$$

- Let's write down how close we came to the optimum $x^* = \arg\min_{x \in \mathbb{R}^n} f(x) = \arg f^*$ on the last iteration:
- For a subgradient: $\langle g_k, x^* - x_k \rangle \leq f(x^*) - f(x_k)$.

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) \\
2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq R^2 + G^2 \sum_{k=0}^{T-1} \alpha_k^2
\end{aligned}
$$

- Let's write down how close we came to the optimum $x^* = \arg\min_{x \in \mathbb{R}^n} f(x) = \arg f^*$ on the last iteration:
- For a subgradient: $\langle g_k, x^* - x_k \rangle \leq f(x^*) - f(x_k)$.
- We additionally assume that $\|g_k\|^2 \leq G^2$

Convergence bound

$$
\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*))
\end{aligned}
$$

$$
2\alpha_k (f(x_k) - f(x^*)) \leq \|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
$$

Let us sum the obtained inequality for $k = 0, \ldots, T-1$:

$$
\begin{aligned}
\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) &\leq \|x_0 - x^*\|^2 - \|x_T - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq \|x_0 - x^*\|^2 + \sum_{k=0}^{T-1} \alpha_k^2 \|g_k\|^2 \\
&\leq R^2 + G^2 \sum_{k=0}^{T-1} \alpha_k^2
\end{aligned}
$$

- Let's write down how close we came to the optimum $x^* = \arg\min_{x \in \mathbb{R}^n} f(x) = \arg f^*$ on the last iteration:
- For a subgradient: $\langle g_k, x^* - x_k \rangle \leq f(x^*) - f(x_k)$.
- We additionally assume that $\|g_k\|^2 \leq G^2$
- We use the notation $R = \|x_0 - x^*\|_2$

Convergence bound

- Finally, note:

$$\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) \geq \sum_{k=0}^{T-1} 2\alpha_k (f_k^{\text{best}} - f(x^*)) = (f_k^{\text{best}} - f(x^*)) \sum_{k=0}^{T-1} 2\alpha_k$$

Convergence bound

- Finally, note:

$$\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) \geq \sum_{k=0}^{T-1} 2\alpha_k (f_k^{\text{best}} - f(x^*)) = (f_k^{\text{best}} - f(x^*)) \sum_{k=0}^{T-1} 2\alpha_k$$

- Which leads to the basic inequality:

$$\boxed{f_k^{\text{best}} - f(x^*) \leq \frac{R^2 + G^2 \sum\limits_{k=0}^{T-1} \alpha_k^2}{2 \sum\limits_{k=0}^{T-1} \alpha_k}}$$

Convergence bound

- Finally, note:

$$\sum_{k=0}^{T-1} 2\alpha_k (f(x_k) - f(x^*)) \geq \sum_{k=0}^{T-1} 2\alpha_k (f_k^{\text{best}} - f(x^*)) = (f_k^{\text{best}} - f(x^*)) \sum_{k=0}^{T-1} 2\alpha_k$$

- Which leads to the basic inequality:

$$\boxed{f_k^{\text{best}} - f(x^*) \leq \frac{R^2 + G^2 \sum\limits_{k=0}^{T-1} \alpha_k^2}{2 \sum\limits_{k=0}^{T-1} \alpha_k}}$$

- From this point we can see, that if the stepsize strategy is such that

$$\sum_{k=0}^{T-1} \alpha_k^2 < \infty, \quad \sum_{k=0}^{T-1} \alpha_k = \infty,$$

then the subgradient method converges (step size should be decreasing, but not too fast).

Different step size strategies



Different step size strategies



Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size α, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{R^2}{2\alpha k} + \frac{\alpha}{2} G^2$$

- Note, that with any constant step size, the first term of the right-hand side is decreasing, but the second term stays constant.

Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size α, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{R^2}{2\alpha k} + \frac{\alpha}{2} G^2$$

- Note, that with any constant step size, the first term of the right-hand side is decreasing, but the second term stays constant.
- Some versions of the subgradient method (e.g., diminishing nonsummable step lengths) work when the assumption on $\|g_k\|_2 \leq G$ doesn't hold; see [1] or [2].

[1] B. Polyak. Introduction to Optimization. Optimization Software, Inc., 1987.
[2] N. Shor. Minimization Methods for Non-differentiable Functions. Springer Series in Computational Mathematics. Springer, 1985.

Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size α, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{R^2}{2\alpha k} + \frac{\alpha}{2} G^2$$

- Note, that with any constant step size, the first term of the right-hand side is decreasing, but the second term stays constant.
- Some versions of the subgradient method (e.g., diminishing nonsummable step lengths) work when the assumption on $\|g_k\|_2 \leq G$ doesn't hold; see [1] or [2].
- Let's find the optimal step size α that minimizes the right-hand side of the inequality.

[1] B. Polyak. Introduction to Optimization. Optimization Software, Inc., 1987.
[2] N. Shor. Minimization Methods for Non-differentiable Functions. Springer Series in Computational Mathematics. Springer, 1985.

Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size $\alpha = \frac{R}{G}\sqrt{\frac{1}{k}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR}{\sqrt{k}}$$

- This version requires knowledge of the number of iterations in advance, which is not usually practical.

Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size $\alpha = \frac{R}{G}\sqrt{\frac{1}{k}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR}{\sqrt{k}}$$

- This version requires knowledge of the number of iterations in advance, which is not usually practical.
- It is interesting to mention, that if you want to find the optimal stepsizes for the whole sequence $\alpha_0, \alpha_1, \dots, \alpha_{k-1}$, you will get the same result.

Convergence bound. Non-smooth convex case. Constant step size

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step size $\alpha = \frac{R}{G}\sqrt{\frac{1}{k}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR}{\sqrt{k}}$$

- This version requires knowledge of the number of iterations in advance, which is not usually practical.
- It is interesting to mention, that if you want to find the optimal stepsizes for the whole sequence $\alpha_0, \alpha_1, \ldots, \alpha_{k-1}$, you will get the same result.
- Why? Because the right-hand side is convex and **symmetric** function of $\alpha_0, \alpha_1, \ldots, \alpha_{k-1}$.

Convergence bound. Non-smooth convex case. Constant step length

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a fixed step length $\gamma = \alpha_k \|g_k\|_2$, i.e. $\alpha_k = \frac{\gamma}{\|g_k\|_2}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR^2}{2\gamma k} + \frac{G\gamma}{2}$$

- Note, that for the subgradient method, we typically can not use the norm of the subgradient as a stopping criterion (imagine $f(x) = |x|$). There are some variants of more advanced stopping criteria, but the convergence is so slow, so typically we just set a maximum number of iterations.

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T);$$

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T); \qquad \sum_{k=0}^{T-1} \alpha_k = \frac{R}{G} \sum_{k=1}^{T} \frac{1}{\sqrt{k}} \geq \frac{R}{G} \int_1^{T+1} \frac{1}{\sqrt{t}}\, dt = \frac{2R}{G}(\sqrt{T+1} - 1).$$

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T); \qquad \sum_{k=0}^{T-1} \alpha_k = \frac{R}{G} \sum_{k=1}^{T} \frac{1}{\sqrt{k}} \geq \frac{R}{G} \int_1^{T+1} \frac{1}{\sqrt{t}}\, dt = \frac{2R}{G}(\sqrt{T+1} - 1).$$

2. We drop the last -1 in the upper bound above and use the basic inequality:

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T); \qquad \sum_{k=0}^{T-1} \alpha_k = \frac{R}{G} \sum_{k=1}^{T} \frac{1}{\sqrt{k}} \geq \frac{R}{G} \int_1^{T+1} \frac{1}{\sqrt{t}}\, dt = \frac{2R}{G}(\sqrt{T+1} - 1).$$

2. We drop the last -1 in the upper bound above and use the basic inequality:

$$f_T^{\text{best}} - f(x^*) \leq \frac{R^2 + G^2 \sum\limits_{k=0}^{T-1} \alpha_k^2}{2 \sum\limits_{k=0}^{T-1} \alpha_k}$$

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T); \qquad \sum_{k=0}^{T-1} \alpha_k = \frac{R}{G} \sum_{k=1}^{T} \frac{1}{\sqrt{k}} \geq \frac{R}{G} \int_1^{T+1} \frac{1}{\sqrt{t}}\, dt = \frac{2R}{G}(\sqrt{T+1} - 1).$$

2. We drop the last -1 in the upper bound above and use the basic inequality:

$$f_T^{\text{best}} - f(x^*) \leq \frac{R^2 + G^2 \sum\limits_{k=0}^{T-1} \alpha_k^2}{2 \sum\limits_{k=0}^{T-1} \alpha_k} \leq \frac{R^2 + R^2(1 + \ln T)}{4\frac{R}{G}(\sqrt{T+1})}$$

Convergence bound. Non-smooth convex case. Practical strategy

Theorem

Let f be a convex G-Lipschitz function and $R = \|x_0 - x^*\|_2$. For a diminishing step size strategy $\alpha_k = \frac{R}{G\sqrt{k+1}}$, subgradient method satisfies

$$f_k^{\text{best}} - f(x^*) \leq \frac{GR(2 + \ln k)}{4\sqrt{k+1}}$$

1. Bounding sums:

$$\sum_{k=0}^{T-1} \alpha_k^2 = \frac{R^2}{G^2} \sum_{k=1}^{T} \frac{1}{k} \leq \frac{R^2}{G^2}(1 + \ln T); \qquad \sum_{k=0}^{T-1} \alpha_k = \frac{R}{G} \sum_{k=1}^{T} \frac{1}{\sqrt{k}} \geq \frac{R}{G} \int_1^{T+1} \frac{1}{\sqrt{t}}\, dt = \frac{2R}{G}(\sqrt{T+1} - 1).$$

2. We drop the last -1 in the upper bound above and use the basic inequality:

$$f_T^{\text{best}} - f(x^*) \leq \frac{R^2 + G^2 \sum\limits_{k=0}^{T-1} \alpha_k^2}{2 \sum\limits_{k=0}^{T-1} \alpha_k} \leq \frac{R^2 + R^2(1 + \ln T)}{4\frac{R}{G}(\sqrt{T+1})} = \frac{GR(2 + \ln T)}{4\sqrt{T+1}}$$

Non-smooth strongly convex case



Non-smooth
Convex



Non-smooth
μ - strongly convex

Non-smooth strongly convex case



Non-smooth
Convex

$$\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$$



Non-smooth
μ - strongly convex

$$\mathcal{O}\left(\frac{1}{k}\right)$$

Non-smooth strongly convex case

Theorem

Let f be μ-strongly convex on a convex set and x, y be arbitrary points. Then for any $g \in \partial f(x)$,

$$\langle g, x - y \rangle \geq f(x) - f(y) + \frac{\mu}{2}\|x - y\|^2.$$

1. For any $\lambda \in [0, 1)$, by μ-strong convexity,

$$f(\lambda x + (1 - \lambda)y) \leq \lambda f(x) + (1 - \lambda)f(y) - \frac{\mu}{2}\lambda(1 - \lambda)\|x - y\|^2.$$

Non-smooth strongly convex case

Theorem

Let f be μ-strongly convex on a convex set and x, y be arbitrary points. Then for any $g \in \partial f(x)$,

$$\langle g, x - y \rangle \geq f(x) - f(y) + \frac{\mu}{2}\|x - y\|^2.$$

1. For any $\lambda \in [0, 1)$, by μ-strong convexity,

$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y) - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2.$$

2. By the subgradient inequality at x, we have

$$f(\lambda x + (1-\lambda)y) \geq f(x) + \langle g, \lambda x + (1-\lambda)y - x \rangle \quad \to \quad f(\lambda x + (1-\lambda)y) \geq f(x) - (1-\lambda)\langle g, x - y \rangle.$$

Non-smooth strongly convex case

Theorem

Let f be μ-strongly convex on a convex set and x, y be arbitrary points. Then for any $g \in \partial f(x)$,

$$\langle g, x - y \rangle \geq f(x) - f(y) + \frac{\mu}{2}\|x - y\|^2.$$

1. For any $\lambda \in [0, 1)$, by μ-strong convexity,

$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y) - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2.$$

2. By the subgradient inequality at x, we have

$$f(\lambda x + (1-\lambda)y) \geq f(x) + \langle g, \lambda x + (1-\lambda)y - x \rangle \quad \rightarrow \quad f(\lambda x + (1-\lambda)y) \geq f(x) - (1-\lambda)\langle g, x - y \rangle.$$

3. Thus,

$$f(x) - (1-\lambda)\langle g, x - y \rangle \leq \lambda f(x) + (1-\lambda)f(y) - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2$$

$$(1-\lambda)f(x) \leq (1-\lambda)f(y) + (1-\lambda)\langle g, x - y \rangle - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2$$

$$f(x) \leq f(y) + \langle g, x - y \rangle - \frac{\mu}{2}\lambda\|x - y\|^2$$

Non-smooth strongly convex case

Theorem

Let f be μ-strongly convex on a convex set and x, y be arbitrary points. Then for any $g \in \partial f(x)$,

$$\langle g, x - y \rangle \geq f(x) - f(y) + \frac{\mu}{2}\|x - y\|^2.$$

1. For any $\lambda \in [0, 1)$, by μ-strong convexity,
$$f(\lambda x + (1-\lambda)y) \leq \lambda f(x) + (1-\lambda)f(y) - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2.$$
2. By the subgradient inequality at x, we have
$$f(\lambda x + (1-\lambda)y) \geq f(x) + \langle g, \lambda x + (1-\lambda)y - x \rangle \quad \to \quad f(\lambda x + (1-\lambda)y) \geq f(x) - (1-\lambda)\langle g, x - y \rangle.$$
3. Thus,
$$f(x) - (1-\lambda)\langle g, x - y \rangle \leq \lambda f(x) + (1-\lambda)f(y) - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2$$
$$(1-\lambda)f(x) \leq (1-\lambda)f(y) + (1-\lambda)\langle g, x - y \rangle - \frac{\mu}{2}\lambda(1-\lambda)\|x - y\|^2$$
$$f(x) \leq f(y) + \langle g, x - y \rangle - \frac{\mu}{2}\lambda\|x - y\|^2$$
4. Letting $\lambda \to 1^-$ gives $f(x) \leq f(y) + \langle g, x - y \rangle - \frac{\mu}{2}\|x - y\|^2 \to \langle g, x - y \rangle \geq f(x) - f(y) + \frac{\mu}{2}\|x - y\|^2$.

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\|x_{k+1} - x^*\|^2 = \|x_k - x^* - \alpha_k g_k\|^2 =$$

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\begin{aligned} \|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\ &= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \end{aligned}$$

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) - \alpha_k \mu \|x_k - x^*\|^2
\end{aligned}$$

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) - \alpha_k \mu \|x_k - x^*\|^2 \\
&= (1 - \mu\alpha_k)\|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \left(f(x_k) - f(x^*)\right)
\end{aligned}$$

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) - \alpha_k \mu \|x_k - x^*\|^2 \\
&= (1 - \mu\alpha_k)\|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \left(f(x_k) - f(x^*)\right) \\
2\alpha_k \left(f(x_k) - f(x^*)\right) &\leq (1 - \mu\alpha_k)\|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2
\end{aligned}$$

Convergence bound. Non-smooth strongly convex case.

Theorem

Let f be a μ-strongly convex function (possibly non-smooth) with minimizer x^* and bounded subgradients $\|g_k\| \leq G$. Using the step size $\alpha_k = \frac{2}{\mu(k+1)}$, the subgradient method guarantees for $k > 0$ that:

$$f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.$$

1. We start with the method formulation as before:

$$\begin{aligned}
\|x_{k+1} - x^*\|^2 &= \|x_k - x^* - \alpha_k g_k\|^2 = \\
&= \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \langle g_k, x_k - x^* \rangle \\
&\leq \|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k (f(x_k) - f(x^*)) - \alpha_k \mu \|x_k - x^*\|^2 \\
&= (1 - \mu\alpha_k)\|x_k - x^*\|^2 + \alpha_k^2 \|g_k\|^2 - 2\alpha_k \left(f(x_k) - f(x^*)\right) \\
2\alpha_k \left(f(x_k) - f(x^*)\right) &\leq (1 - \mu\alpha_k)\|x_k - x^*\|^2 - \|x_{k+1} - x^*\|^2 + \alpha_k^2 \|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{1 - \mu\alpha_k}{2\alpha_k} \|x_k - x^*\|^2 - \frac{1}{2\alpha_k} \|x_{k+1} - x^*\|^2 + \frac{\alpha_k}{2} \|g_k\|^2
\end{aligned}$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$
\begin{aligned}
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2 \\
k\left(f(x_k) - f(x^*)\right) &\leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2
\end{aligned}
$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$
\begin{aligned}
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2 \\
k\left(f(x_k) - f(x^*)\right) &\leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2
\end{aligned}
$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$
\begin{aligned}
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2 \\
k\left(f(x_k) - f(x^*)\right) &\leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2
\end{aligned}
$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$
\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2
$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$
\begin{aligned}
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2 \\
k\left(f(x_k) - f(x^*)\right) &\leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2
\end{aligned}
$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$
\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}
$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

$$k\left(f(x_k) - f(x^*)\right) \leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}$$

$$\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right)$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

$$k\left(f(x_k) - f(x^*)\right) \leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}$$

$$\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right) \leq \sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right)$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

$$k\left(f(x_k) - f(x^*)\right) \leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}$$

$$\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right) \leq \sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq \frac{G^2 T}{\mu}$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

$$k\left(f(x_k) - f(x^*)\right) \leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}$$

$$\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right) \leq \sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq \frac{G^2 T}{\mu}$$

$$f_{T-1}^{\text{best}} - f(x^*) \leq \frac{G^2 T}{\mu \sum_{k=0}^{T-1} k}$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2$$

$$f(x_k) - f(x^*) \leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2$$

$$k\left(f(x_k) - f(x^*)\right) \leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}$$

$$\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right) \leq \sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq \frac{G^2 T}{\mu}$$

$$f_{T-1}^{\text{best}} - f(x^*) \leq \frac{G^2 T}{\mu \sum_{k=0}^{T-1} k} = \frac{2G^2 T}{\mu T(T-1)}$$

Convergence bound. Non-smooth strongly convex case. Proof

2. Substitute the step size $\alpha_k = \frac{2}{\mu(k+1)}$ into the inequality:

$$
\begin{aligned}
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu(k+1)}\|g_k\|^2 \\
f(x_k) - f(x^*) &\leq \frac{\mu(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu k}\|g_k\|^2 \\
k\left(f(x_k) - f(x^*)\right) &\leq \frac{\mu k(k-1)}{4}\|x_k - x^*\|^2 - \frac{\mu k(k+1)}{4}\|x_{k+1} - x^*\|^2 + \frac{1}{\mu}\|g_k\|^2
\end{aligned}
$$

3. Summing up the inequalities for all $k = 0, 1, \ldots, T-1$, we get:

$$
\sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq 0 - \frac{\mu(T-1)T}{4}\|x_T - x^*\|^2 + \frac{1}{\mu}\sum_{k=0}^{T-1}\|g_k\|^2 \leq \frac{G^2 T}{\mu}
$$

$$
\left(f_{T-1}^{\text{best}} - f(x^*)\right)\sum_{k=0}^{T-1} k = \sum_{k=0}^{T-1} k\left(f_{T-1}^{\text{best}} - f(x^*)\right) \leq \sum_{k=0}^{T-1} k\left(f(x_k) - f(x^*)\right) \leq \frac{G^2 T}{\mu}
$$

$$
f_{T-1}^{\text{best}} - f(x^*) \leq \frac{G^2 T}{\mu \sum_{k=0}^{T-1} k} = \frac{2G^2 T}{\mu T(T-1)} \qquad f_k^{\text{best}} - f(x^*) \leq \frac{2G^2}{\mu k}.
$$

Summary. Subgradient method

Problem Type	Stepsize Rule	Convergence Rate	Iteration Complexity
Convex & Lipschitz problems	$\alpha \sim \frac{1}{\sqrt{k}}$	$\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$	$\mathcal{O}\left(\frac{1}{\varepsilon^2}\right)$
Strongly convex & Lipschitz problems	$\alpha \sim \frac{1}{k}$	$\mathcal{O}\left(\frac{1}{k}\right)$	$\mathcal{O}\left(\frac{1}{\varepsilon}\right)$

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\tfrac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 6: Smooth convex case. Sublinear convergence, no convergence in domain

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A \in \mathbb{R}^{m\times n}, \quad \lambda\left(\tfrac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 7: Non-smooth convex case. Small λ value imposes non-smoothness. No convergence with constant step size

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\tfrac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 8: Non-smooth convex case. Larger λ value reveals non-monotonicity of $f(x_k)$. One can see that a smaller constant step size leads to a lower stationary level.

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\tfrac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 9: Non-smooth convex case. Diminishing step size leads to the convergence fot the f_k^{best}

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\frac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 10: Non-smooth convex case. $\frac{\alpha_0}{\sqrt{k}}$ step size leads to the convergence fot the f_k^{best}

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A \in \mathbb{R}^{m\times n}, \quad \lambda\left(\tfrac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 11: Non-smooth convex case. $\frac{\alpha_0}{\sqrt{k}}$ step size leads to the convergence fot the f_k^{best}

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\frac{1}{m}A^TA\right) \in [\mu; L].$$



Рис. 12: Non-smooth strongly convex case. $\frac{\alpha_0}{k}$ step size leads to the convergence fot the f_k^{best}

Numerical experiments

$$f(x) = \frac{1}{2m}\|Ax - b\|_2^2 + \lambda\|x\|_1 \to \min_{x \in \mathbb{R}^n}, \qquad A \in \mathbb{R}^{m \times n}, \quad \lambda\left(\tfrac{1}{m}A^T A\right) \in [\mu; L].$$



Рис. 13: Non-smooth strongly convex case. $\frac{\alpha_0}{\sqrt{k}}$ step size works worse

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 14: Logistic regression with ℓ_1 regularization

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 15: Logistic regression with ℓ_1 regularization

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 16: Logistic regression with ℓ_1 regularization

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 17: Logistic regression with ℓ_1 regularization

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 18: Logistic regression with ℓ_1 regularization

Numerical experiments

$$f(x) = \frac{1}{m}\sum_{i=1}^{m} \log(1 + \exp(-b_i(A_i x))) + \lambda\|x\|_1 \to \min_{x\in\mathbb{R}^n}, \qquad A_i \in \mathbb{R}^n, \quad b_i \in \{-1, 1\}$$



Рис. 19: Logistic regression with ℓ_1 regularization

Lower bounds

Lower bounds

convex (non-smooth) [3]	smooth (non-convex)[4]	smooth & convex[5]	smooth & strongly convex (or PL)[1]
$\mathcal{O}\left(\frac{1}{\sqrt{k}}\right)$	$\mathcal{O}\left(\frac{1}{k^2}\right)$	$\mathcal{O}\left(\frac{1}{k^2}\right)$	$\mathcal{O}\left(\left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}\right)^k\right)$
$k_\varepsilon \sim \mathcal{O}\left(\frac{1}{\varepsilon^2}\right)$	$k_\varepsilon \sim \mathcal{O}\left(\frac{1}{\sqrt{\varepsilon}}\right)$	$k_\varepsilon \sim \mathcal{O}\left(\frac{1}{\sqrt{\varepsilon}}\right)$	$k_\varepsilon \sim \mathcal{O}\left(\sqrt{\kappa}\log\frac{1}{\varepsilon}\right)$

[3] Nesterov, Lectures on Convex Optimization
[4] Carmon, Duchi, Hinder, Sidford, 2017
[5] Nemirovski, Yudin, 1979

Black box iteration

The iteration of gradient descent:

$$
\begin{aligned}
x^{k+1} &= x^k - \alpha^k \nabla f(x^k) \\
&= x^{k-1} - \alpha^{k-1} \nabla f(x^{k-1}) - \alpha^k \nabla f(x^k) \\
&\vdots \\
&= x^0 - \sum_{i=0}^{k} \alpha^{k-i} \nabla f(x^{k-i})
\end{aligned}
$$

Black box iteration

The iteration of gradient descent:

$$
\begin{aligned}
x^{k+1} &= x^k - \alpha^k \nabla f(x^k) \\
&= x^{k-1} - \alpha^{k-1} \nabla f(x^{k-1}) - \alpha^k \nabla f(x^k) \\
&\vdots \\
&= x^0 - \sum_{i=0}^{k} \alpha^{k-i} \nabla f(x^{k-i})
\end{aligned}
$$

Consider a family of first-order methods, where

$$
\begin{aligned}
&x^{k+1} \in x^0 + \text{span}\left\{\nabla f(x^0), \nabla f(x^1), \ldots, \nabla f(x^k)\right\} && f \text{ - smooth} \\
&x^{k+1} \in x^0 + \text{span}\left\{g_0, g_1, \ldots, g_k\right\}, \text{ where } g_i \in \partial f(x^i) && f \text{ - non-smooth}
\end{aligned} \tag{1}
$$

Black box iteration

The iteration of gradient descent:

$$
\begin{aligned}
x^{k+1} &= x^k - \alpha^k \nabla f(x^k) \\
&= x^{k-1} - \alpha^{k-1} \nabla f(x^{k-1}) - \alpha^k \nabla f(x^k) \\
&\vdots \\
&= x^0 - \sum_{i=0}^{k} \alpha^{k-i} \nabla f(x^{k-i})
\end{aligned}
$$

Consider a family of first-order methods, where

$$
\begin{aligned}
&x^{k+1} \in x^0 + \text{span}\left\{\nabla f(x^0), \nabla f(x^1), \dots, \nabla f(x^k)\right\} && f \text{ - smooth} \\
&x^{k+1} \in x^0 + \text{span}\left\{g_0, g_1, \dots, g_k\right\}, \text{ where } g_i \in \partial f(x^i) && f \text{ - non-smooth}
\end{aligned} \tag{1}
$$

To construct a lower bound, we need to find a function f from the corresponding class such that any method from the family 1 will work at least as slowly as the lower bound.

Non-smooth convex case

Theorem

There exists a function f that is G-Lipschitz and convex such that any method 1 satisfies

$$\min_{i \in [1,k]} f(x^i) - \min_{x \in \mathbb{B}(R)} f(x) \geq \frac{GR}{2(1+\sqrt{k})}$$

for $R > 0$ and $k \leq n$, where n is the dimension of the problem.

Non-smooth convex case

Theorem

There exists a function f that is G-Lipschitz and convex such that any method 1 satisfies

$$\min_{i \in [1,k]} f(x^i) - \min_{x \in \mathbb{B}(R)} f(x) \geq \frac{GR}{2(1+\sqrt{k})}$$

for $R > 0$ and $k \leq n$, where n is the dimension of the problem.

Proof idea: build such a function f that, for any method 1, we have

$$\text{span}\{g_0, g_1, \dots, g_k\} \subset \text{span}\{e_1, e_2, \dots, e_i\}$$

where e_i is the i-th standard basis vector. At iteration $k \leq n$, there are at least $n - k$ coordinate of x are 0. This helps us to derive a bound on the error.

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2} \|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2} \|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Key Properties:

- The function $f(x)$ is α-strongly convex due to the quadratic term $\frac{\alpha}{2}\|x\|_2^2$.

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2} \|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Key Properties:

- The function $f(x)$ is α-strongly convex due to the quadratic term $\frac{\alpha}{2}\|x\|_2^2$.
- The function is non-smooth because the first term introduces a non-differentiable point at the maximum coordinate of x.

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2} \|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Key Properties:

- The function $f(x)$ is α-strongly convex due to the quadratic term $\frac{\alpha}{2}\|x\|_2^2$.
- The function is non-smooth because the first term introduces a non-differentiable point at the maximum coordinate of x.

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2}\|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Key Properties:

- The function $f(x)$ is α-strongly convex due to the quadratic term $\frac{\alpha}{2}\|x\|_2^2$.
- The function is non-smooth because the first term introduces a non-differentiable point at the maximum coordinate of x.

Consider the subdifferential of $f(x)$ at x:

$$\begin{aligned}
\partial f(x) &= \partial \left(\beta \max_{i \in [1,k]} x[i] \right) + \partial \left(\frac{\alpha}{2}\|x\|_2^2 \right) \\
&= \beta \partial \left(\max_{i \in [1,k]} x[i] \right) + \alpha x \\
&= \beta \text{conv} \left\{ e_i \mid i : x[i] = \max_j x[j] \right\} + \alpha x
\end{aligned}$$

Non-smooth case (proof)

Consider the function:

$$f(x) = \beta \max_{i \in [1,k]} x[i] + \frac{\alpha}{2}\|x\|_2^2,$$

where $\alpha, \beta \in \mathbb{R}$ are parameters, and $x[1:k]$ denotes the first k components of x.

Key Properties:

- The function $f(x)$ is α-strongly convex due to the quadratic term $\frac{\alpha}{2}\|x\|_2^2$.
- The function is non-smooth because the first term introduces a non-differentiable point at the maximum coordinate of x.

Consider the subdifferential of $f(x)$ at x:

$$\begin{aligned}
\partial f(x) &= \partial \left(\beta \max_{i \in [1,k]} x[i] \right) + \partial \left(\frac{\alpha}{2}\|x\|_2^2 \right) \\
&= \beta \partial \left(\max_{i \in [1,k]} x[i] \right) + \alpha x \\
&= \beta \text{conv} \left\{ e_i \mid i : x[i] = \max_j x[j] \right\} + \alpha x
\end{aligned}$$

It is easy to see, that if $g \in \partial f(x)$ and $\|x\| \leq R$, then

$$\|g\| \leq \alpha R + \beta$$

Thus, f is $\alpha R + \beta$-Lipschitz on $B(R)$.

Non-smooth case (proof)

Next, we describe the first-order oracle for this function. When queried for a subgradient at a point x, the oracle returns

$$\alpha x + \gamma e_i,$$

where i is the *first* coordinate for with $x[i] = \max_{1 \leq j \leq k} x[j]$.

- We ensure that $\|x^0\| \leq R$ by starting from $x^0 = 0$.

Non-smooth case (proof)

Next, we describe the first-order oracle for this function. When queried for a subgradient at a point x, the oracle returns

$$\alpha x + \gamma e_i,$$

where i is the *first* coordinate for with $x[i] = \max_{1 \leq j \leq k} x[j]$.

- We ensure that $\|x^0\| \leq R$ by starting from $x^0 = 0$.
- When the oracle is queried at $x^0 = 0$, it returns e_1. Consequently, x^1 must lie on the line generated by e_1.

Non-smooth case (proof)

Next, we describe the first-order oracle for this function. When queried for a subgradient at a point x, the oracle returns

$$\alpha x + \gamma e_i,$$

where i is the *first* coordinate for with $x[i] = \max_{1 \leq j \leq k} x[j]$.

- We ensure that $\|x^0\| \leq R$ by starting from $x^0 = 0$.
- When the oracle is queried at $x^0 = 0$, it returns e_1. Consequently, x^1 must lie on the line generated by e_1.
- By an induction argument, one shows that for all i, the iterate x^i lies in the linear span of $\{e_1, \dots, e_i\}$. In particular, for $i \leq k$, the $k+1$-th coordinate of x_i is zero and due to the structure of $f(x)$:

$$f(x^i) \geq 0.$$

Non-smooth case (proof)

- It remains to compute the minimal value of f. Define the point $y \in \mathbb{R}^n$ as

$$y[i] = -\frac{\beta}{\alpha k} \quad \text{for } 1 \leq i \leq k, \qquad y[i] = 0 \quad \text{for } k+1 \leq i \leq n.$$

Non-smooth case (proof)

- It remains to compute the minimal value of f. Define the point $y \in \mathbb{R}^n$ as

$$y[i] = -\frac{\beta}{\alpha k} \quad \text{for } 1 \leq i \leq k, \qquad y[i] = 0 \quad \text{for } k+1 \leq i \leq n.$$

- Note, that $0 \in \partial f(y)$:

$$\begin{aligned} \partial f(y) &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = \max_j y[j] \right\} \\ &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = 0 \right\} \\ 0 &\in \partial f(y). \end{aligned}$$

Non-smooth case (proof)

- It remains to compute the minimal value of f. Define the point $y \in \mathbb{R}^n$ as

$$y[i] = -\frac{\beta}{\alpha k} \quad \text{for } 1 \leq i \leq k, \qquad y[i] = 0 \quad \text{for } k+1 \leq i \leq n.$$

- Note, that $0 \in \partial f(y)$:

$$\begin{aligned} \partial f(y) &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = \max_j y[j] \right\} \\ &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = 0 \right\} \\ 0 &\in \partial f(y). \end{aligned}$$

- It follows that the minimum value of $f = f(y) = f(x^*)$ is

$$f(y) = -\frac{\beta^2}{\alpha k} + \frac{\alpha}{2} \cdot \frac{\beta^2}{\alpha^2 k} = -\frac{\beta^2}{2\alpha k}.$$

Non-smooth case (proof)

- It remains to compute the minimal value of f. Define the point $y \in \mathbb{R}^n$ as

$$y[i] = -\frac{\beta}{\alpha k} \quad \text{for } 1 \leq i \leq k, \qquad y[i] = 0 \quad \text{for } k+1 \leq i \leq n.$$

- Note, that $0 \in \partial f(y)$:

$$\begin{aligned} \partial f(y) &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = \max_j y[j] \right\} \\ &= \alpha y + \beta \text{conv} \left\{ e_i \mid i : y[i] = 0 \right\} \\ 0 &\in \partial f(y). \end{aligned}$$

- It follows that the minimum value of $f = f(y) = f(x^*)$ is

$$f(y) = -\frac{\beta^2}{\alpha k} + \frac{\alpha}{2} \cdot \frac{\beta^2}{\alpha^2 k} = -\frac{\beta^2}{2\alpha k}.$$

- Now we have:

$$f(x^i) - f(x^*) \geq 0 - \left(-\frac{\beta^2}{2\alpha k} \right) \geq \frac{\beta^2}{2\alpha k}.$$

Non-smooth case (proof)

We have: $f(x^i) - f(x^*) \geq \frac{\beta^2}{2\alpha k}$, while we need to prove that $\min\limits_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{GR}{2(1+\sqrt{k})}$.

Non-smooth case (proof)

We have: $f(x^i) - f(x^*) \geq \frac{\beta^2}{2\alpha k}$, while we need to prove that $\min\limits_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{GR}{2(1+\sqrt{k})}$.

Convex case

$$\alpha = \frac{G}{R}\frac{1}{1+\sqrt{k}} \quad \beta = \frac{\sqrt{k}}{1+\sqrt{k}}$$

$$\frac{\beta^2}{2\alpha} = \frac{GRk}{2(1+\sqrt{k})}$$

Note, in particular, that $\|y\|_2^2 = \frac{\beta^2}{\alpha^2 k} = R^2$ with these parameters

$$\min_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{\beta^2}{2\alpha k} = \frac{GR}{2(1+\sqrt{k})}$$

Non-smooth case (proof)

We have: $f(x^i) - f(x^*) \geq \frac{\beta^2}{2\alpha k}$, while we need to prove that $\min\limits_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{GR}{2(1+\sqrt{k})}$.

Convex case

$$\alpha = \frac{G}{R}\frac{1}{1+\sqrt{k}} \quad \beta = \frac{\sqrt{k}}{1+\sqrt{k}}$$

$$\frac{\beta^2}{2\alpha} = \frac{GRk}{2(1+\sqrt{k})}$$

Note, in particular, that $\|y\|_2^2 = \frac{\beta^2}{\alpha^2 k} = R^2$ with these parameters

$$\min_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{\beta^2}{2\alpha k} = \frac{GR}{2(1+\sqrt{k})}$$

Strongly convex case

$$\alpha = \frac{G}{2R} \quad \beta = \frac{G}{2}$$

Note, in particular, that $\|y\|_2^2 = \frac{\beta^2}{\alpha^2 k} = \frac{G^2}{4\alpha^2 k} = R^2$ with these parameters

$$\min_{i \in [1,k]} f(x^i) - f(x^*) \geq \frac{G^2}{8\alpha k}$$

Applications

Linear Least Squares with l_1-regularization

$$\min_{x \in \mathbb{R}^n} \frac{1}{2}\|Ax - b\|_2^2 + \lambda \|x\|_1$$

The algorithm will be written as:

$$x_{k+1} = x_k - \alpha_k \left(A^\top (Ax_k - b) + \lambda \text{sign}(x_k) \right),$$

where the signum function is taken element-wise.



Regularized logistic regression

Given $(x_i, y_i) \in \mathbb{R}^p \times \{0, 1\}$ for $i = 1, \dots, n$, the logistic regression function is defined as:

$$f(\theta) = \sum_{i=1}^{n} \left(-y_i x_i^T \theta + \log(1 + \exp(x_i^T \theta))\right)$$

This is a smooth and convex function with its gradient given by:

$$\nabla f(\theta) = \sum_{i=1}^{n} \left(y_i - s_i(\theta)\right) x_i$$

where $s_i(\theta) = \frac{\exp(x_i^T \theta)}{1 + \exp(x_i^T \theta)}$, for $i = 1, \dots, n$. Consider the regularized problem:

$$f(\theta) + \lambda r(\theta) \to \min_{\theta}$$

where $r(\theta) = \|\theta\|_2^2$ for the ridge penalty, or $r(\theta) = \|\theta\|_1$ for the lasso penalty.

Support Vector Machines

Let $D = \{(x_i, y_i) \mid x_i \in \mathbb{R}^n, y_i \in \{\pm 1\}\}$

We need to find $\theta \in \mathbb{R}^n$ and $b \in \mathbb{R}$ such that

$$\min_{\theta \in \mathbb{R}^n, b \in \mathbb{R}} \frac{1}{2}\|\theta\|_2^2 + C \sum_{i=1}^{m} \mathsf{max}[0, 1 - y_i(\theta^\top x_i + b)]$$

References

- Subgradient Methods Stephen Boyd (with help from Jaehyun Park)