

R_Worksheet_Paclibar#1

Jhon Albert Paclibar

2024-09-20

#1. Set up a vector named age, consisting of 34, 28, 22, 36, 27, 18, 52, 39, 42, 29, 35, 31, 27, 22, 37, 34, 19, 20, 57, 49, 50, 37, 46, 25, 17, 37, 42, 53, 41, 51, 35, 24, 33, 41.

```
age <- c(34, 28, 22, 36, 27, 18, 52, 39, 42, 29, 35, 31, 27, 22, 37, 34, 19, 20, 57, 49, 50, 37, 46, 25
```

a. How many data points?

```
length(age)
```

```
## [1] 34
```

2. Find the reciprocal of the values for age.

```
library(MASS)
reciprocal0fage <- 1 / age
fractions(reciprocal0fage)
```

```
## [1] 1/34 1/28 1/22 1/36 1/27 1/18 1/52 1/39 1/42 1/29 1/35 1/31 1/27 1/22 1/37
## [16] 1/34 1/19 1/20 1/57 1/49 1/50 1/37 1/46 1/25 1/17 1/37 1/42 1/53 1/41 1/51
## [31] 1/35 1/24 1/33 1/41
```

3. Assign also new_age <- c(age, 0, age).

```
new_age <- c(age, 0, age)
new_age
```

```
## [1] 34 28 22 36 27 18 52 39 42 29 35 31 27 22 37 34 19 20 57 49 50 37 46 25 17
## [26] 37 42 53 41 51 35 24 33 41 0 34 28 22 36 27 18 52 39 42 29 35 31 27 22 37
## [51] 34 19 20 57 49 50 37 46 25 17 37 42 53 41 51 35 24 33 41
```

What happen to the new_age?

-The new_age vector is a combination of the original age values, 0

4. Sort the values for age.

```
sorted0fage <- sort(age)
sorted0fage
```

```
## [1] 17 18 19 20 22 22 24 25 27 27 28 29 31 33 34 34 35 35 36 37 37 37 39 41 41
## [26] 42 42 46 49 50 51 52 53 57
```

5. Find the minimum and maximum value for age.

```
min0age <- min(age)
min0age
```

```
## [1] 17
```

```
max0fage <- max(age)
max0fage
```

```
## [1] 57
```

6. Set up a vector named data, consisting of 2.4, 2.8, 2.1, 2.5, 2.4, 2.2, 2.5, 2.3, 2.5, 2.3, 2.4, and 2.7.

```
data <- c(2.4, 2.8, 2.1, 2.5, 2.4, 2.2, 2.5, 2.3, 2.5, 2.3, 2.4, 2.7)
```

a. How many data points?

```
length(data)
```

```
## [1] 12
```

7. Generates a new vector for data where you double every value of the data. | What happen to the data?

```
doubledThedata <- data * 2
doubledThedata
```

```
## [1] 4.8 5.6 4.2 5.0 4.8 4.4 5.0 4.6 5.0 4.6 4.8 5.4
```

What happen to the data?

- Every value multiplies to two.

8. Generate a sequence for the following scenario:

8.1 Integers from 1 to 100.

```
seq1To100 <- seq(1, 100)
seq1To100
```

```
## [1] 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18
## [19] 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36
## [37] 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54
```

```
## [55] 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72
## [73] 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90
## [91] 91 92 93 94 95 96 97 98 99 100
```

8.2 Numbers from 20 to 60

```
seq20To60 <- seq(20, 60)
seq20To60
```

```
## [1] 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44
## [26] 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60
```

*8.3 Mean of numbers from 20 to 60

```
mean20To60 <- mean(seq20To60)
mean20To60
```

```
## [1] 40
```

*8.4 Sum of numbers from 51 to 91

```
sum51To91 <- sum(51:91)
sum51To91
```

```
## [1] 2911
```

*8.5 Integers from 1 to 1,000

```
seq_1_1000 <- seq(1, 1000)
```

a. How many data points from 8.1 to 8.4?

-There are 143 data points

c. For 8.5 find only maximum data points until 10.

```
maxTo10 <- max(seq_1_1000[1:10])
maxTo10
```

```
## [1] 10
```

9. *Print a vector with the integers between 1 and 100 that are not divisible by 3, 5 and 7 using filter option.

```
filteredValues <- Filter(function(i) { all(i %% c(3, 5, 7) != 0) }, seq(100))
filteredValues
```

```
## [1] 1 2 4 8 11 13 16 17 19 22 23 26 29 31 32 34 37 38 41 43 44 46 47 52 53
## [26] 58 59 61 62 64 67 68 71 73 74 76 79 82 83 86 88 89 92 94 97
```

10. Generate a sequence backwards of the integers from 1 to 100.

```
backwardSeq <- seq(100,1)
backwardSeq
```

```
## [1] 100 99 98 97 96 95 94 93 92 91 90 89 88 87 86 85 84 83
## [19] 82 81 80 79 78 77 76 75 74 73 72 71 70 69 68 67 66 65
## [37] 64 63 62 61 60 59 58 57 56 55 54 53 52 51 50 49 48 47
## [55] 46 45 44 43 42 41 40 39 38 37 36 35 34 33 32 31 30 29
## [73] 28 27 26 25 24 23 22 21 20 19 18 17 16 15 14 13 12 11
## [91] 10 9 8 7 6 5 4 3 2 1
```

11. List all the natural numbers below 25 that are multiples of 3 or 5.

```
multiplesOf3_5 <- seq(1, 24)[seq(1, 24) %% 3 == 0 | seq(1, 24) %% 5 == 0]
multiplesOf3_5
```

```
## [1] 3 5 6 9 10 12 15 18 20 21 24
```

a. How many data points from 10 to 11?

-There are 111 data points

12. Statements can be grouped together using braces ‘{’ and ‘}’. A group of statements is sometimes called a block. Single statements are evaluated when a new line is typed at the end of the syntactically complete statement. Blocks are not evaluated until a new line is entered after the closing brace.

Enter this statement:

```
x <- {0 + x + 5 + }
```

Describe the output. -It outputs error

13. *Set up a vector named score, consisting of 72, 86, 92, 63, 88, 89, 91, 92, 75, 75 and 77. To access individual elements of an atomic vector, one generally uses the x[i] construction.

Find x[2] and x[3]. Write the R code and its output.

```
score <- c(72, 86, 92, 63, 88, 89, 91, 92, 75, 75, 77)
```

```
score[2]
```

```
## [1] 86
```

```
score[3]
```

```
## [1] 92
```

14. *Create a vector a = c(1,2,NA,4,NA,6,7).

```
a <- c(1,2,NA,4,NA,6,7)
print(a,na.print="-999")
```

```
## [1] 1 2 -999 4 -999 6 7
```

b. Write the R code and its output. Describe the output.

-The NA in the vector a changed to 999.

15. A special type of function calls can appear on the left hand side of the assignment operator as in > class(x) <- “foo”.

#Follow the codes below:

```
name = readline(prompt="Input your name: ")

## Input your name:
age = readline(prompt="Input your age: ")

## Input your age:
print(paste("My name is",name, "and I am",age , "years old. "))

## [1] "My name is  and I am  years old."
print(R.version.string)

## [1] "R version 4.4.1 (2024-06-14)"
```