

Assignment-based Subjective Questions

1. From your analysis of the categorical variables from the dataset, what could you infer about their effect on the dependent variable?

Below are the points we can infer from the visualization data

- Booking seemed to be almost equal either on working day or non-working day.
- When it's not holiday, booking seems to be less in number which seems reasonable as on holidays, people may want to spend time at home and enjoy with family.
- Clear weather attracted more booking which seems obvious
- Most of the bookings has been done during the month of may, june, july, aug, sep and oct. Trend increased starting of the year till mid of the year and then it started decreasing as we approached the end of year.
- Booking count has increased drastically from 2018 to 2019.

2. Why is it important to use `drop_first=True` during dummy variable creation?

`drop_first = True` is important to use, as it helps in reducing the extra column created during dummy variable creation. Hence it reduces the correlations created among dummy variables.

3. Looking at the pair-plot among the numerical variables, which one has the highest correlation with the target variable?

The 'temp' variable has the highest correlation.

4. How did you validate the assumptions of Linear Regression after building the model on the training set?

We have validated with 3 assumptions

- i. Normality of error terms
- ii. Multicollinearity
- iii. Linear relationship validation

5. Based on the final model, which are the top 3 features contributing significantly towards explaining the demand of the shared bikes?

The top 3 features contributing towards the demand of the shared bikes are “temp”, “sep” and “winter”

General Subjective Questions

1. Explain the linear regression algorithm in detail.

Linear regression may be defined as the statistical model that analyses the linear relationship between a dependent variable with given set of independent variables. Linear relationship between variables means that when the value of one or more independent variables will change (increase or decrease), the value of dependent variable will also change accordingly (increase or decrease).

Mathematically the relationship can be represented with the help of following equation

$Y = mX + c$ Here, Y is the dependent variable we are trying to predict.

X is the independent variable we are using to make predictions.

m is the slope of the regression line which represents the effect X has on .

c is a constant, known as the Y-intercept. If $X = 0$, Y would be equal to c.

cc

2. Explain the Anscombe's quartet in detail.

Anscombe's Quartet was developed by statistician Francis Anscombe. It comprises four datasets, each containing eleven (x, y) pairs. The essential thing to note about these datasets is that they share the same descriptive statistics. But things change completely, and I must emphasize COMPLETELY, when they are graphed. Each graph tells a different story irrespective of their similar summary statistics.

3. What is Pearson's R?

Pearson's r is a numerical summary of the strength of the linear association between the variables. If the variables tend to go up and down together, the correlation coefficient will be positive. If the variables tend to go up and down in opposition with low values of one variable associated with high values of the other, the correlation coefficient will be negative.

The Pearson correlation coefficient, r , can take a range of values from +1 to -1. A value of 0 indicates that there is no association between the two variables. A value greater than 0 indicates a positive association; that is, as the value of one variable increases, so does the value of the other variable. A value less than 0 indicates a negative association; that is, as the value of one variable increases, the value of the other variable decreases.

4. What is scaling? Why is scaling performed? What is the difference between normalized scaling and standardized scaling?

Feature Scaling is a technique to standardize the independent features present in the data in a fixed range. It is performed during the data pre-processing to handle highly varying magnitudes or values or units. If feature scaling is not done, then a machine learning algorithm tends to weigh greater values, higher and consider smaller values as the lower values, regardless of the unit of the values.

| S.NO. | Normalized scaling | Standardized scaling |
|-------|--|--|
| 1. | Minimum and maximum value of features are used for scaling | Mean and standard deviation is used for scaling. |
| 2. | It is used when features are of different scales. | It is used when we want to ensure zero mean and unit standard deviation. |
| 3. | Scales values between [0, 1] or [-1, 1]. | It is not bounded to a certain range. |
| 4. | It is really affected by outliers. | It is much less affected by outliers. |
| 5. | Scikit-Learn provides a transformer called MinMaxScaler for Normalization. | Scikit-Learn provides a transformer called StandardScaler for standardization. |

5. You might have observed that sometimes the value of VIF is infinite. Why does this happen?

If there is perfect correlation, then $VIF = \text{infinity}$. A large value of VIF indicates that there is a correlation between the variables. If the VIF is 4, this means that the variance of the model coefficient is inflated by a factor of 4 due to the presence of multicollinearity.

When the value of VIF is infinite it shows a perfect correlation between two independent variables. In the case of perfect correlation, we get R-squared (R^2) = 1, which lead to $1/(1-R^2)$ infinity. To solve this we need to drop one of the variables from the dataset which is causing this perfect multicollinearity. 6.

6. What is a Q-Q plot? Explain the use and importance of a Q-Q plot in linear regression. (3 marks)

The quantile-quantile (q-q) plot is a graphical technique for determining if two data sets come from populations with a common distribution.

Use of Q-Q plot:

A q-q plot is a plot of the quantiles of the first data set against the quantiles of the second dataset. By a quantile, we mean the fraction (or percent) of points below the given value. That is, the 0.3 (or 30%) quantile is the point at which 30% percent of the data fall below and 70% fall above that value. A 45-degree reference line is also plotted. If the two sets come from a population with the same distribution, the points should fall approximately along this reference line. The greater the departure from this reference line, the greater the evidence for the conclusion that the two data sets have come from populations with different distributions. Importance of Q-Q plot: When there are two data samples, it is often desirable to know if the assumption of a common distribution is justified. If so, then location and scale estimators can pool both data sets to obtain estimates of the common location and scale. If two samples do differ, it is also useful to gain some understanding of the differences. The q-q plot can provide more insight into the nature of the difference than analytical methods such as the chi-square and Kolmogorov-Smirnov 2-sample tests.