

Internal Waves in the Ocean

Master 2 – Physique de l’Océan et du Climat

Jonathan Gula (jonathan.gula@univ-brest.fr)


Clément Vic (clement.vic@ifremer.fr)

Gaspard Geoffroy (gaspard.Geoffroy@ifremer.fr)

Outline

1. A general introduction to ocean waves
2. What are internal waves ? Why do we study internal waves ?
3. Internal waves in the two-layer shallow-water model
4. Internal waves in the continuously-stratified model
- 5. Generation of internal waves**
6. Propagation of internal waves
7. Dissipation of internal waves and impacts

5. Generation of internal waves




Three main classes of waves:

- **near-inertial waves**, mostly generated by winds
- **internal tides = baroclinic tides**, generated by the interaction of the barotropic tide with the seafloor topography
- **lee waves**, generated by low-frequency currents interacting with the seafloor topography


5. Generation of internal waves

Global energetics of internal waves:

- internal tides: 1 TW (10^{12} W)
- lee waves: 0.15-0.75 TW



Generation of internal tides




Generation of lee waves


5. Generation of internal waves

Global energetics of internal waves:

- internal tides: 1 TW (10^{12} W)
- lee waves: 0.15-0.75 TW
- near-inertial waves: 0.3-1.5 TW




5. Generation of internal waves




5. Generation of internal waves

Generation of internal tides



5. Generation of internal waves

Generation of internal tides. We first need to know the barotropic tidal currents.




amplitude of the M2 tide

5. Generation of internal waves


Generation of internal tides.

A barotropic velocity profile impinging on a sloping boundary. On the boundary, the velocity vector can only have a component tangential to it and hence over the slope, the velocity profile is no longer barotropic. Energy must go into generating internal waves, and propagate away along characteristics.




5. Generation of internal waves

Generation of internal tides.



5. Generation of internal waves

Generation of internal tides.




5. Generation of internal waves

Generation of internal tides in the deep ocean.

6 dimensional parameters :

- Coriolis frequency f
- Brunt-Vaisala frequency N
- Forcing (tidal) frequency ω
- Topographic height h_0
- Topographic wavenumber $k = 2\pi/l_0$
- Magnitude of the tidal current u_0



4 dimensionless parameters :

- ω/N
- ω/f
- ku_0/ω , ratio of the “tidal excursion” to the topographic length scale

$$\epsilon = \frac{kh_0}{\alpha} = \frac{\text{topographic slope}}{\text{beam slope}}, \text{ with } \alpha = \sqrt{\frac{\omega^2 - f^2}{N^2 - \omega^2}}$$

5. Generation of internal waves

Generation of internal tides in the deep ocean.

Assumptions to derive a linear expression for w :

$$(1) - f < \omega < N$$


5. Generation of internal waves

Generation of internal tides in the deep ocean.

Assumptions to derive a linear expression for w :

(1) - $f < \omega < N$

(2) - The topographic slopes are subcritical : $\epsilon = \frac{kh_0}{\alpha} = \frac{\text{topographic slope}}{\text{beam slope}} < 1$



Beam slope in the deep ocean :
 $N^2 = 10^{-6} - 10^{-7} (\text{rad s}^{-1})^2$
 $f = 10^{-4} \text{ rad s}^{-1}$
 $\omega = 1.4 \times 10^{-4} \text{ rad s}^{-1}$ (M2 tide)
Give
 $\alpha = 0.09 - 0.34 = 9 - 34\%$ (steep !)

Vs typical topographic slopes are ~1-2%

5. Generation of internal waves

Generation of internal tides in the deep ocean.

Assumptions to derive a linear expression for w :

(1) - $f < \omega < N$

(2) - The topographic slopes are subcritical : $\epsilon = \frac{kh_0}{\alpha} = \frac{\text{topographic slope}}{\text{beam slope}} < 1$

(3) - The tidal excursion is small : :

$$\rightarrow \text{the distance traveled by a particle in half a tidal cycle} = u_0 \times \frac{1}{2} \times \frac{2\pi}{\omega} = \frac{\pi u_0}{\omega}$$

$$\text{should be smaller than half the topographic wavelength } \frac{1}{2}l_0 = \frac{\pi}{k}$$

For a semi-diurnal tide with $\omega = 1.4 \times 10^{-4} \text{ rad s}^{-1}$ and $u_0 = 1 - 4 \text{ cm s}^{-1}$,

$$\text{The theory is valid for } l_0 > \frac{2\pi u_0}{\omega} > 400 - 1800 \text{ m,}$$

\rightarrow OK for most topographic structures over mid-ocean ridges.

5. Generation of internal waves

Generation of internal tides in the deep ocean.

For the 2D (x-z) problem with $h = h_0 \cos(kx)$ and a tidal current $u = u_0 \cos(\omega t)$, the response is:

$$u = mh_0 u_0 \cos(kx) \sin(mz + \omega t)$$
$$w = -kh_0 u_0 \sin(kx) \cos(mz + \omega t)$$

The vertical energy flux is $F = \frac{1}{4} \rho_0 \omega^{-1} [(N^2 - \omega^2)(\omega^2 - f^2)]^{1/2} k u_0^2 h_0^2$


Take-home message : the energy flux (conversion) depends on :

- The shape of the seafloor topography k, h_0^2
- The frequencies of the systems ω, f, N
- The squared amplitude of the barotropic tide current u_0^2

5. Generation of internal waves

Generation of internal tides in the deep ocean.

Generalisation to a 3-D problem with random bathymetry.




You must know :

- bathymetry properties (power density spectrum)
- tidal currents

5. Generation of internal waves

Generation of internal tides in the deep ocean.

Generalisation to a 3-D problem with random bathymetry.





The generation can be cast into vertical modes (cf section 4 of this course).

5. Generation of internal waves

Generation of internal tides in the deep ocean.


Generalisation to a 3-D problem with random bathymetry.



5. Generation of internal waves

Generation of internal tides in the deep ocean.


Global prediction of the generation into vertical modes.



5. Generation of internal waves

Generation of internal tides in the deep ocean.

Global prediction of the generation into vertical modes.



Interest of the method: link the generation to local and remote mixing. High modes are unstable and break locally vs low modes are stable and propagate long distances.

5. Generation of internal waves

Generation of internal tides ~~in the deep ocean.~~ in shallow seas.

On continental shelves / shelf breaks, the assumptions to derive a linear theory do not hold. Why ?

5. Generation of internal waves

Generation of internal tides ~~in the deep ocean.~~ in shallow seas.

On continental shelves / shelf breaks, the assumptions to derive a linear theory do not hold. Why ?

- Shallower depths : N is much larger (stratification++) \rightarrow beams are more horizontal
- Topographic slopes are steeper



$$\rightarrow \epsilon = \frac{kh_0}{\alpha} = \frac{\text{topographic slope}}{\text{beam slope}} \text{ can be no longer } < 1$$

\rightarrow critical slopes $\epsilon = 1$ and supercritical slopes $\epsilon > 1$

5. Generation of internal waves


Generation of internal tides in the deep ocean, in shallow seas.

- Internal tides are preferentially generated where the topographic slope equals the beam slope (critical slopes).
- The wave field is more organised in beams (vs individual modes in the deep ocean).




5. Generation of internal waves

Generation of internal tides (different slopes).




5. Generation of internal waves

Generation of internal tides (different slopes).




5. Generation of internal waves

Generation of lee waves



5. Generation of internal waves

Generation of lee waves




Modis, 23/11/2009 - South Atlantic – Sandwich Islands

The lower atmosphere is drier – Downwind of the islands the waves are seen when the air goes up, condensate and form clouds


5. Generation of internal waves

Generation of lee waves



5. Generation of internal waves

Generation of lee waves



5. Generation of internal waves


Generation of lee waves

- Topography $z_b = H \sin(k_x x)$
- In the frame moving at constant speed U , $z_b = H \sin(k_x(x + Ut))$

- We can identify the frequency of motion: $\omega = -k_x U$
- Recall the general equation for internal waves:

$$(w_{xx} + w_{yy} + w_{zz})_{tt} + f^2 w_{zz} + N^2(w_{xx} + w_{yy}) = 0$$


→ with a boundary condition, and some assumptions ($f = 0$, $H \ll U/N$), we can solve the equation assuming a wave form $w = w_0 \exp(i(k_x x + k_z z - \omega t))$



5. Generation of internal waves

Generation of lee waves


- if $N > U k_x = \omega$, the wave radiates



5. Generation of internal waves


Generation of lee waves

- if $N < U k_x = \omega$, the wave is trapped




5. Generation of internal waves

Generation of lee waves




5. Generation of internal waves

Generation of lee waves



5. Generation of internal waves



Generation of lee waves



5. Generation of internal waves

Generation of lee waves


Observation of lee waves by gliders in the Gulf Stream



5. Generation of internal waves

Generation of lee waves


Surface signature of lee waves



5. Generation of internal waves

Generation of lee waves


Surface signature of lee waves



5. Generation of internal waves


Generation of lee waves

Numerical model at high resolution ($\Delta x=200$ m, J. Gula)




5. Generation of internal waves

Generation of near-inertial waves by winds



5. Generation of internal waves

Generation of near-inertial waves by winds



Near-inertial oscillations of surface drifters after the passage of a storm.
The velocity field is polarised in the clockwise direction (northern hemisphere)


5. Generation of internal waves

Generation of near-inertial waves by winds

Near-inertial oscillation in the surface mixed layer (not stratified, so does not allow internal waves) create a pressure perturbation at the mixed-layer base that then propagates in the ocean interior (stratified), becoming **near-inertial waves**.


Near-inertial waves have very large horizontal wavelengths (>100 km) and short vertical wavelengths (~100 m).

They are characterised by strong vertical shear.



5. Generation of internal waves

Generation of near-inertial waves by winds



5. Generation of internal waves

References used in this document:

- MacKinnon et al., BAMS 2017, *Climate process team on internal wave-driven ocean mixing*
- Whalen et al., Nature Reviews 2020, *Internal wave-driven mixing: governing processes and consequences for climate*
- Musgrave et al. Ocean Mixing book 2022, *The lifecycle of topographically-generated internal waves*
- Vic et al., Nature Communication 2019, *Deep-ocean mixing driven by small-scale internal tides*
- Thomas & Zhai, Ocean Mixing book 2022, *The lifecycle of surface-generated near-inertial waves*
- Garrett & Kunze, ARFM 2007, *Internal tide generation in the deep ocean*
- Vic et al., JPO 2018, *The Lifecycle of Semidiurnal Internal Tides over the Northern Mid-Atlantic Ridge*
- de Lavergne et al., OM 2019, *Toward global maps of internal tide energy sinks*
- de Lavergne et al., JAMES 2020, *A Parameterization of Local and Remote Tidal Mixing*
- Rapaka et al., JFM 2013, *Tidal conversion and turbulence at a model ridge: direct and large eddy simulations*
- de Marez et al., SR 2020, *Interaction of the Gulf Stream with small scale topography: a focus on lee waves*
- Dossmann et al. JGR 2016, *Experiments with mixing in stratified flow over a topographic ridge*

5. Generation of internal waves

References used in this document:

- Todd, GRL 2017, *High-frequency internal waves and thick bottom mixed layers observed by gliders in the Gulf Stream*
-