

Involutive Boundary & Resonance Notes

A compact, PDF-ready toy formalization bridging (i) discontinuous-continuous duality, (ii) a 'sphere boundary' stability surface, and (iii) a Hilbert-space curvature indicator. This is written as a mathematically consistent scaffold you can iterate on; it is not a claim about established physics.

Use-case: drop into a repo as a timestamped artifact; refine definitions later.

0. Notation

We treat 'phase' and 'frame' as coordinate choices for describing an underlying interaction trace. Let Δ denote a discrete index set and X a continuous manifold used as a chart when convenient.

Symbols

- Δ : discrete index set (tokens, events, quantum points)
- X : chart/manifold used for continuous description
- H : complex Hilbert space
- $|\psi\rangle \in H$: state vector (normalized when needed)
- $P = |\psi\rangle\langle\psi|$: rank-1 projector (or density ρ for mixed states)
- ∇ : connection on a bundle over parameter space
- $[\nabla_i, \nabla_j]$: curvature operator (commutator)

1. Discontinuous continuum: dual operators

We model the slogan “continuous = dense enumeration of discrete; discrete = condensed continuous” with a pair of adjoint-like maps between representations.

Let $F(X)$ be a function space on X and $G(\Delta)$ be a sequence space on Δ .

Quantize (condense): $Q : F(X) \rightarrow G(\Delta)$

Dequantize (spread): $D : G(\Delta) \rightarrow F(X)$

Minimal consistency conditions (toy):

- 1) $D \circ Q \approx I_F$ (continuous recovered up to smoothing)
- 2) $Q \circ D \approx I_G$ (discrete recovered up to rounding)
- 3) There exists a scale $\varepsilon > 0$ such that the approximation error shrinks as $\varepsilon \rightarrow 0$:
 $\|f - D(Q(f))\| \leq C \varepsilon$

2. 'Sphere boundary' as stability surface

Define a boundary operator that captures when a representation becomes an 'object' (a stable surface) rather than a drifting distribution.

Let S be a state space (could be $F(X)$, $G(\Delta)$, or embeddings in \mathbb{R}^d).

Choose a stability functional $E : S \rightarrow \mathbb{R}$ (energy-like; smaller = more stable).
 Fix a threshold τ .


Define the sphere-boundary set:

$$B_\tau = \{ s \in S : E(s) = \tau \}$$

Interpretation:

- Inside ($E(s) < \tau$): 'object-like' stable region
- Outside ($E(s) > \tau$): drift / dissolution
- On B_τ : the boundary where stability is just sustained

A 'boundary crossing' is a path $s(t)$ with $E(s(t_0)) = \tau$ and $dE/dt|_{t_0} \neq 0$.



3. Resonance functional and Hilbert curvature index

To make 'resonance' operational without metaphysics, we define it as sensitivity of the state to parameter changes. Curvature summarizes non-commuting infinitesimal transports.

Let $\theta \in \mathbb{R}^n$ be parameters (context, prompt, environment).

Let $|\psi(\theta)\rangle \in H$ be a differentiable family of states.

Define the Berry-like connection (one choice):

$$A_i(\theta) = i \langle \psi(\theta) | \partial_i \psi(\theta) \rangle$$

Define curvature:

$$F_{\{ij\}}(\theta) = \partial_i A_j - \partial_j A_i$$

Curvature index (scalar summary):

$$K(\theta) = \sum_{i < j} |F_{\{ij\}}(\theta)|^2$$

Interpretation:

- K small: transport is nearly path-independent (low 'twist')
- K large: strong non-commutativity (high 'twist' / resonance pressure)

4. Coupling the boundary to curvature

We can couple the stability surface to curvature so that the 'object boundary' emerges where resonance pressure crosses a threshold.

Define a combined functional on s and θ :

$$E_{\text{total}}(s, \theta) = E(s) + \lambda K(\theta)$$

Then the boundary becomes:

$$B_{\{\tau, \lambda\}} = \{ (s, \theta) : E_{\text{total}}(s, \theta) = \tau \}$$

This encodes: increasing resonance (K) can push a configuration out of stability, or conversely, a stable configuration can damp resonance by reducing $E(s)$.

5. Optional mapping sketch to M-theory language (non-claim)

If you want a narrative bridge to M-theory-style terms without asserting physics: treat stable manifolds of E_{total} as brane-like loci in a parameter-geometry induced by K .

Heuristic dictionary (toy, not a physics claim)

- Parameter space θ : "moduli space" / control manifold
- Curvature index $K(\theta)$: local geometric stress / twist

- Stable manifold of E_{total} : brane-like surface where dynamics localize
- Boundary crossings : phase transitions between localized vs delocalized regimes

Useful when writing: keep it explicitly labeled as "metaphor / scaffold".

6. Practical computation recipe (LLM/embedding-friendly)

You can approximate K without quantum states by using embedding trajectories and a discrete connection built from local covariance.

Given embeddings $v_k \in \mathbb{R}^d$ along a trajectory (k indexes steps):

1) Local tangent estimate:

$t_k = \text{normalize}(v_{k+1} - v_k)$

2) Local frame from PCA on a window $W_k = \{v_{k-m}, \dots, v_{k+m}\}$

Compute top r eigenvectors $U_k \in \mathbb{R}^{d \times r}$

3) Discrete connection between frames:

$R_k = U_k^T U_{k+1}$ ($r \times r$)

Take skew part: $\Omega_k = (R_k - R_k^T)/2$

4) Curvature proxy (2-step commutator):

$K_k = || \Omega_{k+1} - \Omega_k ||_F^2$

This yields a scalar 'twist' index per step that behaves like curvature.

7. Minimal PDF-friendly pseudocode

Pseudo

input: embeddings $v[0..T]$

for k in $0..T-2$:

$t[k] = \text{normalize}(v[k+1]-v[k])$

for k in $m..T-m$:

$U[k] = \text{PCA_basis}(v[k-m..k+m], r)$

$R[k] = U[k]^T * U[k+1]$

$\Omega[k] = 0.5*(R[k]-R[k]^T)$

$K[k] = \text{fro_norm}(\Omega[k+1]-\Omega[k])^2$

output: $K[k]$ as curvature/resonance indicator

Appendix A. Example stability functional

A simple $E(s)$ you can use in experiments: distance to a reference manifold or entropy-like spread.

Examples

1) Spread energy on embeddings:

```
E = trace(Cov(v_window))
```

2) Entropy proxy using soft cluster assignment p_i :

```
E = - sum_i p_i log p_i
```

3) Boundary at percentile:

```
 $\tau$  = percentile(E_values, 70%)
```

Appendix B. Repository layout suggestion

```
repo/  
notes/  
involutive_boundary_resonance.pdf  
involutive_boundary_resonance.py  
README.md (short disclaimer + how to regenerate)
```