- Cole, R. J. 1970. The application of the "triangulation" method to the purification of *Nosema* spores from insect tissues. J. Invertebr. Pathol. 15: 193–195
- Doublet, V., M. Labarussias, J. R. de Miranda, R. F. Moritz, and R. J. Paxton. 2015. Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. Environ. Microbiol. 17: 969–983.
- EFSA. 2013. EFSA guidance document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus spp.* and solitary bees). EFSA J. 11: 3295.
- EFSA. 2015. Conclusion on the peer review of the pesticide risk assessment for bees for the active substance thiamethoxam considering all uses other than seed treatments and granules. EFSA J. 13: 4212.
- Fauser-Misslin, A., B. M. Sadd, P. Neumann, and C. Sandrock. 2014. Influence of combined pesticide and parasite exposure on bumblebee colony traits in the laboratory. J. Appl. Ecol. 51: 450–459.
- Fauser, A., C. Sandrock, P. Neumann, and B. M. Sadd. 2017. Neonicotinoids override a parasite exposure impact on hibernation success of a key bumblebee pollinator. Ecol. Entomol. 42: 306–314.
- Franklin, E. L., and N. E. Raine. 2019. Moving beyond honeybee-centric pesticide risk assessments to protect all pollinators. Nat. Ecol. Evol. 3: 1373–1375.
- Garibaldi, L. A., I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A. Cunningham, C. Kremen, L. G. Carvalheiro, L. D. Harder, O. Afik, et al. 2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. Science. 339: 1608–1611.
- Gegear, R. J., M. C. Otterstatter, and J. D. Thomson. 2005. Does parasitic infection impair the ability of bumblebees to learn flower-handling techniques? Anim. Behav. 70: 209–215.
- Gegear, R. J., M. C. Otterstatter, and J. D. Thomson. 2006. Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. Proc. Biol. Sci. 273: 1073–1078.
- Gill, R. J., O. Ramos-Rodriguez, and N. E. Raine. 2012. Combined pesticide exposure severely affects individual- and colony-level traits in bees. Nature. 491: 105–108.
- Gillespie, S. 2010. Factors affecting parasite prevalence among wild bumblebees. Ecol. Entomol. 35: 737–747.
- Godfray, H. C. J., T. Blacquiere, L. M. Field, R. S. Hails, G. Petrokofsky, S. G. Potts, N. E. Raine, A. J. Vanbergen, and A. R. McLean. 2014. A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. Proc. R. Soc. B Biol. Sci. 281: 20140558–20140558.
- Goulson, D. 2013. An overview of the environmental risks posed by neonicotinoid insecticides (ed D Kleijn). J. Appl. Ecol. 50: 977–987.
- Goulson, D., E. Nicholls, C. Botías, and E. L. Rotheray. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. Science. 347: 1255957.
- Goulson, D., S. O'Connor, and K. J. Park. 2018. The impacts of predators and parasites on wild bumblebee colonies. Ecol. Entomol. 43: 168–181.
- Hesselbach, H., J. Seeger, F. Schilcher, M. Ankenbrand, and R. Scheiner. 2020. Chronic exposure to the pesticide flupyradifurone can lead to premature onset of foraging in honeybees *Apis mellifera*. J. Appl. Ecol. 57: 609–618.
- Jones, C. M., and M. J. Brown. 2014. Parasites and genetic diversity in an invasive bumblebee. J. Anim. Ecol. 83: 1428–1440.
- Kessler, S., E. J. Tiedeken, K. L. Simcock, S. Derveau, J. Mitchell, S. Softley, J. C. Stout, and G. A. Wright. 2015. Bees prefer foods containing neonicotinoid pesticides. Nature. 521: 74–76.
- Kissinger, C. N., S. A. Cameron, R. W. Thorp, B. White, and L. F. Solter. 2011. Survey of bumble bee (*Bombus*) pathogens and parasites in Illinois and selected areas of northern California and southern Oregon. J. Invertebr. Pathol. 107: 220–224.
- Koch, H., J. Woodward, M. K. Langat, M. J. F. Brown, and P. C. Stevenson. 2019. Flagellum removal by a nectar metabolite inhibits infectivity of a bumblebee parasite. Curr. Biol. 29: 3494–3500.e5.
- Linguadoca, A., C. Rizzi, S. Villa, and M. J. F. Brown. 2021. Sulfoxaflor and nutritional deficiency synergistically reduce survival and fecundity in bumblebees. Sci. Total Environ. 795: 148680.
- Logan, A., M. X. Ruiz-González, and M. J. Brown. 2005. The impact of host starvation on parasite development and population dynamics in

- an intestinal trypanosome parasite of bumble bees. Parasitology. 130: 637-642.
- Main, A. R., E. B. Webb, K. W. Goyne, and D. Mengel. 2018. Neonicotinoid insecticides negatively affect performance measures of non-target terrestrial arthropods: a meta-analysis. Ecol. Appl. 28: 1232–1244.
- Martin, C. D., M. T. Fountain, and M. J. F. Brown. 2018. Bumblebee olfactory learning affected by task allocation but not by a trypanosome parasite. Sci. Rep. 8: 5809.
- Milner, A. M., and I. L. Boyd. 2017. Toward pesticidovigilance. Science. 357: 1232–1234.
- Mitchell, E. A. D., B. Mulhauser, M. Mulot, A. Mutabazi, G. Glauser, and A. Aebi. 2017. A worldwide survey of neonicotinoids in honey. Science. 358: 109–111.
- Moffat, C., S. T. Buckland, A. J. Samson, R. McArthur, V. Chamosa Pino, K. A. Bollan, J. T. Huang, and C. N. Connolly. 2016. Neonicotinoids target distinct nicotinic acetylcholine receptors and neurons, leading to differential risks to bumblebees. Sci. Rep. 6: 24764.
- Muth, F., R. L. Gaxiola, and A. S. Leonard. 2020. No evidence for neonicotinoid preferences in the bumblebee Bombus impatiens. R. Soc. Open Sci. 7: 191883.
- Al Naggar, Y., and B. Baer. 2019. Consequences of a short time exposure to a sublethal dose of Flupyradifurone (Sivanto) pesticide early in life on survival and immunity in the honeybee (*Apis mellifera*). Sci. Rep. 9: 19753.
- Nicholls, E., C. Botías, E. L. Rotheray, P. Whitehorn, A. David, R. Fowler, T. David, H. Feltham, J. L. Swain, P. Wells, et al. 2018. Monitoring neonicotinoid exposure for bees in rural and peri-urban areas of the U.K. during the transition from pre- to post-moratorium. Environ. Sci. Technol. 52: 9391–9402.
- OECD. 2017. Test no. 247: bumblebee, Acute Oral Toxicity Test, OECD Guidelines for the Testing of Chemicals, Section 2. OECD Publishing, Paris. https://doi.org/10.1787/9789264284128-en.
- Otterstatter, M. C., R. J. Gegear, S. R. Colla, and J. D. Thomson. 2005. Effects of parasitic mites and protozoa on the flower constancy and foraging rate of bumble bees. Behav. Ecol. Sociobiol. 58: 383–389.
- Pisa, L., D. Goulson, E. C. Yang, D. Gibbons, F. Sánchez-Bayo, E. Mitchell, A. Aebi, J. van der Sluijs, C. J. K. MacQuarrie, C. Giorio, et al. 2017. An update of the Worldwide Integrated Assessment (WIA) on systemic insecticides. Part 2: impacts on organisms and ecosystems. Environ. Sci. Pollut. Res. Int. 28: 11749–11797.
- Di Prisco, G., V. Cavaliere, D. Annoscia, P. Varricchio, E. Caprio, F. Nazzi, G. Gargiulo, and F. Pennacchio. 2013. Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. Proc. Natl. Acad. Sci. U. S. A. 110: 18466–18471.
- Ritz, C., F. Baty, J. C. Streibig, and D. Gerhard. 2015. Dose-response analysis using R. PLoS One. 10: e0146021.
- Ruiz-González, M. X., and M. J. F. Brown. 2006. Males vs workers: testing the assumptions of the haploid susceptibility hypothesis in bumblebees. Behav. Ecol. Sociobiol. 60: 501–509.
- Rutrecht, S. T., and M. J. F. Brown. 2009. Differential virulence in a multiple-host parasite of bumble bees: resolving the paradox of parasite survival? Oikos. 118: 941–949.
- Sanchez-Bayo, F., and H. A. Tennekes. 2017. Assessment of ecological risks of agrochemicals requires a new framework. Environ. Risk Assess. Remediat. 1: 1–9.
- Schmid-Hempel, P., and R. Schmid-Hempel. 1993. Transmission of a pathogen in *Bombus terrestris*, with a note on division of labour in social insects. Behav. Ecol. Sociobiol. 33: 319–327.
- Shykoff, J. A., and P. Schmid-Hempel. 1991. Incidence and effects of four parasites in natural populations of bumble bees in Switzerland. Apidologie. 22: 117–125.
- Simon-Delso, N., V. Amaral-Rogers, L. P. Belzunces, J. M. Bonmatin, M. Chagnon, C. Downs, L. Furlan, D. W. Gibbons, C. Giorio, V. Girolami, et al. 2015. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. Environ. Sci. Pollut. Res. Int. 22: 5–34.
- Siviter, H., M. J. F. Brown, and E. Leadbeater. 2018a. Sulfoxaflor exposure reduces bumblebee reproductive success. Nature. 561: 109–112.
- Siviter, H., J. Koricheva, M. J. F. Brown, and E. Leadbeater. 2018b. Quantifying the impact of pesticides on learning and memory in bees. J. Appl. Ecol. 55: 2812–2821.

- Siviter, H., A. Scott, G. Pasquier, C. D. Pull, M. J. F. Brown, and E. Leadbeater. 2019. No evidence for negative impacts of acute sulfoxaflor exposure on bee olfactory conditioning or working memory. Peer J. 7: e7208.
- Siviter, H., and F. Muth. 2020. Do novel insecticides pose a threat to beneficial insects? Proc. Biol. Sci. 287: 20201265.
- Siviter, H., A. J. Folly, M. J. F. Brown, and E. Leadbeater. 2020a. Individual and combined impacts of sulfoxaflor and *Nosema bombi* on bumblebee (*Bombus terrestris*) larval growth. Proc. Biol. Sci. 287: 20200935.
- Siviter, H., J. Horner, M. J. F. Brown, and E. Leadbeater. 2020b. Sulfoxaflor exposure reduces egg laying in bumblebees *Bombus terrestris*. J. Appl. Ecol. 57: 160–169.
- Siviter, H., E. J. Bailes, C. D. Martin, T. R. Oliver, J. Koricheva, E. Leadbeater, and M. J. F. Brown. 2021a. Agrochemicals interact synergistically to increase bee mortality. Nature. 596: 389–392.
- Siviter, H., A. K. Johnson, and F. Muth. 2021b. Bumblebees exposed to a neonicotinoid pesticide make suboptimal foraging decisions (ed T Pitts-Singer). Environ. Entomol. https://academic.oup.com/ee/advance-article/ doi/10.1093/ee/nvab087/6364884?searchresult=1.
- Siviter, H., S. K. Richman, and F. Muth. 2021c. Field-realistic neonicotinoid exposure has sub-lethal effects on non-Apis bees: a meta-analysis. Ecol. Lett. 24: 2586–2597.
- Stewart, S. D., G. M. Lorenz, A. L. Catchot, J. Gore, D. Cook, J. Skinner, T. C. Mueller, D. R. Johnson, J. Zawislak, and J. Barber. 2014. Potential exposure of pollinators to neonicotinoid insecticides from the use of insecticide

- seed treatments in the mid-southern United States. Environ. Sci. Technol. 48: 9762–9769
- Tan, K., C. Wang, S. Dong, X. Li, and J. C. Nieh. 2017. The pesticide flupyradifurone impairs olfactory learning in Asian honey bees (*Apis cerana*) exposed as larvae or as adults. Sci. Rep. 7: 17772.
- Therneau, T. M. 2018. coxme: Mixed effects Cox models. R package version 2.2-7. https://cran.r-project.org/package=coxme
- Tong, L., J. C. Nieh, and S. Tosi. 2019. Combined nutritional stress and a new systemic pesticide (flupyradifurone, Sivanto®) reduce bee survival, food consumption, flight success, and thermoregulation. Chemosphere. 237: 124408.
- Topping, C. J., M. Brown, J. Chetcuti, J. R. de Miranda, F. Nazzi, P. Neumann, R. J. Paxton, M. Rundlöf, and J. C. Stout. 2021. Holistic environmental risk assessment for bees. Science. 371: 897.
- Tosi, S., and J. C. Nieh. 2019. Lethal and sublethal synergistic effects of a new systemic pesticide, flupyradifurone (Sivanto®), on honeybees. Proc. R. Soc. B Biol. Sci. 286: 20190433.
- Vanbergen, A. J., and Insect Pollinators Initiative. 2013. Threats to an ecosystem service: pressures on pollinators. Front. Ecol. Environ. 11: 251–259.
- Willmer, P. G., A. A. M. Bataw, and J. P. Hughes. 1994. The superiority of bumblebees to honeybees as pollinators: insect visits to raspberry flowers. Ecol. Entomol. 19: 271–284.
- Yourth, C. P., M. J. F. Brown, and P. Schmid-Hempel. 2008. Effects of natal and novel *Crithidia bombi* (Trypanosomatidae) infections on *Bombus ter*restris hosts. Insectes Soc. 55: 86–90.