

interactions between agrochemicals and parasites (Fauser-Misslin et al. 2014, Siviter et al. 2021a). While our methodology could be used within the regulatory process, future research should be focused on developing methodologies that assess the potential sub-lethal interactions between agrochemicals and parasites on bees.

We found that *C. bombi* intensity was significantly higher in bumblebees that had been fed high dosages of thiamethoxam and that had subsequently died. *C. bombi* intensity typically increases for up to 7 d after inoculation and plateaus between 7- and 10-days post inoculation (Logan et al. 2005). We exposed bumblebees to thiamethoxam 7 d post inoculation and found that bees that died (on day 7) had a higher intensity of *C. bombi* than bees that survived (Fig. 4). One explanation for this is that acute exposure to thiamethoxam exerts long-term inhibition on the growth of *C. bombi*, and that this could only occur in bees that survived exposure. Alternatively, higher *C. bombi* counts in bumblebees exposed to lethal acute doses could be due to rapidly enhanced production or release of the parasite from the gut lining (Koch et al. 2019). Future experiments are needed to determine the mechanism behind this interaction. However, as we found no effect of sub-lethal thiamethoxam dosages on *C. bombi* intensity, this suggests that at field-realistic levels, thiamethoxam is unlikely to impact *C. bombi* intensity.

Neonicotinoids are the most commonly used insecticides in the world and understanding the interaction between them and bumblebee pathogens is therefore vitally important. However, as the number of insect pests that are resistant to neonicotinoids increase, and bans/restrictions on their use increase globally, novel insecticides such as sulfoxaflor or flupyradifurone could replace them over large geographical areas (Brown et al. 2016, Siviter and Muth 2020). Sulfoxaflor exposure can have significant sub-lethal impacts on bumblebee (*B. terrestris*) reproduction (Siviter et al. 2018a, Siviter et al. 2020 a,b; Linguadoca et al. 2021) (but see [Siviter et al. 2019]) and flupyradifurone exposure can impair honeybee larval development (Tan et al. 2017, Al Naggar and Baer 2019), and adult behaviour (Tosi and Nieh 2019, Tong et al. 2019, Hesselbach et al. 2020) (recently reviewed in [Siviter and Muth 2020]). Novel insecticides could also interact with bee pathogens, for example, bumblebee larvae fed sulfoxaflor in isolation showed no evidence of an increase in larval mortality, but when coexposed to sulfoxaflor, and the common bumblebee parasite *Nosema bombi*, there was a significant increase in larval mortality (Siviter et al. 2020a). Similarly, honeybees (*A. mellifera*) fed flupyradifurone and inoculated with *N. ceranae* had lower survival than unexposed bees, and those exposed to each stressor in isolation (Al Naggar and Baer 2019). While we found no interaction between *C. bombi* and the neonicotinoid thiamethoxam on bee mortality, future research should focus on understanding how novel insecticides, such as sulfoxaflor and flupyradifurone, interact with common bee parasites (Siviter and Muth 2020).

Global bee declines are thought to be driven by multiple anthropogenic stressors, including agrochemicals and parasites (Vanbergen & Insect Pollinators Initiative 2013, Goulson et al. 2015, Siviter et al. 2021a) which suggests that the agrochemical regulatory process should consider how insecticides interact with commonly occurring bee parasites (Siviter and Muth 2020). Here we show how toxicity tests, such as LD₅₀ experiments, can be modified to consider the interactions between agrochemicals and parasites. This methodology could easily be modified to test other parasites depending on the life history of the parasites. However, the sheer number of bee parasites (known and unknown), and the range of different agrochemicals used in intensive agriculture means that testing every potential interaction between parasites and agrochemicals is impractical, and in some cases, when we do not have an understanding of the parasite life-history, impossible. In these cases, post-authorisation monitoring observations, which are currently nonexistent (Milner and Boyd 2017), should be

carried out that monitor interactions between pesticides and pathogens. More broadly a move towards a more holistic approach to environmental risk assessment, that considers the interactions between multiple stressors, and models their impact on wild bees, is required to better safe-guard bees, and other pollinators, from the potential harm of agrochemicals (Siviter and Muth 2020, Topping et al. 2021).

Supplementary Data

Supplementary data are available at *Environmental Entomology* online.

Acknowledgments

We thank Alberto Linguadoca for suggestions made for data presentation and statistical analysis, Callum Martin, and Christopher Pull for assisting with the statistical analysis, Arran Folly for assisting with the parasite work, Tom Oliver for helpful general discussion, and two anonymous reviewers for helpful comments. H.S. was supported by a Royal Holloway University of London Reid PhD Scholarship, by contributions from High Wycombe Beekeeper's Association, and by a Stengl-Wyer Scholarship. This project has received funding from the European Horizon 2020 research and innovation programme under grant agreement no.773921 and from the Biotechnology and Biological Sciences Research Council, Grant/Award Number:BB/N000668/1.

Data Availability

Raw data available here <https://osf.io/vaute/>.

References Cited

- Arce, A. N., A. Ramos Rodrigues, J. Yu, T. J. Colgan, Y. Wurm, and R. J. Gill. 2018. Foraging bumblebees acquire a preference for neonicotinoid-treated food with prolonged exposure. *Proc. R. Soc B Biol. Sci.* 285: 20180655.
- Baron, G. L., V. A. A. Jansen, M. J. F. Brown, and N. E. Raine. 2017. Pesticide reduces bumblebee colony initiation and increases probability of population extinction. *Nat. Ecol. Evol.* 1: 1308–1316.
- Barton, K. 2016. MuMIn: multi-model inference. R package version 1.15.6. <https://cran.r-project.org/package=MUMIN>
- Bass, C., I. Denholm, M. S. Williamson, and R. Nauen. 2015. The global status of insect resistance to neonicotinoid insecticides. *Pestic. Biochem. Physiol.* 121: 78–87.
- Bates, D., M. Mächler, B. Bolker, and S. Walker. 2015. Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* 67: 51.
- Bonmatin, J. M., C. Giorio, V. Girolami, D. Goulson, D. P. Kreutzweiser, C. Krupke, M. Liess, E. Long, M. Marzaro, E. A. Mitchell, et al. 2015. Environmental fate and exposure; neonicotinoids and fipronil. *Environ. Sci. Pollut. Res. Int.* 22: 35–67.
- Botías, C., A. David, E. M. Hill, and D. Goulson. 2016. Contamination of wild plants near neonicotinoid seed-treated crops, and implications for non-target insects. *Sci. Total Environ.* 566–567: 269–278.
- Botías, C., A. David, E. M. Hill, and D. Goulson. 2017. Quantifying exposure of wild bumblebees to mixtures of agrochemicals in agricultural and urban landscapes. *Environ. Pollut.* 222: 73–82.
- Brown, M. J. F., R. Schmid-Hempel, and P. Schmid-Hempel. 2003. Strong context-dependent virulence in a host-parasite system: reconciling genetic evidence with theory. *J. Anim. Ecol.* 72: 994–1002.
- Brown, M. J. F., R. Loosli, and P. Schmid-Hempel. 2000. Condition-dependent expression of virulence in a trypanosome infecting bumblebees. *Oikos* 91: 421–427.
- Brown, M. J., L. V. Dicks, R. J. Paxton, K. C. Baldock, A. B. Barron, M. P. Chauzat, B. M. Freitas, D. Goulson, S. Jepsen, C. Kremen, et al. 2016. A horizon scan of future threats and opportunities for pollinators and pollination. *PeerJ*. 4: e2249.
- Burnham, K. P., and D. R. Anderson. 2002. Model selection and multimodel inference: a practical information-theoretic approach, 2nd ed. New York: Springer-verlag.

- Cole, R. J. 1970. The application of the “triangulation” method to the purification of *Nosema* spores from insect tissues. *J. Invertebr. Pathol.* 15: 193–195.
- Doublet, V., M. Labarussias, J. R. de Miranda, R. F. Moritz, and R. J. Paxton. 2015. Bees under stress: sublethal doses of a neonicotinoid pesticide and pathogens interact to elevate honey bee mortality across the life cycle. *Environ. Microbiol.* 17: 969–983.
- EFSA. 2013. EFSA guidance document on the risk assessment of plant protection products on bees (*Apis mellifera*, *Bombus spp.* and solitary bees). EFSA J. 11: 3295.
- EFSA. 2015. Conclusion on the peer review of the pesticide risk assessment for bees for the active substance thiamethoxam considering all uses other than seed treatments and granules. EFSA J. 13: 4212.
- Fausser-Misslin, A., B. M. Sadd, P. Neumann, and C. Sandrock. 2014. Influence of combined pesticide and parasite exposure on bumblebee colony traits in the laboratory. *J. Appl. Ecol.* 51: 450–459.
- Fausser, A., C. Sandrock, P. Neumann, and B. M. Sadd. 2017. Neonicotinoids override a parasite exposure impact on hibernation success of a key bumblebee pollinator. *Ecol. Entomol.* 42: 306–314.
- Franklin, E. L., and N. E. Raine. 2019. Moving beyond honeybee-centric pesticide risk assessments to protect all pollinators. *Nat. Ecol. Evol.* 3: 1373–1375.
- Garibaldi, L. A., I. Steffan-Dewenter, R. Winfree, M. A. Aizen, R. Bommarco, S. A. Cunningham, C. Kremen, L. G. Carvalheiro, L. D. Harder, O. Afik, et al. 2013. Wild pollinators enhance fruit set of crops regardless of honey bee abundance. *Science*. 339: 1608–1611.
- Gegeer, R. J., M. C. Otterstatter, and J. D. Thomson. 2005. Does parasitic infection impair the ability of bumblebees to learn flower-handling techniques? *Anim. Behav.* 70: 209–215.
- Gegeer, R. J., M. C. Otterstatter, and J. D. Thomson. 2006. Bumble-bee foragers infected by a gut parasite have an impaired ability to utilize floral information. *Proc. Biol. Sci.* 273: 1073–1078.
- Gill, R. J., O. Ramos-Rodriguez, and N. E. Raine. 2012. Combined pesticide exposure severely affects individual- and colony-level traits in bees. *Nature*. 491: 105–108.
- Gillespie, S. 2010. Factors affecting parasite prevalence among wild bumblebees. *Ecol. Entomol.* 35: 737–747.
- Godfray, H. C. J., T. Blacquiere, L. M. Field, R. S. Hails, G. Petrokofsky, S. G. Potts, N. E. Raine, A. J. Vanbergen, and A. R. McLean. 2014. A restatement of the natural science evidence base concerning neonicotinoid insecticides and insect pollinators. *Proc. R. Soc. B Biol. Sci.* 281: 20140558–20140558.
- Goulson, D. 2013. An overview of the environmental risks posed by neonicotinoid insecticides (ed D Kleijn). *J. Appl. Ecol.* 50: 977–987.
- Goulson, D., E. Nicholls, C. Botías, and E. L. Rotheray. 2015. Bee declines driven by combined stress from parasites, pesticides, and lack of flowers. *Science*. 347: 1255957.
- Goulson, D., S. O'Connor, and K. J. Park. 2018. The impacts of predators and parasites on wild bumblebee colonies. *Ecol. Entomol.* 43: 168–181.
- Hesselbach, H., J. Seeger, F. Schilcher, M. Ankenbrand, and R. Scheiner. 2020. Chronic exposure to the pesticide flupyradifurone can lead to premature onset of foraging in honeybees *Apis mellifera*. *J. Appl. Ecol.* 57: 609–618.
- Jones, C. M., and M. J. Brown. 2014. Parasites and genetic diversity in an invasive bumblebee. *J. Anim. Ecol.* 83: 1428–1440.
- Kessler, S., E. J. Tiedeken, K. L. Simcock, S. Derveau, J. Mitchell, S. Softley, J. C. Stout, and G. A. Wright. 2015. Bees prefer foods containing neonicotinoid pesticides. *Nature*. 521: 74–76.
- Kissinger, C. N., S. A. Cameron, R. W. Thorp, B. White, and L. F. Solter. 2011. Survey of bumble bee (*Bombus*) pathogens and parasites in Illinois and selected areas of northern California and southern Oregon. *J. Invertebr. Pathol.* 107: 220–224.
- Koch, H., J. Woodward, M. K. Langat, M. J. F. Brown, and P. C. Stevenson. 2019. Flagellum removal by a nectar metabolite inhibits infectivity of a bumblebee parasite. *Curr. Biol.* 29: 3494–3500.e5.
- Linguadoca, A., C. Rizzi, S. Villa, and M. J. F. Brown. 2021. Sulfoxaflor and nutritional deficiency synergistically reduce survival and fecundity in bumblebees. *Sci. Total Environ.* 795: 148680.
- Logan, A., M. X. Ruiz-González, and M. J. Brown. 2005. The impact of host starvation on parasite development and population dynamics in an intestinal trypanosome parasite of bumble bees. *Parasitology*. 130: 637–642.
- Main, A. R., E. B. Webb, K. W. Goyne, and D. Mengel. 2018. Neonicotinoid insecticides negatively affect performance measures of non-target terrestrial arthropods: a meta-analysis. *Ecol. Appl.* 28: 1232–1244.
- Martin, C. D., M. T. Fountain, and M. J. F. Brown. 2018. Bumblebee olfactory learning affected by task allocation but not by a trypanosome parasite. *Sci. Rep.* 8: 5809.
- Milner, A. M., and I. L. Boyd. 2017. Toward pesticide vigilance. *Science*. 357: 1232–1234.
- Mitchell, E. A. D., B. Mulhauser, M. Mulot, A. Mutabazi, G. Glauser, and A. Aebi. 2017. A worldwide survey of neonicotinoids in honey. *Science*. 358: 109–111.
- Moffat, C., S. T. Buckland, A. J. Samson, R. McArthur, V. Chamosa Pino, K. A. Bolland, J. T. Huang, and C. N. Connolly. 2016. Neonicotinoids target distinct nicotinic acetylcholine receptors and neurons, leading to differential risks to bumblebees. *Sci. Rep.* 6: 24764.
- Muth, F., R. L. Gaxiola, and A. S. Leonard. 2020. No evidence for neonicotinoid preferences in the bumblebee *Bombus impatiens*. *R. Soc. Open Sci.* 7: 191883.
- Al Naggari, Y., and B. Baer. 2019. Consequences of a short time exposure to a sublethal dose of Flupyradifurone (Sivanto) pesticide early in life on survival and immunity in the honeybee (*Apis mellifera*). *Sci. Rep.* 9: 19753.
- Nicholls, E., C. Botías, E. L. Rotheray, P. Whitehorn, A. David, R. Fowler, T. David, H. Feltham, J. L. Swain, P. Wells, et al. 2018. Monitoring neonicotinoid exposure for bees in rural and peri-urban areas of the U.K. during the transition from pre- to post-moratorium. *Environ. Sci. Technol.* 52: 9391–9402.
- OECD. 2017. Test no. 247: bumblebee, Acute Oral Toxicity Test, OECD Guidelines for the Testing of Chemicals, Section 2. OECD Publishing, Paris. <https://doi.org/10.1787/9789264284128-en>.
- Otterstatter, M. C., R. J. Gegeer, S. R. Colla, and J. D. Thomson. 2005. Effects of parasitic mites and protozoa on the flower constancy and foraging rate of bumble bees. *Behav. Ecol. Sociobiol.* 58: 383–389.
- Pisa, L., D. Goulson, E. C. Yang, D. Gibbons, F. Sánchez-Bayo, E. Mitchell, A. Aebi, J. van der Sluijs, C. J. K. MacQuarrie, C. Giorio, et al. 2017. An update of the Worldwide Integrated Assessment (WIA) on systemic insecticides. Part 2: impacts on organisms and ecosystems. *Environ. Sci. Pollut. Res. Int.* 28: 11749–11797.
- Di Prisco, G., V. Cavaliere, D. Annoscia, P. Varricchio, E. Caprio, F. Nazzi, G. Gargiulo, and F. Pennacchio. 2013. Neonicotinoid clothianidin adversely affects insect immunity and promotes replication of a viral pathogen in honey bees. *Proc. Natl. Acad. Sci. U. S. A.* 110: 18466–18471.
- Ritz, C., F. Baty, J. C. Streibig, and D. Gerhard. 2015. Dose-response analysis using R. *PLoS One*. 10: e0146021.
- Ruiz-González, M. X., and M. J. F. Brown. 2006. Males vs workers: testing the assumptions of the haploid susceptibility hypothesis in bumblebees. *Behav. Ecol. Sociobiol.* 60: 501–509.
- Rutrecht, S. T., and M. J. F. Brown. 2009. Differential virulence in a multiple-host parasite of bumble bees: resolving the paradox of parasite survival? *Oikos*. 118: 941–949.
- Sanchez-Bayo, F., and H. A. Tennekes. 2017. Assessment of ecological risks of agrochemicals requires a new framework. *Environ. Risk Assess. Remediat.* 1: 1–9.
- Schmid-Hempel, P., and R. Schmid-Hempel. 1993. Transmission of a pathogen in *Bombus terrestris*, with a note on division of labour in social insects. *Behav. Ecol. Sociobiol.* 33: 319–327.
- Shykoff, J. A., and P. Schmid-Hempel. 1991. Incidence and effects of four parasites in natural populations of bumble bees in Switzerland. *Apidologie*. 22: 117–125.
- Simon-Delso, N., V. Amaral-Rogers, L. P. Belzunces, J. M. Bonmatin, M. Chagnon, C. Downs, L. Furlan, D. W. Gibbons, C. Giorio, V. Girolami, et al. 2015. Systemic insecticides (neonicotinoids and fipronil): trends, uses, mode of action and metabolites. *Environ. Sci. Pollut. Res. Int.* 22: 5–34.
- Siviter, H., M. J. F. Brown, and E. Leadbeater. 2018a. Sulfoxaflor exposure reduces bumblebee reproductive success. *Nature*. 561: 109–112.
- Siviter, H., J. Koricheva, M. J. F. Brown, and E. Leadbeater. 2018b. Quantifying the impact of pesticides on learning and memory in bees. *J. Appl. Ecol.* 55: 2812–2821.