

3.2.P.5.1 Specification

AMLOZA (Amlodipine Besylate Tablets, USP) 5 mg / 10 mg — ANDA

Description	White, round, biconvex, uncoated tablets, debossed as applicable
Identification	HPLC retention time matches reference standard; UV/IR conforms
Assay	95.0–105.0% of label claim, by HPLC
Content uniformity	Per USP <905>, Acceptance Value criteria
Dissolution	Example method (for portfolio purposes): USP Apparatus 2 (Paddle), 50 rpm; medium: 900 mL Purified Water; temperature 37.0°C ± 0.5°C; sampling at 30 minutes; acceptance criterion: Not less than 80% (Q) of the labeled amount of amlodipine dissolved in 30 minutes. The final validated method would be established during formal method development, referencing the USP Amlodipine Tablets monograph and/or FDA product-specific dissolution guidance as applicable.
Related substances	Individual degradation product ≤ 0.2%; total degradation products ≤ 1.0% (indicative; to be finalized against ICH Q3B thresholds for max daily dose)
Water content	≤ [X]% w/w by Karl Fischer (limit to be set from stability data)
Weight variation / uniformity of dosage units	Per USP <905>
Microbial limits	Per USP <1111>, <61>, <62> — routine testing at every batch not typically required for this dosage form category provided development controls are demonstrated; tested if warranted per risk assessment

ACADEMIC PORTFOLIO EXERCISE — NOT FOR REGULATORY SUBMISSION