
a
PBMC
FTD
BT474
K562
HG002
UHRR
All
Pile−up + cap
+ CAGE or refTSS
+ not canonical TSS
+ protein altered
+ N−truncated
5,263
177
146
75
24
16
6,222
83
70
29
7
4
460
90
81
54
14
8
519
58
55
34
9
7
880
91
89
54
14
10
734
73
64
40
10
7
1
100
10,000
strict ISM TSSs
b
Same−exon random
Site
Capped pile−up
Uncapped pile−up
% within 10 nt of a CAGE peak
100
75
50
25
0
PBMC
FTD
BT474
K562
HG002
UHRR
c
% internal TSSs kept
% uncapped pile−ups kept
IsoSeq
LRAA
Mandalorion
ESPRESSO
Bambu
StringTie
IsoQuant
0
2
10
50
100
d
COMT, BT474: 271 −> 221 aa
COMT−L
COMT−S
Read starts
600
400
200
0
−200
−100
100
200
nt from the site
e
MBP: no qualifying S/L pair in LRAA v0.50.0 models
f
CORO1B, PBMC: 489 −> 171 aa
CORO1B−L
CORO1B−S
Read starts
nt from the site
g
RECQL4, BT474: 1208 −> 851 aa
RECQL4−L
RECQL4−S
Read starts
50
40
30
20
10
nt from the site
Site, capped
Site, not capped
Site, no cap information
Starts elsewhere
Through L intron