

a
Classical monocytes
Non-classical monocytes
Dendritic cells
B cells
T cells
CD4+
CD8+
NK
b
LINC01550
CEACAM21
ACYP1
SACM1L
SVBP
MIR4435−2HG
DHRS4−AS1
NMRAL1
GFUS
LPCAT1
PHF21A
IGFLR1
RDX
CNST
C0 C1 C2 C3 C4 C5 C6 C7 C8 C9 C10 C11 C12 C13 all cells
isoform fraction
>=5%
4%
3%
2%
1%
0%
c
AIF1^6633d
AIF1^a9489
Partial EF-hand Domain
Full EF-hand Domain
Superimposed
C-terminus Disordered Region
Shared Sequences
AIF^a9489-specific Sequences
Other Sequences
d
read 5' ends at the gene's TSS sites, per site (within 25 bp)
Naive CD4 T−2 (C7)
CD14 Mono (C1)
31,615,500
31,616,000
31,616,500
31,617,000
e
usage
AIF1−L
AIF1−S
C1 C11 C13 C6 C9 C3 C7 C0 C5 C12 C2 C4 C8 C10
isoform fraction
cells with reads
25%
50%
75%
100%
f
AIF1−L
AIF1−S
g
read 5' ends at the gene's TSS sites, per site (within 25 bp)
Naive CD4 T−2 (C7)
CD16 Mono (C11)
57,741,500
57,742,000
57,742,500
57,743,000
h
usage
SELENOH−U
SELENOH−D
C1 C11 C13 C6 C9 C3 C7 C0 C5 C12 C2 C4 C8 C10
isoform fraction
cells with reads
20%
40%
60%
80%
i
SELENOH−U
SELENOH−D