

a
Classical monocytes
Non-classical monocytes
Dendritic cells
B cells
T cells
CD4+
CD8+
NK
b
LINC01550
CEACAM21
LACTB
SACM1L
SVBP
MIR4435–2HG
DHRS4–AS1
SLC50A1
GFUS
LPCAT1
APTX
OXNAD1
CCNDBP1
PRPF38A
C0 C1 C2 C3 C4 C5 C6 C7 C8 C9 C10 C11 C12 C13 all cells
isoform fraction
>=5%
4%
3%
2%
1%
0%
c
AIF1-S
AIF1-L
Partial EF-hand Domain
Full EF-hand Domain
Superimposed
C-terminus Disordered Region
Shared Sequences
AIF1-L Specific Sequences
d
read 5' ends at the gene's TSS sites, per site (within 25 bp)
Naive CD4 T–2 (C7)
CD14 Mono (C1)
31,615,500
31,616,000
31,616,500
31,617,000
e
usage
AIF1–L
AIF1–S
C1 C11 C13 C6 C9 C3 C7 C0 C5 C12 C2 C4 C8 C10
isoform fraction
cells with reads
25%
50%
75%
100%
f
AIF1–L
AIF1–S
g
read 5' ends at the gene's TSS sites, per site (within 25 bp)
Naive CD4 T–2 (C7)
CD16 Mono (C11)
57,741,500
57,742,000
57,742,500
57,743,000
h
usage
SELENOH–U
SELENOH–D
C1 C11 C13 C6 C9 C3 C7 C0 C5 C12 C2 C4 C8 C10
isoform fraction
cells with reads
20%
40%
60%
80%
i
SELENOH–U
SELENOH–D