

MATH 243: Discussion Worksheet 1

Topics: Review of differentiation and integration; 12.1 Three-Dimensional Coordinate Systems

Please put away all electronic devices, including cell phones and calculators.

This content is protected and may not be shared, uploaded, or distributed.

1. Find the derivative of $f(x) = \cos(x^2 + e^x)$.
2. Given that $x \ln(y) + e^y = 3$, find $\frac{dy}{dx}$ by implicit differentiation.
3. Suppose that $f(x)$ is a differentiable function such that $f(1) = 7$ and $f'(1) = 4$. Find an equation of the tangent line to $y = \sqrt{4 + 3f(x)}$ at $x = 1$.
4. Evaluate the indefinite integrals.
 - (a) $\int e^{\tan x} \sec^2(x) dx$
 - (b) $\int (2x^4 + 1) \ln(x) dx$
5. Evaluate the definite integrals.
 - (a) $\int_0^{\pi/2} \sin^2 \theta \cos \theta d\theta$
 - (b) $\int_0^3 \frac{x}{\sqrt{x^2 + 1}} dx$
6. A particle is moving with acceleration at time t given by
$$a(t) = 3 \cos t - 2 \sin t.$$
Given that $s(0) = 0$ and $v(0) = 4$, determine the position of the particle $s(t)$ at any time t .
7. Consider the point $P(3, 4, 5)$.
 - (a) What is the projection of the point onto the xy -plane?
 - (b) What is the projection of the point onto the xz -plane?
 - (c) Find the length of the line segment \overline{OP} .
 - (d) Find the position vector for the point $P(3, 4, 5)$ in $\mathbf{i}, \mathbf{j}, \mathbf{k}$ -form.
8. (a) Find an equation of the sphere that passes through the point $(1, 8, 5)$ and has center $(3, 1, -3)$.
(b) Using your answer from part (a), find an equation describing the intersection of the sphere with the yz -plane. If the sphere does not intersect the yz -plane, write DNE.
9. Find an equation of a sphere if one of its diameters has endpoints at $(1, 2, 4)$ and $(4, 3, 10)$.

SOME USEFUL DEFINITIONS, THEOREMS, AND NOTATION:

Differentiation Formulas

$$\frac{d}{dx}(c) = 0$$

$$\frac{d}{dx}(b^x) = b^x \ln b$$

$$\frac{d}{dx}(f(x) + g(x)) = f'(x) + g'(x)$$

$$\frac{d}{dx}(x^n) = nx^{n-1}$$

$$\frac{d}{dx}(cf(x)) = cf'(x)$$

$$\frac{d}{dx}(f(x) - g(x)) = f'(x) - g'(x)$$

$$\frac{d}{dx}(e^x) = e^x$$

The Product Rule

$$\frac{d}{dx}(f(x)g(x)) = f'(x)g(x) + f(x)g'(x)$$

The Quotient Rule

$$\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$$

The Chain Rule

$$\frac{d}{dx}(f(g(x))) = f'(g(x))g'(x)$$

Derivatives of Trigonometric Functions

$$\frac{d}{dx}(\sin x) = \cos x$$

$$\frac{d}{dx}(\sec x) = \sec x \tan x$$

$$\frac{d}{dx}(\csc x) = -\csc x \cot x$$

$$\frac{d}{dx}(\tan x) = \sec^2 x$$

$$\frac{d}{dx}(\cos x) = -\sin x$$

$$\frac{d}{dx}(\cot x) = -\csc^2 x$$

The Chain Rule for the power of a function

$$\frac{d}{dx}[f(x)^n] = n[f(x)]^{n-1} f'(x)$$

Derivatives of Inverse Trigonometric Functions

$$\frac{d}{dx}(\arcsin x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\arctan x) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\arccos x) = -\frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}(\text{arccot } x) = -\frac{1}{1+x^2}$$

Derivatives of Logarithmic Functions

$$\frac{d}{dx}(\log_b x) = \frac{1}{x \ln b}$$

$$\frac{d}{dx}(\ln x) = \frac{1}{x}$$

$$\frac{d}{dx}[\ln f(x)] = \frac{f'(x)}{f(x)}$$

Integration Formulas

- a. If $n \neq -1$, $\int x^n dx = \frac{x^{n+1}}{n+1} + C$
- b. $\int e^x dx = e^x + C$
- c. $\int \sin x dx = -\cos x + C$
- d. $\int \sec^2 x dx = \tan x + C$
- e. $\int \sec x \tan x dx = \sec x + C$
- f. $\int \sec x dx = \ln |\sec x + \tan x| + C$
- g. $\int \tan x dx = \ln |\sec x| + C$
- i. $\int \frac{1}{x} dx = \ln |x| + C$
- j. $\int b^x dx = \frac{b^x}{\ln b} + C$
- k. $\int \cos x dx = \sin x + C$
- l. $\int \csc^2 x dx = -\cot x + C$
- m. $\int \csc x \cot x dx = -\csc x + C$
- n. $\int \csc x dx = -\ln |\csc x + \cot x| + C$
- o. $\int \cot x dx = \ln |\sin x| + C$

where C is any real constant.

The Substitution Rule: This rule is useful when the integrand contains a function and (a constant multiple of) its derivative. Let $u = g(x)$, so $du = g'(x) dx$. Then

$$\int f(g(x)) g'(x) dx = \int f(u) du.$$

Integration by Parts: This rule is useful for integrals that look like a product of two functions. Choose u and dv from the integrand so that $du = u' dx$ and $v = \int dv$. Then

$$\int u dv = uv - \int v du.$$

Suggested Textbook Problems

Section 12.1: 1-46