

Topics: 13.2 Derivatives and Integrals of Vector Functions; 13.3 Arc Length and Curvature; 13.4 Motion in Space: Velocity and Acceleration

Please put away all electronic devices, including cell phones and calculators.

This content is protected and may not be shared, uploaded, or distributed.

1. Find the derivative of the vector function

$$\mathbf{r}(t) = \langle e^{t^2+2t}, \ln(\cos(t)), t \arctan(t) \rangle.$$

2. Evaluate the definite integral

$$\int_0^1 \left(\frac{1}{t+1} \mathbf{i} + \frac{1}{t^2+1} \mathbf{j} + \frac{t}{t^2+1} \mathbf{k} \right) dt.$$

3. Find a vector equation of the tangent line to the curve

$$\mathbf{r}(t) = t \cos(2t) \mathbf{i} + 4t \mathbf{j} + \sin(t) \mathbf{k}$$

at the point corresponding to $t = \frac{\pi}{4}$.

4. Find the length of the curve $\mathbf{r}(t) = 2t \mathbf{i} + t^2 \mathbf{j} + \frac{1}{3}t^3 \mathbf{k}$ for the interval $0 \leq t \leq 1$.
5. A particle starts at the point $(0, 0, 3)$ and moves along the curve $\mathbf{r}(t) = \langle 3 \sin t, 4t, 3 \cos t \rangle$ in the direction of increasing t . Find the position of the particle after it has traveled a distance of 5π units.
6. Consider the vector function $\mathbf{r}(t) = \langle t, t^2, 4 \rangle$.
 - (a) Find the unit tangent vector $\mathbf{T}(t)$.
 - (b) Find the unit normal vector $\mathbf{N}(t)$.
 - (c) Find the curvature $\kappa(t)$.
7. Find the vectors \mathbf{T} , \mathbf{N} , and \mathbf{B} at the point $(1, \frac{2}{3}, 1)$ for the vector function

$$\mathbf{r}(t) = \langle t^2, \frac{2}{3}t^3, t \rangle.$$

8. A particle moves in space with position function

$$\mathbf{r}(t) = \langle t \ln t, t, e^{-t} \rangle.$$

Find the velocity, speed, and acceleration of the particle.

9. The acceleration of an object is given by $\mathbf{a}(t) = t \mathbf{i} + e^t \mathbf{j} + e^{-t} \mathbf{k}$. Find the object's position function $\mathbf{r}(t)$ if the initial velocity is $\mathbf{v}(0) = \mathbf{k}$ and the initial position is $\mathbf{r}(0) = \mathbf{j} + \mathbf{k}$.
10. At a point P , the velocity and acceleration vectors of a particle moving in space are $\mathbf{v} = \mathbf{i} - \mathbf{j} + \mathbf{k}$ and $\mathbf{a} = 2\mathbf{i} - 3\mathbf{k}$, respectively. Determine the curvature of the particle's path at P .

SOME USEFUL DEFINITIONS, THEOREMS, AND NOTATION

Arc Length. If \mathbf{r} is a smooth vector function on the interval $[a, b]$, then the arc length of the curve described by \mathbf{r} between the points with position vectors $\mathbf{r}(a)$ and $\mathbf{r}(b)$ is

$$L = \int_a^b |\mathbf{r}'(t)| dt.$$

Unit Tangent, Unit Normal, and Binormal Vectors. Let \mathbf{r} be a smooth vector function. The **unit tangent vector** at time t is defined by

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}.$$

The **(principal) unit normal vector** is defined by

$$\mathbf{N}(t) = \frac{\mathbf{T}'(t)}{|\mathbf{T}'(t)|}.$$

The **binormal vector** is defined by

$$\mathbf{B}(t) = \mathbf{T}(t) \times \mathbf{N}(t).$$

The vectors \mathbf{T} , \mathbf{N} , and \mathbf{B} are orthogonal unit vectors.

Curvature. The **curvature** of a smooth curve is given by

$$\kappa(t) = \left| \frac{d\mathbf{T}}{ds} \right| = \frac{|\mathbf{T}'(t)|}{|\mathbf{r}'(t)|} = \frac{|\mathbf{r}'(t) \times \mathbf{r}''(t)|}{|\mathbf{r}'(t)|^3}.$$

Curvature is a scalar quantity that measures how rapidly a smooth curve changes direction at a given point. Intuitively, it describes how sharply the curve bends.

Position, Velocity, and Acceleration. If a particle moves in space so that its position vector at time t is $\mathbf{r}(t)$, then the **velocity vector** of the particle is $\mathbf{v}(t) = \mathbf{r}'(t)$, its **speed** is $|\mathbf{v}(t)|$, and its **acceleration vector** is $\mathbf{a}(t) = \mathbf{r}''(t)$.

In terms of velocity and acceleration, the third curvature formula may be written as

$$\kappa(t) = \frac{|\mathbf{v}(t) \times \mathbf{a}(t)|}{|\mathbf{v}(t)|^3}.$$

Suggested Textbook Problems

Section 13.2: 1-44, 49-52

Section 13.3: 1-30, 40-45, 51, 52

Section 13.4: 1-32, 36