

13 Vector Functions





13.2

Derivatives and Integrals of Vector Functions



Derivatives

Derivatives (1 of 4)

The derivative \mathbf{r}' of a vector function \mathbf{r} is defined in much the same way as for real-valued functions:

$$1 \quad \frac{d\mathbf{r}}{dt} = \mathbf{r}'(t) = \lim_{h \rightarrow 0} \frac{\mathbf{r}(t+h) - \mathbf{r}(t)}{h}$$

if this limit exists. The geometric significance of this definition is shown in Figure 1.



Figure 1

Derivatives (2 of 4)

If the points P and Q have position vectors $\mathbf{r}(t)$ and $\mathbf{r}(t + h)$, then \overrightarrow{PQ} represents the vector $\mathbf{r}(t + h) - \mathbf{r}(t)$, which can therefore be regarded as a secant vector.

If $h > 0$, the scalar multiple $\left(\frac{1}{h}\right)(\mathbf{r}(t + h) - \mathbf{r}(t))$ has the same direction as

$\mathbf{r}(t + h) - \mathbf{r}(t)$. As $h \rightarrow 0$, it appears that this vector approaches a vector that lies on the tangent line.

For this reason, the vector $\mathbf{r}'(t)$ is called the **tangent vector** to the curve defined by \mathbf{r} at the point P , provided that $\mathbf{r}'(t)$ exists and $\mathbf{r}'(t) \neq \mathbf{0}$.

The **tangent line** to C at P is defined to be the line through P parallel to the tangent vector $\mathbf{r}'(t)$.

Derivatives (3 of 4)

The following theorem gives us a convenient method for computing the derivative of a vector function \mathbf{r} : just differentiate each component of \mathbf{r} .

2 Theorem If $\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}$, where f , g , and h are differentiable functions, then

$$\mathbf{r}'(t) = \langle f'(t), g'(t), h'(t) \rangle = f'(t)\mathbf{i} + g'(t)\mathbf{j} + h'(t)\mathbf{k}$$

A unit vector that has the same direction as the tangent vector is called the **unit tangent vector \mathbf{T}** and is defined by

$$\mathbf{T}(t) = \frac{\mathbf{r}'(t)}{|\mathbf{r}'(t)|}$$

Example 1

- (a) Find the derivative of $\mathbf{r}(t) = (1 + t^3) \mathbf{i} + te^{-t} \mathbf{j} + \sin 2t \mathbf{k}$.
- (b) Find the unit tangent vector at the point where $t = 0$.

Solution:

- (a) According to Theorem 2, we differentiate each component of \mathbf{r} :

$$\mathbf{r}'(t) = 3t^2 \mathbf{i} + (1 - t)e^{-t} \mathbf{j} + 2 \cos 2t \mathbf{k}$$

Example 1 – Solution

(b) Since $\mathbf{r}(0) = \mathbf{i}$ and $\mathbf{r}'(0) = \mathbf{j} + 2\mathbf{k}$, the unit tangent vector at the point $(1, 0, 0)$ is

$$\begin{aligned} \mathbf{T}(0) &= \frac{\mathbf{r}'(0)}{|\mathbf{r}'(0)|} \\ &= \frac{\mathbf{j} + 2\mathbf{k}}{\sqrt{1 + 4}} \\ &= \frac{1}{\sqrt{5}}\mathbf{j} + \frac{2}{\sqrt{5}}\mathbf{k} \end{aligned}$$

Derivatives (4 of 4)

Just as for real-valued functions, the **second derivative** of a vector function \mathbf{r} is the derivative of \mathbf{r}' , that is, $\mathbf{r}'' = (\mathbf{r}')'$.

For instance, the second derivative of the function, $\mathbf{r}(t) = \langle 2 \cos t, \sin t, t \rangle$, is

$$\mathbf{r}''(t) = \langle -2 \cos t, -\sin t, 0 \rangle$$



Differentiation Rules

Differentiation Rules (1 of 3)

The next theorem shows that the differentiation formulas for real-valued functions have their counterparts for vector-valued functions.

3 Theorem Suppose \mathbf{u} and \mathbf{v} are differentiable vector functions, c is a scalar, and f is a real-valued function. Then

$$1. \frac{d}{dt}[\mathbf{u}(t) + \mathbf{v}(t)] = \mathbf{u}'(t) + \mathbf{v}'(t)$$

$$4. \frac{d}{dt}[\mathbf{u}(t) \cdot \mathbf{v}(t)] = \mathbf{u}'(t) \cdot \mathbf{v}(t) + \mathbf{u}(t) \cdot \mathbf{v}'(t)$$

$$2. \frac{d}{dt}[c\mathbf{u}(t)] = c\mathbf{u}'(t)$$

$$5. \frac{d}{dt}[\mathbf{u}(t) \times \mathbf{v}(t)] = \mathbf{u}'(t) \times \mathbf{v}(t) + \mathbf{u}(t) \times \mathbf{v}'(t)$$

$$3. \frac{d}{dt}[f(t)\mathbf{u}(t)] = f'(t)\mathbf{u}(t) + f(t)\mathbf{u}'(t)$$

$$6. \frac{d}{dt}[\mathbf{u}(f(t))] = f'(t)\mathbf{u}'(f(t)) \quad (\text{Chain Rule})$$

Differentiation Rules (2 of 3)

We use Formula 4 to prove the following theorem.

4 Theorem if $|\mathbf{r}(t)| = c$ (a constant), then $\mathbf{r}'(t)$ is orthogonal to $\mathbf{r}(t)$ for all t .

PROOF

Since $\mathbf{r}(t) \cdot \mathbf{r}(t) = |\mathbf{r}(t)|^2 = c^2$

and c^2 is a constant, Formula 4 of Theorem 3 gives

$$0 = \frac{d}{dt}[\mathbf{r}(t) \cdot \mathbf{r}(t)] = \mathbf{r}'(t) \cdot \mathbf{r}(t) + \mathbf{r}(t) \cdot \mathbf{r}'(t) = 2\mathbf{r}'(t) \cdot \mathbf{r}(t)$$

Thus $\mathbf{r}'(t) \cdot \mathbf{r}(t) = 0$, which says that $\mathbf{r}'(t)$ is orthogonal to $\mathbf{r}(t)$.

Differentiation Rules (3 of 3)

Geometrically, this result says that if a curve lies on a sphere with center the origin, then the tangent vector $\mathbf{r}'(t)$ is always perpendicular to the position vector $\mathbf{r}(t)$. (See Figure 4.)




Figure 4



Integrals

Integrals (1 of 3)

The **definite integral** of a continuous vector function $\mathbf{r}(t)$ can be defined in much the same way as for real-valued functions except that the integral is a vector.

But then we can express the integral of \mathbf{r} in terms of the integrals of its component functions f , g , and h as follows.

$$\begin{aligned}\int_a^b \mathbf{r}(t) dt &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \mathbf{r}(t_i^*) \Delta t \\ &= \lim_{n \rightarrow \infty} \left[\left(\sum_{i=1}^n f(t_i^*) \Delta t \right) \mathbf{i} + \left(\sum_{i=1}^n g(t_i^*) \Delta t \right) \mathbf{j} + \left(\sum_{i=1}^n h(t_i^*) \Delta t \right) \mathbf{k} \right]\end{aligned}$$

Integrals (2 of 3)

So

$$\int_a^b \mathbf{r}(t) dt = \left(\int_a^b f(t) dt \right) \mathbf{i} + \left(\int_a^b g(t) dt \right) \mathbf{j} + \left(\int_a^b h(t) dt \right) \mathbf{k}$$

This means that we can evaluate an integral of a vector function by integrating each component function.

Integrals (3 of 3)

We can extend the Fundamental Theorem of Calculus to continuous vector functions as follows:

$$\int_a^b \mathbf{r}(t) dt = \left[\mathbf{R}(t) \right]_a^b = \mathbf{R}(b) - \mathbf{R}(a)$$

where \mathbf{R} is an antiderivative of \mathbf{r} , that is, $\mathbf{R}'(t) = \mathbf{r}(t)$.

We use the notation $\int \mathbf{r}(t) dt$ for indefinite integrals (antiderivatives).

Example 4

If $\mathbf{r}(t) = 2 \cos t \mathbf{i} + \sin t \mathbf{j} + 2t \mathbf{k}$, then

$$\begin{aligned}\int \mathbf{r}(t) dt &= \left(\int 2 \cos t \, dt \right) \mathbf{i} + \left(\int \sin t \, dt \right) \mathbf{j} + \left(\int 2t \, dt \right) \mathbf{k} \\ &= 2 \sin t \mathbf{i} - \cos t \mathbf{j} + t^2 \mathbf{k} + \mathbf{C}\end{aligned}$$

where \mathbf{C} is a vector constant of integration, and

$$\begin{aligned}\int_0^{\frac{\pi}{2}} \mathbf{r}(t) dt &= [2 \sin t \mathbf{i} - \cos t \mathbf{j} + t^2 \mathbf{k}]_0^{\frac{\pi}{2}} \\ &= 2\mathbf{i} + \mathbf{j} + \frac{\pi^2}{4} \mathbf{k}\end{aligned}$$