

Topics: 12.5 Equations of Lines and Planes; 13.1 Vector Functions and Space Curves

Please put away all electronic devices, including cell phones and calculators.

This content is protected and may not be shared, uploaded, or distributed.

1. Find the following equations of the line through the point $P(2, 2, 4)$ that is parallel to the vector $\mathbf{a} = 4\mathbf{i} - \mathbf{j} + 7\mathbf{k}$. For parts (a) and (b), use the parameter t .
 - (a) Vector equation
 - (b) Parametric equations
 - (c) Symmetric equations

The position vector of the point P is $\overrightarrow{OP} = 2\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}$. Since the line is parallel to \mathbf{a} , we may take the direction vector to be $\mathbf{v} = \mathbf{a} = 4\mathbf{i} - \mathbf{j} + 7\mathbf{k}$. Any point on the line has position vector \mathbf{r} .

(a) A vector equation of the line through P with direction vector \mathbf{v} is

$$\mathbf{r}(t) = \overrightarrow{OP} + t\mathbf{v} = (2\mathbf{i} + 2\mathbf{j} + 4\mathbf{k}) + t(4\mathbf{i} - \mathbf{j} + 7\mathbf{k}).$$

Equivalently,

$$\mathbf{r}(t) = (2 + 4t)\mathbf{i} + (2 - t)\mathbf{j} + (4 + 7t)\mathbf{k}.$$

(b) Parametric equations are

$$x(t) = 2 + 4t, \quad y(t) = 2 - t, \quad z(t) = 4 + 7t.$$

(c) Symmetric equations are

$$\frac{x - 2}{4} = \frac{y - 2}{-1} = \frac{z - 4}{7}.$$

2. (a) Find symmetric equations of the line through the point $P(3, 4, 0)$ that is perpendicular to both vectors $2\mathbf{i} + 2\mathbf{j}$ and $\mathbf{j} + \mathbf{k}$.

A direction vector \mathbf{v} for the desired line must be perpendicular to both given vectors, so we can take the cross product:

$$\mathbf{v} = (2\mathbf{i} + 2\mathbf{j}) \times (\mathbf{j} + \mathbf{k}) = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 2 & 0 \\ 0 & 1 & 1 \end{vmatrix} = 2\mathbf{i} - 2\mathbf{j} + 2\mathbf{k}.$$

Thus a direction vector is $\langle 2, -2, 2 \rangle$. The symmetric equations are

$$\frac{x - 3}{2} = \frac{y - 4}{-2} = \frac{z - 0}{2}.$$

- (b) Find the point where the line from part (a) intersects the yz -plane.

The yz -plane is given by $x = 0$. Using the symmetric equations,

$$\frac{0 - 3}{2} = \frac{y - 4}{-2} = \frac{z - 0}{2}.$$

So

$$-\frac{3}{2} = \frac{y - 4}{-2} \quad \text{and} \quad -\frac{3}{2} = \frac{z}{2}.$$

Solving gives $y = 7$ and $z = -3$. Therefore, the line intersects the yz -plane at

$$(0, 7, -3).$$

3. Determine whether the lines L_1 and L_2 are parallel, intersecting, or skew.

(a) $L_1 : \mathbf{r}(t) = \langle -1 + 3t, 2 + 4t, 3 - 2t \rangle, \quad L_2 : \frac{x - 1}{2} = \frac{y}{-3} = \frac{z + 1}{-3}$

Write L_2 in parametric form by letting

$$\frac{x - 1}{2} = \frac{y}{-3} = \frac{z + 1}{-3} = s.$$

Then

$$x = 1 + 2s, \quad y = -3s, \quad z = -1 - 3s,$$

so a direction vector for L_2 is $\mathbf{v}_2 = \langle 2, -3, -3 \rangle$. A direction vector for L_1 is $\mathbf{v}_1 = \langle 3, 4, -2 \rangle$.

Since \mathbf{v}_1 is not a scalar multiple of \mathbf{v}_2 , the lines are not parallel. To check whether they intersect, solve

$$-1 + 3t = 1 + 2s, \quad 2 + 4t = -3s, \quad 3 - 2t = -1 - 3s.$$

From the first two equations, we obtain $t = \frac{2}{17}$ and $s = -\frac{14}{17}$. Substituting into the third equation gives

$$3 - 2\left(\frac{2}{17}\right) \neq -1 - 3\left(-\frac{14}{17}\right),$$

so the system is inconsistent and the lines do not intersect. Therefore, the lines are **skew**.

(b) $L_1 : x = 2t, y = -3 + t, z = 5 - t, \quad L_2 : x = 3 - 3s, y = 2 - \frac{3}{2}s, z = \frac{3}{2}s$

Direction vectors are $\mathbf{v}_1 = \langle 2, 1, -1 \rangle$ and $\mathbf{v}_2 = \langle -3, -\frac{3}{2}, \frac{3}{2} \rangle$. Since

$$\mathbf{v}_1 = -\frac{2}{3}\mathbf{v}_2,$$

the lines are **parallel**. Note that they are not the same line (for example, $(0, -3, 5)$ lies on L_1 when $t = 0$, but it does not satisfy the equations of L_2 for any s).

4. Find an equation of the plane through the points $A(0, 1, 2)$, $B(1, 2, 3)$, and $C(2, 3, 5)$.

First find direction vectors in the plane:

$$\overrightarrow{AB} = \langle 1, 1, 1 \rangle, \quad \overrightarrow{AC} = \langle 2, 2, 3 \rangle.$$

A normal vector is their cross product:

$$\mathbf{n} = \overrightarrow{AB} \times \overrightarrow{AC} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 1 & 1 & 1 \\ 2 & 2 & 3 \end{vmatrix} = (1 \cdot 3 - 1 \cdot 2)\mathbf{i} - (1 \cdot 3 - 1 \cdot 2)\mathbf{j} + (1 \cdot 2 - 1 \cdot 2)\mathbf{k} = \langle 1, -1, 0 \rangle.$$

Using point $A(0, 1, 2)$, the plane equation is

$$1(x - 0) + (-1)(y - 1) + 0(z - 2) = 0.$$

This can be rewritten as:

$$x - y = -1.$$

5. Find an equation of the plane through the points $(0, -2, 5)$ and $(-1, 3, 1)$ that is perpendicular to the plane $2z = 5x + 4y$.

Rewrite the given plane as $5x + 4y - 2z = 0$, so a normal vector is $\mathbf{n}_1 = \langle 5, 4, -2 \rangle$.

Let $A(0, -2, 5)$ and $B(-1, 3, 1)$. Then

$$\overrightarrow{AB} = \langle -1, 5, -4 \rangle$$

lies in the desired plane, so it is perpendicular to the desired plane's normal vector \mathbf{n}_2 . Also, since the planes are perpendicular, \mathbf{n}_2 is perpendicular to \mathbf{n}_1 . Hence \mathbf{n}_2 is perpendicular to both \overrightarrow{AB} and \mathbf{n}_1 , so we can take

$$\mathbf{n}_2 = \overrightarrow{AB} \times \mathbf{n}_1 = \langle -1, 5, -4 \rangle \times \langle 5, 4, -2 \rangle = \langle 6, -22, -29 \rangle.$$

Using point $A(0, -2, 5)$, the plane equation is

$$6(x - 0) - 22(y + 2) - 29(z - 5) = 0.$$

This can be rewritten as:

$$6x - 22y - 29z = -101.$$

6. Find parametric equations for the line of intersection of the planes

$$2x + 3y + 5z = 7 \quad \text{and} \quad x - y + 2z = 3.$$

Normal vectors to the planes are

$$\mathbf{n}_1 = \langle 2, 3, 5 \rangle, \quad \mathbf{n}_2 = \langle 1, -1, 2 \rangle.$$

A direction vector for the line of intersection is

$$\mathbf{v} = \mathbf{n}_1 \times \mathbf{n}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 2 & 3 & 5 \\ 1 & -1 & 2 \end{vmatrix} = 11\mathbf{i} + \mathbf{j} - 5\mathbf{k} = \langle 11, 1, -5 \rangle.$$

To find a point on the line, solve the system: Setting $y = 0$ gives

$$2x + 5z = 7, \quad x + 2z = 3,$$

which yields $x = 1$ and $z = 1$. Thus $P_0(1, 0, 1)$ is on the line. Therefore,

$$\langle x, y, z \rangle = \langle 1, 0, 1 \rangle + t\langle 11, 1, -5 \rangle$$

so parametric equations are

$$x = 1 + 11t, \quad y = t, \quad z = 1 - 5t.$$

7. Find an equation of the plane through $P(7, -2, -4)$ that is parallel to the plane $z = 4x - 5y$.

Rewrite the equation of the given plane as $4x - 5y - z = 0$, so a normal vector is $\mathbf{n} = \langle 4, -5, -1 \rangle$.

A parallel plane through $P(7, -2, -4)$ has the same normal vector:

$$4(x - 7) - 5(y + 2) - (z + 4) = 0.$$

This can be rewritten as:

$$4x - 5y - z = 42.$$

8. Find an equation of the plane that passes through the point $P(10, -1, 5)$ and contains the line with symmetric equations $\frac{x}{4} = y + 6 = \frac{z}{5}$.

Setting $\frac{x}{4} = y + 6 = \frac{z}{5} = t$, we get a parametric form of the line:

$$x = 4t, \quad y = t - 6, \quad z = 5t,$$

so a direction vector is $\mathbf{v}_1 = \langle 4, 1, 5 \rangle$. Taking $t = 0$ gives a point on the line: $A(0, -6, 0)$.

The vector from A to P is

$$\mathbf{v}_2 = \overrightarrow{AP} = \langle 10 - 0, -1 - (-6), 5 - 0 \rangle = \langle 10, 5, 5 \rangle,$$

which also lies in the plane. Therefore a normal vector is

$$\mathbf{n} = \mathbf{v}_1 \times \mathbf{v}_2 = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ 4 & 1 & 5 \\ 10 & 5 & 5 \end{vmatrix} = \langle -20, 30, 10 \rangle.$$

Using point $P(10, -1, 5)$, the plane equation is

$$-20(x - 10) + 30(y + 1) + 10(z - 5) = 0.$$

This can be rewritten as:

$$-20x + 30y + 10z = -180.$$

9. Determine whether the line L given by $\mathbf{r}(t) = \langle -2t, 2 + 7t, -1 - 4t \rangle$ intersects the plane given by $4x + 9y - 2z + 8 = 0$.

Write parametric equations for L :

$$x = -2t, \quad y = 2 + 7t, \quad z = -1 - 4t.$$

Substitute into the plane equation:

$$4(-2t) + 9(2 + 7t) - 2(-1 - 4t) + 8 = 0$$

$$-8t + 18 + 63t + 2 + 8t + 8 = 0 \implies 63t + 28 = 0 \implies t = -\frac{4}{9}.$$

Since a real value of t exists, the line intersects the plane. The intersection point is

$$(x, y, z) = \left(-2 \left(-\frac{4}{9} \right), 2 + 7 \left(-\frac{4}{9} \right), -1 - 4 \left(-\frac{4}{9} \right) \right) = \left(\frac{8}{9}, -\frac{10}{9}, \frac{7}{9} \right).$$

10. Find the domain of the vector function

$$\mathbf{r}(t) = \left\langle \ln(t+1), \frac{t}{4-t^2}, \sqrt{4-t} \right\rangle.$$

We first find the domain of each component of the vector function.

The domain of $\ln(t+1)$ is given by $t+1 > 0$, so it is the set $(-1, \infty)$.

The domain of $\frac{t}{4-t^2} = \frac{t}{(2+t)(2-t)}$ is given by $t \neq 2$ and $t \neq -2$. In interval notation, that is $(-\infty, -2) \cup (-2, 2) \cup (2, \infty)$.

The domain of $\sqrt{4-t}$ is given by $4-t \geq 0$, so it is the set $(-\infty, 4]$.

The domain of $\mathbf{r}(t)$ is the intersection of the domains of its components. So the domain of $\mathbf{r}(t)$ is

$$(-1, 2) \cup (2, 4].$$

11. Evaluate the limit $\lim_{t \rightarrow 1} \mathbf{r}(t)$, where

$$\mathbf{r}(t) = \left\langle \frac{t^2 - 1}{t^2 - 3t + 2}, \frac{t - 1}{\sqrt{t + 3} - 2}, \frac{\sin(t - 1)}{t - 1} \right\rangle.$$

To find the limit of a vector-valued function, we must compute

$$\lim_{t \rightarrow 1} \mathbf{r}(t) = \left\langle \lim_{t \rightarrow 1} \frac{t^2 - 1}{t^2 - 3t + 2}, \lim_{t \rightarrow 1} \frac{t - 1}{\sqrt{t + 3} - 2}, \lim_{t \rightarrow 1} \frac{\sin(t - 1)}{t - 1} \right\rangle.$$

For the first component,

$$\lim_{t \rightarrow 1} \frac{t^2 - 1}{t^2 - 3t + 2} = \lim_{t \rightarrow 1} \frac{(t+1)(t-1)}{(t-1)(t-2)} = \lim_{t \rightarrow 1} \frac{t+1}{t-2} = \frac{2}{-1} = -2.$$

For the second component, rationalize:

$$\lim_{t \rightarrow 1} \frac{t-1}{\sqrt{t+3}-2} = \lim_{t \rightarrow 1} \frac{(t-1)(\sqrt{t+3}+2)}{(t+3)-4} = \lim_{t \rightarrow 1} (\sqrt{t+3}+2) = \sqrt{4}+2 = 4.$$

For the third component, use L'Hôpital's rule:

$$\lim_{t \rightarrow 1} \frac{\sin(t-1)}{t-1} = \lim_{t \rightarrow 1} \frac{\cos(t-1)}{1} = \frac{\cos(0)}{1} = 1$$

Therefore,

$$\lim_{t \rightarrow 1} \mathbf{r}(t) = \langle -2, 4, 1 \rangle.$$

SOME USEFUL DEFINITIONS, THEOREMS, AND NOTATION

Equations of a line. The line through the point (x_0, y_0, z_0) and parallel to the vector $\langle a, b, c \rangle$ has **vector equation**

$$\mathbf{r}(t) = \langle x_0, y_0, z_0 \rangle + t\langle a, b, c \rangle.$$

The **parametric equations** for this line are

$$x = x_0 + at \quad y = y_0 + bt \quad z = z_0 + ct.$$

We can eliminate t to obtain the **symmetric equations**

$$\frac{x - x_0}{a} = \frac{y - y_0}{b} = \frac{z - z_0}{c}.$$

provided a , b , and c are all nonzero.

Equations of a plane. The plane through (x_0, y_0, z_0) with normal vector $\langle a, b, c \rangle$ is given by

$$a(x - x_0) + b(y - y_0) + c(z - z_0) = 0.$$

By collecting terms, we can rewrite this equation in the form

$$ax + by + cz = d$$

where $d = ax_0 + by_0 + cz_0$ is a constant. Two planes are parallel if their normal vectors are parallel, and they are perpendicular if their normal vectors are orthogonal.

Vector functions. A **vector-valued function** (or **vector function**) is a function whose domain is a set of real numbers and whose range is a set of vectors. In three dimensions, a vector-valued function is written as

$$\mathbf{r}(t) = \langle f(t), g(t), h(t) \rangle = f(t)\mathbf{i} + g(t)\mathbf{j} + h(t)\mathbf{k}.$$

where f , g , and h are real-valued **component functions**. The domain of $\mathbf{r}(t)$ is the set of all values t for which $f(t)$, $g(t)$, and $h(t)$ are all defined. Limits of vector functions are defined component-wise: For a real number a ,

$$\lim_{t \rightarrow a} \mathbf{r}(t) = \left\langle \lim_{t \rightarrow a} f(t), \lim_{t \rightarrow a} g(t), \lim_{t \rightarrow a} h(t) \right\rangle,$$

provided the three component limits exist.

Suggested Textbook Problems

Section 12.5: 1-72

Section 12.6: 1-49

Section 13.1: 1-58