

## MATH 243 Quiz 1 Solutions

1. From the cross product product rule, We have  $\mathbf{w} = \mathbf{r}' \times \mathbf{T} + \mathbf{r} \times \mathbf{T}'$ . From  $\mathbf{T} = \frac{\mathbf{r}'}{\|\mathbf{r}'\|}$ , the vectors  $\mathbf{r}'$ ,  $\mathbf{T}$  are parallel, so the 1st term is  $\mathbf{0}$ .

From  $\mathbf{N} = \frac{\mathbf{T}'}{\|\mathbf{T}'\|}$ , we can rewrite the last term as  $\|\mathbf{T}'\|(\mathbf{r} \times \mathbf{N})$ . So  $\mathbf{w} = \|\mathbf{T}'\|(\mathbf{r} \times \mathbf{N})$  is parallel to  $\mathbf{r} \times \mathbf{N}$ , which is perpendicular to  $\mathbf{N}$  since the cross product of two vectors is perpendicular to both of the original vectors. Thus, the answer is D.

Since  $\mathbf{w}$  is perpendicular to  $\mathbf{r}$ , B is false. If  $\mathbf{r} \neq \mathbf{0}$ , then  $\mathbf{w} \neq \mathbf{0}$ , so it can't be both perpendicular and parallel to  $\mathbf{r}$ , meaning C is false. Finally, A is false because the unit tangent to  $\mathbf{v}$  is  $\mathbf{z} = \frac{\mathbf{v}'}{\|\mathbf{v}'\|}$ , which has no relation to  $\mathbf{w}$ . You can see that  $\mathbf{w} \neq \mathbf{z}$  by choosing some  $\mathbf{r}$  and calculating both vectors.

2.  $\int \mathbf{r}(t) dt = \langle t^2, \frac{t^3}{3}, 2t \rangle + \mathbf{c}$ , so A is false since the vector of integration is not present. B is true by taking the derivative. We compute  $\|\mathbf{r}(t)\| = \sqrt{t^4 + 4t^2 + 4} = \sqrt{(t^2 + 2)^2} = t^2 + 2$  to see that C is true. We can compute  $\|\mathbf{s}'(t)\| = \|\langle 0, 2, 0 \rangle\| = 2$  to see that the arc length represented by  $\mathbf{s}$  from  $t = 0$  to  $t = 1$  is  $\int_0^1 \|\mathbf{s}'(t)\| dt = 2 \neq 4$ , so D is false. Thus, the correct options are B, C.

3. Recall that  $\|\mathbf{u} \cdot \mathbf{v}\| = \|u\|\|v\|\cos(\theta)$  and  $\|\mathbf{u} \times \mathbf{v}\| = \|u\|\|v\|\sin(\theta)$ . By dividing,  $\tan(\theta) = \frac{\|\mathbf{u} \times \mathbf{v}\|}{\|\mathbf{u} \cdot \mathbf{v}\|} = \frac{6+3\sqrt{3}}{3} = 2 + \sqrt{3}$ , so  $\theta = \tan^{-1}(2 + \sqrt{3}) = 75^\circ$  and the final answer is 75.

To see that  $\tan(75^\circ) = 2 + \sqrt{3}$ , you can remember the value from your classes on trigonometry, remember the value  $\tan(15^\circ) = 2 - \sqrt{3}$  and do  $\tan(75^\circ) = \frac{1}{\tan 15^\circ}$ . by remembering/using  $\tan(90 - x) = \frac{\sin(90-x)}{\cos(90-x)} = \frac{\cos(x)}{\sin(x)} = \cot(x)$ , or use the sum of angles formula for  $\tan$  :

$$\tan(30 + 45) = \frac{\tan(30) + \tan(45)}{1 - \tan(30)\tan(45)} = \frac{1/\sqrt{3} + 1}{1 - 1/\sqrt{3}} = 2 + \sqrt{3}.$$

There are even more ways to do this, but these are the easiest.

4. Let  $\mathbf{r}(t) = \langle x(t), y(t), z(t) \rangle$ . By trying to match the points up, we notice that  $\mathbf{r}(0) = (0, 1, 12)$ ,  $\mathbf{r}(1) = (e^4 \sin(3), e^4 \cos(3), 12)$ , so the arc length we seek is  $L = \int_0^1 \|\mathbf{r}'(t)\| dt$ .

Now we compute the magnitude:  $\mathbf{r}'(t) = \langle e^{4t}(3 \cos(3t) + 4 \sin(3t)), e^{4t}(-3 \sin(3t) + 4 \cos(3t)), 0 \rangle = e^{4t} \langle 3 \cos(3t) + 4 \sin(3t), -3 \sin(3t) + 4 \cos(3t), 0 \rangle$ , so

$$\|\mathbf{r}'(t)\| = e^{4t} \sqrt{(3 \cos(3t) + 4 \sin(3t))^2 + (-3 \sin(3t) + 4 \cos(3t))^2} = e^{4t} \sqrt{25 \sin^2(3t) + 25 \cos^2(3t)} = 5e^{4t}.$$

Plug this in to get  $L = \int_0^1 5e^{4t} dt = \frac{5}{4}(e^4 - 1)$ , so  $(p, q, r, s) = (5, 4, 4, 1)$  and the answer is 5441.

5. For shorthand, we write  $s = \sin, c = \cos$ . Compute  $\mathbf{r}'(t) = (e^t(s + c), 0, e^t(c - s)) = e^t(c + s, 0, c - s)$  and


$$\|\mathbf{r}'(t)\| = e^t \|(c + s, 0, c - s)\| = e^t \sqrt{(c + s)^2 + (c - s)^2} = e^t \sqrt{2s^2 + 2c^2} = e^t \sqrt{2}.$$

By definition,  $\mathbf{T} = \frac{\mathbf{r}'}{\|\mathbf{r}'\|} = \frac{1}{\sqrt{2}}(\cos t + \sin t, 0, \cos t - \sin t)$ . Compute  $\mathbf{T}' = \frac{1}{\sqrt{2}}(-s + c, 0, -s - c)$  and  $\|\mathbf{T}\| = 1$ , so  $\mathbf{N} = \frac{\mathbf{T}'}{\|\mathbf{T}'\|} = \frac{1}{\sqrt{2}}(-\sin(t) + \cos(t), 0, -\sin(t) - \cos(t))$ . Finally, compute  $\mathbf{B} = \mathbf{T} \times \mathbf{N} = \frac{1}{2}((c + s, 0, c - s) \times (-s + c, 0, -s - c)) = \frac{1}{2}(0, 2, 0) = (0, 1, 0)$ .

We saw one definition and two formulas for curvature. Since we have already found  $\|\mathbf{r}'\|$  and  $\|\mathbf{T}'\|$ , the

formula  $\kappa = \frac{\|\mathbf{T}'\|}{\|\mathbf{r}'\|} = \frac{1}{e^t\sqrt{2}} = e^{-t}\sqrt{2}$  is most convenient.

6. Let  $\theta$  be the angle between  $P_1$  and  $P_2$ . Draw a 2D cross section of the planes intersecting and draw normal vectors  $\mathbf{n}_1, \mathbf{n}_2$  to the planes sticking out the intersection point in the cross section. Your diagram may look something like the one below:



From the equations of the planes, we obtain  $\mathbf{n}_1 = (2, -1, 1), \mathbf{n}_2 = (1, 1, 2)$ .

If  $\mathbf{n}_1, \mathbf{n}_2$  are chosen to be on opposite sides of the angle like in the diagram, you get that the angle between  $\mathbf{n}_1, \mathbf{n}_2$  is  $180 - \theta$ . Thus,  $-\cos(\theta) = \cos(180 - \theta) = \frac{\mathbf{n}_1 \cdot \mathbf{n}_2}{\|\mathbf{n}_1\| \|\mathbf{n}_2\|} = \frac{(2, -1, 1) \cdot (1, 1, 2)}{\|(2, -1, 1)\| \|(1, 1, 2)\|} = \frac{3}{6} = 0.5$ , so  $\theta = \cos^{-1}(-0.5) = 120^\circ$ . Since we want the smaller intersection angle, our answer is  $180 - 120 = \boxed{60^\circ}$ .

To find a plane  $P$  perpendicular to  $P_1, P_2$ , note that by applying our diagram to perpendicular planes, we see that two planes are perpendicular if and only if their normal vectors are perpendicular. So we need  $P$  to have a normal vector perpendicular to  $\mathbf{n}_1, \mathbf{n}_2$ . We may take  $\mathbf{n} = \mathbf{n}_1 \times \mathbf{n}_2 = (-3, -3, 3)$ .

We're given that the origin  $(0, 0, 0)$  lies on the plane, so one equation for  $P$  is  $-3(x-0) - 3(y-0) + 3(z-0) = 0$ , which simplifies to  $\boxed{x + y = z}$ .