

FYS 4150 - Computational Physics

Project 1: Solving Poisson's equation in one dimension

MAREN RASMUSSEN

MARKUS LEIRA ASPRUSTEN

METIN SAN

4. September 2018

ABSTRACT

This project involves solving the one-dimensional Poisson equation with Dirichlet boundary conditions using two different algorithms. The first method is the tridiagonal matrix algorithm while the second is the LU decomposition. The conclusion of the project is that a specialized version of the tridiagonal algorithm is much faster.

1. INTRODUCTION

Hello, here is some text without a meaning. This text should show what a printed text will look like at this place. If you read this text, you will get no information. Really? Is there no information? Is there a difference between this text and some nonsense like "Huardest gefburn"? Kjift – not at all! A blind text like this gives you information about the selected font, how the letters are written and an impression of the look. This text should contain all letters of the alphabet and it should be written in of the original language. There is no need for special content, but the length of words should match the language.

2. THEORY

2.1. The Poisson Equation. Poisson's equation is a classical and well known differential equation from electromagnetism. The equation describes the potential field Φ generated from a charge distribution $\rho(\mathbf{r})$. For three dimensions, the equation is given by

$$\nabla^2 \Phi = -4\pi\rho(\mathbf{r}),$$

where $\nabla =$ is the Laplace operator. If both Φ and $\rho(\mathbf{r})$ is spherically symmetric, the equation can be simplified to an one dimensional equation in r ,

$$\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d\Phi}{dr} \right) = -4\pi\rho(r).$$

By substituting $\Phi = \phi(r)/r$, we can simplify the equation even more, giving

$$\frac{d^2\phi}{dr^2} = -4\pi\rho(r).$$

The final equation can be written in a general form by letting $\phi \rightarrow u$ and $r \rightarrow x$, and defining the right hand side of the equation as f ,

$$-u''(x) = f(x).$$

In this study we will assume that the source term is $f(x) = 100e^{-10x}$ with $x \in (0, 1)$. We will also assume that we have Dirichlet boundary conditions, such that $u(0) = u(1) = 0$. With these assumptions the equation have an exact solution on the form $u(x) = 1 - (1 - e^{-10})x - e^{-10x}$.

2.2. Discretization of the problem.