Exercises from Ch. 1: Groups, Lang

Matthew Gergley

Last Updated: March 6, 2026

Disclaimer: The proofs given are completed by myself. Thus, they are also reviewed by myself to ensure no logic flaws and incorrect arguments/deductions/conclusions. If you find any error in my proofs, please do not hesitate to contact me so we can discuss any potential errors and I can update the document as needed.

Exercises

Exercise 0.1. (Exercise 1) Show that every group of order ≤ 5 is abelian.

Proof. First, we will take care of the cases where $|G|$ is prime (i.e. 2, 3, 5). We prove this by showing that the groups of these orders are cyclic and thus abelian (cyclic $\Rightarrow$ abelian). Suppose that $|G| = p \leq 5$, a prime. Then, by Lagrange's Theorem, for all $x \in G$, $|x| \mid |G|$. Hence, $|x| \mid p$. Therefore, $|x| = 1$ or $|x| = p$. If $|x| = 1$, then $x = e$ and we get $|G| = 1$ which is not a prime, but is also trivially abelian. If $|x| = p$, then let $G = \langle x \rangle$. Thus, $G = \left\{e = x^{|G|},\, x,\, x \cdot x = x^2\,, x \cdot x \cdot x = x^3,\, \ldots\,, x^{|G|-1}\right\}$. Clearly, G is abelian since $x \cdot x^j = x^j \cdot x$ for all $x \in G$ and $1 \leq j \leq |G|$. Thus, if $|G| \leq 5$ is prime, then G is abelian. Next suppose $|G| = 4 = 2^2$. We prove a general case that all groups of order p^2 for a prime p are abelian and thus it would follow that G is abelian if $|G| = 4$.
Suppose $|G| = p^2$ for a prime p. Hence, G is a $p - group$ and hence $Z(G)$ is not trivial. Recall that the class equation is defined as

$$|G| = |Z(G)| + \sum_{x \in C} (G\colon G_x)$$

where C is a set of representatives for the distinct conjugacy classes and G_x is the isotropy group of x in G defined by $G_x = \{y \in G :\ x \cdot y = y\}$. But here we can consider G acting on itself by conjugation and thus $G_x = \{y \in G :\ yxy^{-1} = x\} = C_G(x) = \{g \in G :\ gx = xg\}$. Each term in $\sum\limits_{x \in C} (G\colon G_x) > 1$ and divides $|G| = p^2$. Hence, $\sum\limits_{x \in C} (G\colon G_x) \equiv 0 \pmod p$. Therefore, $|G| = |Z(G)| + kp$ ($k \in \mathbb{Z}^+$, and $kp = \sum (G\colon G_x)$). Then, $|Z(G)| = |G| - kp \Rightarrow |Z(G)| \equiv 0 \pmod p$. Thus, $p \mid |Z(G)| \Rightarrow |Z(G)| = p$ or p^2 (since $|Z(G)| \leq |G|$).
<u>Case 1:</u> Suppose $|Z(G)| = p^2$. Then $|Z(G)| = |G| \Rightarrow Z(G) = G$. Thus, G is abelian.
<u>Case 2:</u> Suppose $|Z(G)| = p$ Consider the quotient group $G/Z(G)$ (since $Z(G) \triangleleft G$). Thus, $|G/Z(G)| = \frac{|G|}{|Z(G)|} = \frac{p^2}{p} = p$. Hence, $G/Z(G)$ is cyclic by above and thus $G/Z(G)$ is abelian. Now we will show that the fact that $G/Z(G)$ is abelian implies that G is abelian.
Since $G/Z(G)$ is cyclic, $G/Z(G) = \langle gZ(G) \rangle$. Thus, $xZ(G) = (gZ(G))^k = g^k Z(G)$ for some $k \in \mathbb{Z}$. By coset equality, $xZ(G) = g^k Z(G) \iff x(g^k)^{-1} \in Z(G) \iff x = g^k z_1$ for some $z_1 \in Z(G)$. Hence, every element of $x \in G$ can be written as $x = g^k z_1$ for some $k \in \mathbb{Z}$ and $z_1 \in Z(G)$. Similarly, let $y \in G$ and thus $y = g^\ell z_2$ for some $\ell \in \mathbb{Z}$ and $z_2 \in Z(G)$. We want to show that $xy = yx$. Then,

$$xy = g^k z_1 g^\ell z_2 = g^{k+\ell} z_1 z_2 = g^{\ell+k} z_2 z_1 = g^\ell z_2 g^k z_1 = yx$$

Therefore, G is abelian and note that we were able to commute elements since $z_1, z_2 \in Z(G)$. Thus, we have shown that if $G/Z(G)$ is cyclic, then G is abelian. We are done.

□

Exercise 0.2. (Exercise 3)

Let G be a group. A **commutator** in G is an element of the form $aba^{-1}b^{-1}$ with $a, b \in G$. Let G^cbe the subgroup generated by the commutators. Then G^c is called the **commutator subgroup**. Show that G^c is normal. Show that any homomorphism of G into an abelian group factors through G/G^c.

Proof. By definition, $G^c = \{[a_1, b_1]^{c_1} \cdots [a_n, b_n]^{c_n} : n \in \mathbb{N},\ a_i, b_i \in G,\ c_i = \pm 1\}$. Thus, by construction, $G^c \leqslant G$. We show that this subgroup is normal. First, observe that for a single commutator,

$$\begin{aligned} g[a,b]g^{-1} &= g(aba^{-1}b^{-1})g^{-1} \\ &= ga(g^{-1}g)b(g^{-1}g)a^{-1}(g^{-1}g)b^{-1}g^{-1} \\ &= (gag^{-1})(gbg^{-1})(ga^{-1}g^{-1})(gb^{-1}g^{-1}) \\ &= \left[gag^{-1}, gbg^{-1}\right] \in G^c \end{aligned}$$

Conjugation is a homomorphism (distributes over products), and thus, since an element of G^c is a finite product of commutators, we have

$$\begin{aligned} g\left([a_1,b_1]^{c_1}[a_2,b_2]^{c_2}\cdots[a_n,b_n]^{c_n}\right)g^{-1} &= g\left(\left(a_1b_1a_1^{-1}b_1^{-1}\right)\cdots\left(a_nb_na_n^{-1}b_n^{-1}\right)\right)g^{-1} \\ &= g\left(a_1b_1a_1^{-1}b_1^{-1}\right)g^{-1}g(a_2b_2a_2^{-1}b_2^{-1})g^{-1}\cdots g(a_nb_na_n^{-1}b_n^{-1})g^{-1} \\ &= \left(g[a_1,b_1]g^{-1}\right)\left(g[a_2,b_2]g^{-1}\right)\cdots\left(g[a_n,b_n]g^{-1}\right) \end{aligned}$$

and our previous remarks, for all $1 \leq i \leq n$, $g[a_i, b_i]g^{-1} = \left[ga_ig^{-1}, gb_ig^{-1}\right]$. Then we get

$$g\left([a_1,b_1]\cdots[a_n,b_n]\right)g^{-1} = \left[ga_1g^{-1}, gb_1g^{-1}\right]\cdots\left[ga_ng^{-1}, gb_ng^{-1}\right] \in G^c$$

Note that since conjugation preserves products and inverses, we do not necessarily need to address the exponents $c_i = \pm 1$. Thus, $gG^cg^{-1} \subseteq G^c$ for all $g \in G$. Hence $G^c \triangleleft G$.
Let $f\colon G \to A$ be a homomorphism of G into an abelian group A. We want to show that the following diagram commutes:

$$\begin{tikzcd} G \arrow[rr, "f"] \arrow[dr, "\pi"'] & & A \\ & G/G^c \arrow[ur, "\bar{f}"'] & \end{tikzcd}$$

Here π is the canonical homomorphism (surjective). Observe that since f is a homomorphism and A is abelian,

$$\begin{aligned} f\left([a,b]\right) = f\left(aba^{-1}b^{-1}\right) &= f(a)f(b)f(a)^{-1}f(b)^{-1} \\ &= f(a)f(a)^{-1}f(b)f(b)^{-1} \\ &= e. \end{aligned}$$

Since every $x \in G^c$ is a finite product of commutators, by the fact that f is a homomorphism and by the above computation,

$$f(x) = e \quad \forall x \in G^c$$

Define

$$\bar{f}\colon G/G^c \to A$$

by $gG^c \mapsto f(g)$. First, we show that $\bar{f}$ is well-defined. Suppose $gG^c = hG^c$. Thus, by coset equality, $gh^{-1} \in G^c$. Then

$$\begin{aligned} f\left(gh^{-1}\right) = f(g)f(h)^{-1} &= e \\ \Rightarrow f(g) &= f(h) \end{aligned}$$

Therefore, $\bar{f}$ is well-defined. Now, let $gG^c, hG^c \in G/G^c$. Then

$$\begin{aligned}\bar{f}\left(\left(gG^c\right)\left(hG^c\right)\right) &= \bar{f}\left(ghG^c\right)\\ &= f(gh)\\ &= f(g)f(h)\\ &= \bar{f}\left(gG^c\right)\bar{f}\left(hG^c\right)\end{aligned}$$

Thus, $\bar{f}$ is a homomorphism. Hence, for all $g \in G$,

$$\left(\bar{f} \circ \pi\right)(g) = \bar{f}(\pi(g)) = \bar{f}\left(gG^c\right) = f(g)$$

Thus, $\bar{f} \circ \pi = f$. Therefore, every homomorphism of G into an abelian group factors through G/G^c.

□

Exercise 0.3. (Exercise 4)
Let H, K be subgroups of a finite group G with $K \subset N_H$. Show that

$$|HK| = \frac{|H| \cdot |K|}{|H \cap K|}.$$

Proof. Since $K \subseteq N_H$, we have $HK \leqslant G$. Define

$$f \colon H \times K \to HK$$

by $(h, k) = hk$. Let $x \in HK$. Then $x = hk$ for some $h \in H$ and $k \in K$. Hence, $f((h, k)) = hk = x$. Thus, f is surjective. Let $(h, k) \in \ker f$. Then

$$\begin{aligned}hk &= e\\ k &= h^{-1} \in K\end{aligned}$$

Thus, since $K \leqslant G$ and $h^{-1} \in K$, we must also have $h \in K$. Therefore

$$\ker f = \left\{\left(h, h^{-1}\right) \in H \times K \colon h \in H \cap K\right\} \triangleleft H \times K$$

We can define a bijection between $H \cap K$ and $\ker f$ by $h \mapsto \left(h, h^{-1}\right)$. Thus, $|\ker f| = |H \cap K|$. By the First Isomorphism Theorem,

$$\begin{aligned}H \times K/\ker f &\cong f\left(H \times K\right)\\ H \times K/H \cap K &\cong HK\end{aligned}$$

Thus,

$$\begin{aligned}|H \times K/H \cap K| &= |HK|\\ \frac{|H \times K|}{|H \cap K|} &= |HK|\\ \frac{|H| \cdot |K|}{|H \cap K|} &= |HK|\end{aligned}$$

□

Exercise 0.4. (Exercise 7)
Let G be a group such that $Aut(G)$ is cyclic. Prove that G is abelian.

Proof. Suppose $\mathrm{Aut}(G)$ is cyclic. Hence, $\mathrm{Inn}(G)$ is cyclic since $\mathrm{Inn}(G) \subseteq \mathrm{Aut}(G)$. Consider the map $\psi : G \to \mathrm{Inn}(G)$ given by $g \mapsto \varphi_g$ for all $g \in G$ where $\varphi_g : G \to G$ by $x \mapsto gxg^{-1}$ for all $x \in G$. Notice that $\mathrm{Ker}\,\psi = Z(G)$, since if $\varphi_g(x) = gxg^{-1} = x$, then $gx = xg \Rightarrow g \in Z(G)$. Thus, by the First Isomorphism Theorem, $G/\mathrm{Ker}\,\psi \cong \mathrm{Inn}(G) \Rightarrow G/Z(G) \cong \mathrm{Inn}(G)$. Since we have this isomorphism, $G/Z(G)$ and $\mathrm{Inn}(G)$ have the same group structure, i.e. $G/Z(G)$ is cyclic and thus $G/Z(G)$ is abelian which implies from **Exercise 1** that G is abelian.

□

Exercise 0.5. (Exercise 9)

a) Let G be a group and H a subgroup of finite index. Show that there exists a normal subgroup N of G contained in H and also of finite index. [Hint: If $(G : H) = n$, find a homomorphism of G into S_n whose kernel is contained in H.]

b) Let G be a group and let H_1, H_2 be subgroups of finite index. Prove that $H_1 \cap H_2$ has finite index.

Proof. a) Let G be a group and $H \leqslant G$ such that $(G\colon H) = n$. Consider the left cosets $G/H = \{g_1H, \ldots, g_nH\}$. Define an action on these cosets by

$$\varphi_g\colon G/H \to G/H$$

by $\varphi_g(xH) = gxH$. This is well-defined for if $xH = yH$, then $x - y \in H$ and hence $gx - gy \in H \Rightarrow gxH = gyH \Rightarrow \varphi_g(xH) = \varphi_g(yH)$. The homomorphism φ_g can be shown to be bijective as well . Thus, $\varphi \in \mathrm{Sym}(G/H) \cong S_n$. Define

$$\psi\colon G \to S_n \cong \mathrm{Sym}(G/H)$$

by $\psi(g) = \varphi_g$. This is a homomorphism since for $g_1, g_2 \in G$,

$$\begin{aligned}
\psi(g_1g_2) &= \varphi_{g_1g_2}(xH) \quad \text{for any } xH \in G/H \\
&= g_1g_2xH \\
&= g_1\varphi_{g_2}(xH) \\
&= \varphi_{g_1}(\varphi_{g_2}(xH)) \\
&= \varphi_{g_1} \circ \varphi_{g_2}
\end{aligned}$$

The kernel, $\ker\psi$, is defined to be $g \in G$ such that $\psi(g) = id$. Hence, if $\psi(g) = id$ (i.e. $g \in \ker\psi$), we get

$$\begin{aligned}
&\psi(g) = id \\
\Rightarrow &\varphi_g(xH) = xH \quad \forall xH \in G/H \\
\Rightarrow &gxH = xH \\
\Rightarrow &x^{-1}gxH = H \quad \forall x \in G \\
\Rightarrow &x^{-1}gx \in H \quad \forall x \in G \\
\Rightarrow &g \in xHx^{-1} \quad \forall x \in G \\
\Rightarrow &g \in \bigcap_{x\in G} xHx^{-1}
\end{aligned}$$

Thus, $\ker\psi = \bigcap_{x\in G} xHx^{-1} \subseteq H$. Furthermore, $N = \ker\psi \triangleleft G$. By the First Isomorphism Theorem, $G/\ker\psi \cong \psi(G) \leqslant S_n$. Hence, $(G\colon N) = |G/N| = |\psi(G)| \leq |S_n| = n! < \infty$.

b) Suppose $H_1, H_2 \leqslant G$ such that $(G\colon H_1) = n$, $(G\colon H_2) = m$. Consider the cosets of $H_1 \cap H_2$. Any left coset of $H_1 \cap H_2$ has the form $g(H_1 \cap H_2) = gH_1 \cap gH_2$ for $g \in G$. Hence, every left coset of $H_1 \cap H_2$ is the intersection of a left coset of H_1 with a left coset of H_2. From $(G\colon H_1) = n$, $(G\colon H_2) = m$, there are at most nm distinct intersections and thus at most nm left cosets of $H_1 \cap H_2$. Hence, $(G\colon H_1 \cap H_2) \leq nm < \infty$.

□

Exercise 0.6. (Exercise 10) Let G be a group and let H be a subgroup of finite index. Prove that there is only a finite number of right cosets of H, and that the number of right cosets is equal to the number of left cosets.

Proof. By assumption, $(G\colon H) = n < \infty$. The group G acts on G/H in two ways:

$$\lambda : G \to \mathrm{Perm}(G/H)$$

by $x \mapsto x(gH)$ for some $g \in G$ and

$$\rho : G \to \mathrm{Perm}(G/H)$$

by $x \mapsto (gH)x$ for some $g \in G$. Consider $eH = H \in G/H$. The isotropy group of eH in G is

$$G_{eH} = \{g \in G : \ g(eH) = eH = H\} = H.$$

The orbit of eH under G is

$$G \cdot eH = \{g(eH) : \ g \in G\} = G/H.$$

By the orbit-stabilizer theorem and Lagrange's Theorem,

$$|G/H| = |G \cdot eH| = \underbrace{(G\colon G_{eH})}_{\text{under } \lambda} = (G\colon H) = n < \infty.$$

Now, under ρ, $G_{eH} = \{g \in G : \ (eH)g = H\}$. If $Hg = H$, then $g = eg \in Hg = H$, so $g \in H$. Conversely, if $g \in G$, then clearly $Hg \subseteq H$ and $g^{-1} \in H$ gives $H \subseteq Hg$, so $H = Hg$. Hence, $G_{eH} = H$. The orbit of eH under ρ in G is

$$G \cdot (eH) = \{(eH)g : \ g \in G\} = Hg = H/G \quad \text{(notation in Lang for right cosets)}$$

Again by the orbit-stabilizer theorem and Lagrange's

$$|H/G| = |G \cdot eH| = \underbrace{(G\colon G_{eH})}_{\text{under } \rho} = (G\colon H) = n < \infty.$$

Therefore, $|H/G| = |G/H| = n < \infty$, i.e. there is a finite number of right cosets and the number of left cosets is the same as the number of right cosets.

□

Exercise 0.7. (Exercise 11) Let G be a group and A a normal abelian subgroup. Show that G/A operates on A by conjugation, and in this manner get a homomorphism of G/A into $\mathrm{Aut}(A)$.

Proof. Define a map $\varphi : G/A \to \mathrm{Aut}(A)$ by $\varphi(gA) = gAg^{-1}$, $g \in G$. (Well-defined) Suppose that $gA = hA$. Then $h = ga \in H$ for some $a \in A$. For any $x \in A$, we have $hxh^{-1} = (ga)x(ga)^{-1} = g(axa^{-1})g^{-1}$. Since A is abelian, $axa^{-1} = x$. Thus, $hxh^{-1} = gxg^{-1}$. Thus, the definition of φ does not depend on the choice of representative in G/A. ($\mathrm{Im}\,\varphi \subseteq Aut(A)$) For each $g \in G$, the map $a \mapsto gag^{-1}$ is a homomorphism $A \to A$ since conjugation preserves products. It is bijective with inverse $a \mapsto g^{-1}ag$. Hence, $\varphi(gA) \in Aut(A)$. (Homomorphism) For $gA, hA \in G/A$ and $x \in A$,

$$\begin{aligned}(\varphi(gA) \circ \varphi(hA))(x) &= g(hxh^{-1})g^{-1} \\ &= (gh)x(gh)^{-1} = \varphi(ghA)(x).\end{aligned}$$

Therefore, φ is a group homomorphism. Thus, G/A acts on A by conjugation, and we obtain a homomorphism

$$\varphi : G/A \to \mathrm{Aut}(A)$$

Its kernel is $\mathrm{Ker}\,\varphi = C_G(A)/A$, where $C_G(A)$ is the centralizer of A in G.

□

Semidirect Product

Exercise 0.8. (Exercise 12) Let G be a group and let H, N be subgroups with N normal. Let γ_x be conjugation by an element $x \in G$.

a) Show that $x \mapsto \gamma_x$ induces a homomorphism $f : H \to Aut(N)$.

b) If $H \cap N = \{e\}$, show that the map $H \times N \to HN$ given by $(x, y) \mapsto xy$ is a bijection, and that this map is an isomorphism if and only if f is trivial, i.e. $f(x) = id_N$ for all $x \in H$.

c) We define G to be the **semidirect product** of H and N if $G = HN$ and $H \cap N = \{e\}$. Now, conversely, let N, H be groups, and let $\psi : H \to Aut(N)$ be a given homomorphism. Construct a semidirect product as follows. Let G be the set of pairs (x, h) with $x \in N$, $h \in H$. Define the composition law

$$(x_1, h_1)(x_2, h_2) = (x_1\psi(h_1)x_2, h_1h_2)$$

Show that this is a group law, and yields a semidirect product of N and H, identifying N with the set of elements $(x, 1)$ and H with the set of elements $(1, h)$.

Proof. a) First, let $e_H \in H$ be the identity element in H. Then, $f(e_H) = \gamma_{e_H}$. Let $n \in N$. Thus, $\gamma_{e_H}(n) = e_H n e_H^{-1} = n$ (since N normal). Thus, $f(e_H) = \gamma_{e_H} = id_N$. It is also well-defined since if $h_1 = h_2$, then $f(h_1) = \gamma_{h_1}(n) = h_1 n h_1^{-1}$ and $f(h_2) = \gamma_{h_2}(n) = h_2 n h_2^{-1}$. Clearly, we then have $f(h_1) = f(h_2)$. Let $x, y \in H$. Then, $f(xy) = \gamma_{xy}$. Take $n \in N$, $\gamma_{xy}(n) = xyn(xy)^{-1} = xyny^{-1}x^{-1} = x(\gamma_y(n))x^{-1} = \gamma_x(\gamma_y(n)) = (\gamma_x \circ \gamma_y)(n) = f(x) \circ f(y)$ as desired.

b) (Well-defined) For any $h \in H$, $n \in N$, $hn \in HN = \{xy : x \in H, y \in N\}$. (Injective) Suppose $(x_1, y_1), (x_2, y_2) \in H \times N$ and suppose $\varphi((x_1, y_1)) = \varphi((x_2, y_2)) \Rightarrow x_1y_1 = x_2y_2 \Rightarrow x_2^{-1}x_1 = y_2y_1^{-1}$. Note $x_2^{-1}x_1 \in H$ and $y_2y_1^{-1} \in N$. Thus, $x_2^{-1}x_1 \in H \cap N$ and $y_2y_1^{-1} \in H \cap N$. But $H \cap N = \{e\}$ by assumption, so $x_2^{-1}x_1 = e \Rightarrow x_1 = x_2$ and $y_2y_1^{-1} = e \Rightarrow y_1 = y_2$. Hence, $(x_1, y_1) = (x_2, y_2)$. (Surjective) Let $z \in HN$. Then, $z = hn$ for some $h \in H$, $n \in N$. Hence, $\varphi((h, n)) = hn = z$ with (h, n) clearly an element of $H \times N$. Thus, $\varphi : H \times N \to HN$ is a bijection.

Suppose $\varphi((x_1, y_1)(x_2, y_2)) = \varphi((x_1, y_1))\varphi((x_2, y_2))$ for $(x_i, y_i) \in H \times N$. This is if and only if

$$\begin{aligned} & x_1x_2y_1y_2 = x_1y_1x_2y_2 \\ \iff & x_2y_1 = y_1x_2 \quad \text{(left/right cancellation)} \\ \iff & y_1 = x_2y_1x_2^{-1} \end{aligned}$$

Since x_1, x_2, y_1, y_2 are arbitrary, this must hold for all $x_2 \in H$ and $y_1 \in N$. Hence, $\gamma_x |_N = \mathrm{id}_N \Rightarrow f(x) = \mathrm{id}_N$ for all $x \in H$.

c) Let $\psi : H \to Aut(N)$ be given by $\psi(h) = \varphi_h$ with $\varphi_h(n) = n' \in N$. Thus, we define the composition of $(x_1, h_1), (x_2, h_2) \in N \times H$ by

$$\begin{aligned} (x_1, h_1)(x_2, h_2) &= (x_1\psi(h_1)(x_2), h_1h_2) \\ &= (x_1\varphi_{h_1}(x_2), h_1h_2) \end{aligned}$$

(Identity) Let $e = (e_N, e_H) \in G = N \times H$. Then, $(x_1, h_1)(e_N, e_H) = (x_1\psi(h_1)(e_N), h \cdot e_N) = (x_1\varphi_{h_1}(e_N), h_1) = (x_1 \cdot e_N, h_1) = (x_1, h_1)$ (since $\varphi \in Aut(N)$ fixes the identity). Likewise, $(e_N, e_H)(x_1, h_1) = (x_1, h_1)$ and thus $e \in G$ which implies that $G \neq \varnothing$. (Closed) Observe that

$$\begin{aligned} (x_1, h_1)(x_2, h_2) &= (x_1\psi(h_1)(x_2), h_1h_2) \\ &= (x_1\varphi_{h_1}(x_2), h_1h_2) \end{aligned}$$

and since $\varphi_{h_1} \in Aut(N)$, $\varphi_{h_1}(x_2) \in N$. Hence, $x_1\varphi_{h_1}(x_2) \in N$ and $h_1h_2 \in H \Rightarrow (x_1\varphi_{h_1}(x_2), h_1h_2) \in G = N \times H$. (Associativity) Let $(x_1, h_1), (x_2, h_2), (x_3, h_3) \in G$. Then,

$$\begin{aligned}
(x_1, h_1)\left[(x_2, h_2)(x_3, h_3)\right] &= (x_1, h_1)(x_2\psi(h_2)(x_3), h_2h_3) \\
&= (x_1, h_1)(x_2\varphi_{h_2}(x_3), h_2h_3) = (x_1\psi(h_1)\left[x_2\varphi_{h_2}(x_3)\right], h_1h_2h_3) \\
&= (x_1\varphi_{h_1}(x_2\varphi_{h_2}(x_3)), h_1h_2h_3) \\
&= (x_1\varphi_{h_1}(x_2)\varphi_{h_1}(\varphi_{h_2}(x_3)), h_1h_2h_3)
\end{aligned}$$

Now,

$$\begin{aligned}
\left[(x_1, h_1)(x_2, h_2)\right](x_3, h_3) &= (x_1\psi(h_1)(x_2), h_1h_2)(x_3, h_3) \\
&= (x_1\varphi_{h_1}(x_2), h_1h_2)(x_3, h_3) \\
&= (x_1\varphi_{h_1}(x_2)\psi(h_1h_2)(x_3), h_1h_2h_3) \\
&= (x_1\varphi_{h_1}(x_2)(\psi(h_1)\psi(h_2))(x_3), h_1h_2h_3) \\
&= (x_1\varphi_{h_1}(x_2)\varphi_{h_1}(\varphi_{h_2}(x_3)), h_1h_2h_3)
\end{aligned}$$

Thus, $(x_1, h_1)\left[(x_2, h_2)(x_3, h_3)\right] = \left[(x_1, h_1)(x_2, h_2)\right](x_3, h_3)$. (Inverses) Suppose that $(x_1, h_1)(x_2, h_2) = (e_N, e_H)$. Thus,

$$\begin{aligned}
(x_1, h_1)(x_2, h_2) &= (e_N, e_H) \\
\Rightarrow (x_1\psi(h_1)(x_2), h_1h_2) &= (e_N, e_H) \\
\Rightarrow (x_1\varphi_{h_1}(x_2), h_1h_2) &= (e_N, e_H) \\
\Rightarrow x_1\varphi_{h_1}(x_2) = e_N \quad &\text{and} \quad h_1h_2 = e_H \\
\Rightarrow \varphi_{h_1}(x_2) = x_1^{-1} \qquad &\Rightarrow h_2 = h_1^{-1} \in H \\
\because \varphi_{h_1} \in Aut(N), &\varphi_{h_1^{-1}} \in Aut(N) \\
\Rightarrow \varphi_{h_1^{-1}}(\varphi_{h_1}(x_2)) = \varphi_{h_1^{-1}}(x_1^{-1}) &\Rightarrow x_2 = \varphi_{h_1^{-1}}(x_1^{-1}) \in N.
\end{aligned}$$

Therefore, $(x_1, h_1)^{-1} = (\varphi_{h_1^{-1}}(x_1^{-1}, h_1^{-1}) = (\psi(h_1^{-1})(x_1^{-1}), h_1^{-1}) \in G$. ($G = HN$) Let $(x, h) \in G = N \times H$. Then,

$$\begin{aligned}
(x, h) &= (x\varphi_{e_H}(e_N), e_H \cdot h) \\
&= (x\psi(e_H)(e_N), e_H \cdot h) \\
&= \underbrace{(x, e_H)}_{\in N}\underbrace{(e_N, h)}_{\in H}
\end{aligned}$$

Thus, $G = NH$. ($N \cap H = \{e\}$) Note that $(x, e_H) \in H \iff x \in e_N$ and $(e_N, h) \in N \iff h = e_H$. Thus, $N \cap H = \{(e_N, e_H)\} = \{e\}$. Therefore,

$$G = N \rtimes_\psi H.$$

□

Exercise 0.9. (Exercise 13)

a) Let H, N be normal subgroups of a finite group G. Assume that the orders of H, N are relatively prime. Prove that $xy = yx$ for all $x \in H$ and $y \in N$, and that $H \times N \cong HN$.

b) Let $H_1, \ldots, H_r$ be normal subgroups of G such that the order of H_i is relatively prime to the order of H_j for $i \neq j$. Prove that

$$H_1 \times \cdots \times H_r \cong H_1 \cdots H_r.$$

Example. If the Sylow subgroups of a finite group are normal, then G is the direct product of its Sylow subgroups.

Proof. a) Assume $H, N \triangleleft G$ with $(|H|, |N|) = 1$. Consider the commutator,

$$[x, y] \coloneqq xyx^{-1}y^{-1} \quad \forall x \in H, y \in N$$

Thus, $xyx^{-1}y^{-1} \in H$ since $yx^{-1}y^{-1} \in H$ by the fact that $H \triangleleft G$ and therefore $x(yx^{-1}y^{-1}) \in H$. Similarly $xyx^{-1}y^{-1} \in N$ since $xyx^{-1} \in N$ by the fact that $N \triangleleft G$. Thus, $[x, y] = xyx^{-1}y^{-1} \in H \cap N$. By Lagrange's Theorem,

$$|H \cap N| \Big| |H| \quad \text{and} \quad |H \cap N| \Big| |N|$$

and thus,

$$|H \cap N| \Big| \underbrace{(|H|, |N|)}_{=1}$$

Hence, we must have $|H \cap N| = 1 \Rightarrow H \cap N = \{e\}$. Then since $[x, y] \in H \cap N$, it follows that $xyx^{-1}y^{-1} = e \Rightarrow xy = yx$. Finally, by *Exercise 12*, $H \times N \cong HN$.

b) Assume $H_1, \ldots, H_r \triangleleft G$ with $(|H_i|, |H_j|) = 1$ if $i \neq j$. From *part (a)*, for any H_i, H_j with $i \neq j$, $H_i \cap H_j = \{e\}$ and $xy = yx$ for all $x \in H_i$, $y \in H_j$. Hence, $H_i \times H_j \cong H_iH_j$. We proceed by induction. Suppose that $H_1 \times \cdots \times H_{r-1} \cong H_1 \cdots H_{r-1}$. Then, $(|H_r|, |H_1 \cdots H_{r-1}|) = 1$ and $H_r \triangleleft G$. Thus, by the same argument as above, $H_1 \times \cdots \times H_r \cong H_1 \cdots H_r$.

□

Exercise 0.10. (Exercise 14)
Let G be a finite group and let N be a normal subgroup such that N and G/N have relatively prime orders.

a) Let H be a subgroup of G having the same order as G/N. Prove that $G = HN$.

b) Let g be an automorphism of G. Prove that $g(N) = N$.

Proof. Let G be a finite group of order n and let $N \triangleleft G$ such that $(|N|, |G/N|) = 1$.

a) Assume that $H \leqslant G$ such that $|H| = |G/N| = (G\colon N)$. Then $(|H|, |N|) = 1$. We have

$$\begin{aligned} |HN| &= \frac{|H| \cdot |N|}{|H \cap N|} \\ &= \frac{|G/N| \cdot |N|}{|H \cap N|} \\ &= \frac{\frac{|G|}{|N|} \cdot |N|}{|H \cap N|} \\ &= \frac{|G|}{|H \cap N|} \end{aligned}$$

Then, if $|HN| = |G|$, we would require that $|H \cap N| = 1 \Rightarrow H \cap N = \{e\}$. Hence, since $HN \subseteq G$, we have that $G = HN$ if and only if $H \cap N = \{e\}$. Thus, we show the converse of this biconditional.

By Lagrange's Theorem, $|H \cap N| \Big| |H|$ and $|H \cap N| \Big| |N|$, and thus $|H \cap N| \Big| \underbrace{(|H|, |N|)}_{=1}$. Hence, $|H \cap N| = 1$ and therefore, $H \cap N = \{e\}$. It follows from the above biconditional that $G = HN$.

b) Let $g \in \mathrm{Aut}(G)$. Then $g(N) \leqslant G$ with $|g(N)| = |N|$. By our assumption, $(|N|, |G/N|) = 1$, we have $(|g(N)|, |G/g(N)|) = 1$. Let $H \leqslant G$ such that $|H| = |G/N|$ as in *part (a)* which implies that $G = HN$ and $H \cap N = \{e\}$. Then $|g(H)| = |H| = |G/N|$ and applying *part (a)* to the normal subgroup $g(N)$ gives that $G = g(H) \cdot g(N)$ and $g(H) \cap g(N) = \{e\}$. Hence, since $N, g(N) \triangleleft G$ are the same order, admitting a complement of order $|G/N|$. The order of $Ng(N)$ is

$$|Ng(N)| = \frac{|N| \cdot |g(N)|}{|N \cap g(N)|} = \frac{|N|^2}{|N \cap g(N)|} \leq |G| = |N| \cdot |G/N|$$

Hence, we have that $|N \cap g(N)| \geq \frac{|N|}{|G/N|}$. Since $(|N|, |G/N|) = 1$, this implies that $|N \cap g(N)| = |N|$, so $N \subseteq g(N)$. By symmetry, using g^{-1} (since g is an automorphism), we have $g(N) \subseteq N$. Therefore, $g(N) = N$.

□

Exercise 0.11. (Exercise 20) Let P be a p-group. Let A be a normal subgroup of order p. Prove that A is contained in the center of P.

Proof. Assume $|A| = p$. Thus, $A \cong C_p$, the cyclic group of order p (since prime order implies cyclic). Thus, $\mathrm{Aut}(A)$ is cyclic of order $p - 1$. Define

$$\varphi : P \to \mathrm{Aut}(A)$$

by $g \mapsto c_g$, conjugation by g in A. Since $A \triangleleft P$, and by previous exercises, c_g is a bijective homomorphism of A into itself, hence $c_g \in \mathrm{Aut}(A)$. Our map φ is a homomorphism since for $g, h \in P$ we have

$$\varphi(gh) = c_{gh} = c_g \circ c_h = \varphi(g)\varphi(h).$$

By definition, $\varphi(P) \leq \mathrm{Aut}(A)$, thus $|\varphi(P)| \Big| |\mathrm{Aut}(A)| \Rightarrow |\varphi(P)| \Big| p - 1$. But $\varphi(P)$ is also a homomorphic image of the p-group P, hence $|\varphi(P)| = p^k$ for some k. This is due to the First Isomorphism Theorem which states

$$P/\ker\varphi \cong \varphi(P).$$

Since $|P| = p^n$ for some n by assumption, and $\ker\varphi \leq P$, by Lagrange's Theorem, $|\ker\varphi| = p^k$ for $k \leq n$. Then, by counting sizes via the isomorphism,

$$|\varphi(P)| = |P/\ker\varphi| = \frac{|P|}{|\ker\varphi|} = \frac{p^n}{p^k} = p^{n-k} \qquad \text{(a power of } p\text{)}$$

Whence, the only positive integer that is both a power of p and a divisor $p - 1$ is 1. Thus, $|\varphi(P)| = p^0 = 1$. Thus means $\varphi(P)$ is trivial, i.e. $\varphi(g) = \mathrm{id}_A$ for all $g \in P$. Then, $(\varphi(g))(a) = c_g(a) = gag^{-1} = a$ (since $\varphi(g)(a) = a$ by $\varphi(g) = \mathrm{id}_A$). Hence, $ga = ag$. Therefore every element of A commutes with every element of P. Therefore,

$$A \subseteq Z(P).$$

□

Exercise 0.12. (Exercise 21) Let G be a finite group and H a subgroup. Let P_H be a p-Sylow subgroup of H. Prove that there exists a p-Sylow subgroup P of G such that $P_H = P \cap H$.

Proof. Let G be a finite group, $H \leq G$, and let P_H be a p-Sylow subgroup of H. By Theorem 6.4, there exists a p-Sylow subgroup $P \leq G$ such that every p-subgroup of G is contained in P. Hence, since $H \leq G$, P_H is a p-subgroup of G (p-Sylow subgroup of H, by assumption) and thus

$$P_H \subseteq P$$

Furthermore, $P_H \subseteq H \Rightarrow P_H \subseteq P \cap H$. Observe that $P \cap H \leq H$ and thus since P is a p-subgroup (also Sylow), $P \cap H$ is a p-subgroup of H and therefore $P \cap H \subseteq P_H$. Hence,

$$P \cap H = P_H.$$

□

Exercise 0.13. (Exercise 22)
Let H be a normal subgroup of a finite group G and assume that $\#(H) = p$. Prove that H is contained in every p-Sylow subgroup of G.

Proof. Assume that $H \triangleleft G$ and suppose $|H| = p$. This follows nicely from all parts of Theorem 6.4. Thus, H is a p-subgroup and for all $x \in G$,

$$xHx^{-1} = H \quad \text{(normality)}$$

Then by definition, $N(H) = G$. By Theorem 6.4(i), since H is a p-subgroup of G, H is contained in some p-Sylow subgroup, say K. Then, by Theorem 6.4(ii), all p-Sylow subgroups are conjugate and thus $K' = gKg^{-1}$ is a p-Sylow subgroup for some $g \in G$. Conjugating the inclusion $H \subseteq K$, we get

$$H = xHx^{-1} \subseteq xKx^{-1} = K'$$

Hence, $H \subseteq K'$ with K' a different, unless K normal, p-Sylow subgroup. Thus, we can continue this process until we get all $pk + 1$ p-Sylow subgroups ($k \in \mathbb{Z}$, by Theorem 6.4(iii) which states the number of p-Sylow subgroups of G is $\equiv 1 \mod p$). Thus, it follows that H is contained in every p-Sylow subgroup.

□

Exercise 0.14. (Exercise 23) Let P, P' be p-Sylow subgroups of a finite group G.

a) If $P' \subseteq N(P)$ (normalizer of P), then $P' = P$.

b) If $N(P') = N(P)$, then $P' = P$.

c) We have $N(N(P)) = N(P)$.

Proof. a) Suppose $P' \subseteq N(P)$. Then for all $x \in P'$, we have

$$xPx^{-1} = P$$

and thus $P \triangleleft P'$. Since P, P' are p-groups and P is a normal subgroup of P', the quotient P'/P is also a p-group. By definition, $|P| = |P'| = p^n$ with p^n the highest power of p dividing $|G|$. Hence, $|P'/P| = \frac{|P'|}{|P|} = 1$. Thus, $P = P'$.

b) We know $P' \subseteq N(P')$ and $P \subseteq N(P)$. By assumption $N(P') = N(P)$, so

$$P' \subseteq N(P') = N(P)$$

Thus, by part (a), $P' = P$.

c) ($N(P) \subseteq N(N(P))$) This inclusion follows trivially from the fact that for any subgroup $H \leqslant G$, $H \subseteq N(H)$, i.e. every subgroup is a subset of its normalizer. Thus, if you let $K = N(H)$, then we have $K \subseteq N(K)$.

($N(N(P)) \subseteq N(P)$) Let $x \in N(N(P))$. Then

$$xN(P)x^{-1} = N(P)$$

and in particular,

$$xPx^{-1} \subseteq N(P)$$

Thus, xPx^{-1} is a p-Sylow subgroup of $N(P)$. Since $P \subseteq N(P)$, P is also a p-Sylow subgroup of $N(P)$. By Theorem 6.4, there exists $y \in N(P)$ such that

$$y(xPx^{-1})y^{-1} = P$$

Then $yx \in N(P)$ and since $N(P)$ is a subgroup, we have that $x \in N(P)$ as was to be shown. Therefore, $N(N(P)) = N(P)$.

□

Exercise 0.15. (Exercise 24)
Let p be a prime number. Show that a group of order p^2 is abelian, and that there are only two such groups up to isomorphism.

Proof. Suppose $|G| = p^2$ for a prime p. Thus, G is a p-group and hence $Z(G)$ is not trivial. Recall that the class equation is defined as

$$|G| = |Z(G)| + \sum_{x \in C} (G\colon G_x)$$

where C is a set of representatives for the distinct conjugacy classes and G_x is the isotropy group of x in G defined as $G_x = \{x \in G : \ xy = y\}$ with $y \in G$ (here G is acting on itself via conjugation). Each term in $\sum_{x \in C} (G\colon G_x) > 1$ and divides $|G| = p^2$. Hence, $\sum_{x \in C} (G\colon G_x) \equiv 0 \pmod p$. Thus, $|G| = |Z(G)| + kp \ (k \in \mathbb{Z}^+) \left[kp = \sum_{x \in C} (G\colon G_x)\right]$. Therefore, $|Z(G)| = |G| - kp \Rightarrow |Z(G)| \equiv 0 \pmod p$. Thus, $p \,\big|\, |Z(G)| \Rightarrow |Z(G)| = p$ or p^2.
But by **Exercise 1**, Cases 1 and 2 cover this, thus, it follows that G is abelian.
If G has an element g of order p^2, then the order of $\langle g \rangle$ is p^2 and thus $G = \langle g \rangle$. Therefore, in this case, $G \cong \mathbb{Z}/p^2\mathbb{Z}$.
If no element (not including the identity) of G has order p^2, then it must have order p. Choose $a \in G$ such that $|\langle a \rangle| = p$ and thus $(G\colon \langle a \rangle) = p$. Choose $b \notin \langle a \rangle$. Hence, we also have $|b| = p$. Hence, we have $\langle a \rangle \cap \langle b \rangle = \{e\}$ with the product $\langle a \rangle \times \langle b \rangle$ having order p^2. Therefore, $G = \langle a \rangle \times \langle b \rangle \cong \mathbb{Z}/p\mathbb{Z} \times \mathbb{Z}/p\mathbb{Z}$. Then these two possibilities of G can not be isomorphic since one is cyclic and one is not, and cyclicity is invariant under isomorphisms.

□

Exercise 0.16. (Exercise 25)
Let G be a group of order p^3, where p is prime, and G is not abelian. Let Z be its center. Let C be a cyclic group of order p.

a) Show that $Z \cong C$ and $G/Z \cong C \times C$.

b) Every subgroup G of order p^2 contains Z and is normal.

c) Suppose $x^p = 1$ for all $x \in G$. Show that G contains a normal subgroup $H \cong C \times C$.

Proof. a) Assume that G is not an abelian group with order p^3. Since $|G| = p^3$, G is a p-group. By Theorem 6.5, since $p^3 > 1$, G has a nontrivial center. By the assumption that G is not abelian, $G \neq Z$. Thus, $|Z| = p$ or p^2. If $|Z| = p^2$, then $|G/Z| = p$ and thus G/Z is cyclic. By **Exercise 1**, it follows that G is abelian, a contradiction! Therefore $|Z| = p$ and thus Z is cyclic of order p, i.e. $Z \cong C$. Then $|G/Z| = p^2$. By **Exercise 24**, there are two abelian groups, up to isomorphism, of order p^2, a cyclic group and a non-cyclic group. If G/Z is cyclic, then by **Exercise 1**, G is abelian, a contradiction! Hence, G/Z is not cyclic which implies that $G/Z \cong C \times C$.

b) Let $H \leqslant G$ such that $|H| = p^2$. Thus, by **Exercise 24**, H is abelian. It follows that $H \cong C_{p^2} \cong \mathbb{Z}/p^2\mathbb{Z}$ or $H \cong C \times C$. From part (a), $|Z| = p$. Hence, $|G/Z| = p^2$. In part (a), we showed that

$$G/Z \cong C \times C.$$

Consider

$$\pi\colon G \to G/Z$$

given by $g \mapsto gZ$. Thus

$$\ker \pi = \{x \in G\colon xZ = Z\} = G \cap Z = Z$$

Let $\pi|_H$ be the restriction to H. Then

$$\ker \pi|_H = H \cap Z \subseteq Z.$$

Since $\ker \pi|_H \triangleleft H$, Lagrange's and the containment, $|H \cap Z| \big| |Z| \Rightarrow |H \cap Z| = 1$ or p. By the First Isomorphism Theorem,

$$\begin{aligned} &H/\ker \pi|_H \cong \pi(H) \\ \Rightarrow &H/H \cap Z \cong \pi(H) \end{aligned}$$

Therefore,

$$|H/H \cap Z| = |\pi(H)| \Rightarrow \frac{|H|}{|H \cap Z|} = |\pi(H)|.$$

If $|H \cap Z| = 1$, then $|\pi(H)| = |H| = p^2 = |G/Z|$. This would mean that H maps surjectively onto G/Z but $H \leqslant G$. Thus, we must have $|H \cap Z| = p$. Hence since $Z \cong C$, we have $H \cap Z = Z$ and therefore $Z \subseteq H$. Then $|\pi(H)| = p$. Thus, $\pi(H) \leqslant G/Z \cong C \times C$. Since G/Z is abelian , $\pi(H) \triangleleft G/Z$. The pre-image of a normal subgroup is normal, hence $\pi^{-1}(\pi(H)) \triangleleft G$. The pre-image is defined as

$$\begin{aligned} \pi^{-1}(\pi(G)) &= \{x \in G \colon xZ \in \pi(H)\} \\ &= \{x \in G \colon xZ = yZ \quad \text{for some } h \in H\} \end{aligned}$$

Let $g \in \pi^{-1}(\pi(H))$. Thus, $\pi(g) \in \pi(H)$ which implies that $\pi(g) = \pi(h)$ for some $h \in H$. Then $gZ = hZ$. Hence, there exists some $x \in Z$ such that $g = hx$. Since $Z \subseteq H$, we have $g = hx \in H$. Hence, $\pi^{-1}(\pi(H)) \subseteq H$. Now let $h \in H$. Then $\pi(h) = hZ \in \pi(H)$. Thus, $H \subseteq \pi^{-1}(\pi(H))$. Therefore, $\pi^{-1}(\pi(H)) = H$. Then from

$$\pi^{-1}(\pi(H)) \triangleleft G,$$

we have $H \triangleleft G$. Therefore, $Z \subseteq H$ and H is normal in G.

c) Since $G/Z \cong C \times C$ from part (a), G/Z contains a cyclic subgroup of order p, say

$$\langle aZ \rangle \leqslant G/Z.$$

Let

$$\begin{aligned} H = \pi^{-1}\left(\langle aZ \rangle\right) &= \{x \in G \colon \pi(x) \in \langle aZ \rangle\} \\ &= \{x \in G \colon xZ = a^k Z,\ k \in \mathbb{Z}\} \\ &= \{x \in G \colon x = a^k z,\ k \in \mathbb{Z},\ z \in Z\}. \end{aligned}$$

Thus, $|H| = |\langle aZ \rangle| \cdot |Z| = p^2$. By part (b), $H \triangleleft G$ and $Z \subseteq H$. By assumption, $x^p = 1$ for all $x \in G$, so all elements of H have orders dividing p. Therefore, since $|H| = p^2$ and no element of H has order p^2 ($p^2 \nmid p$), H cannot be cyclic. Thus, by **Exercise 24**, $H \cong C \times C$.

□

Exercise 0.17. (Exercise 26)

a) Let G be a group of order pq, where p, q are primes and $p < q$. Assume that $q \not\equiv 1 \mod p$. Prove that G is cyclic.

b) Show that every group of order 15 is cyclic.

Proof. a) Let $|G| = pq$ with $p < q$ distinct primes and assume that $q \not\equiv 1 \mod p$. Thus, there exists p, q-Sylow subgroups H_p, H_q respectively with $|H_p| = p$ and $|H_q| = q$. By Lemma 6.7, since $(G\colon H_q) = p < q$ is the smallest prime dividing the order of G, we have $H_q \triangleleft G$. Furthermore, since H_p, H_q are both prime order, they are cyclic (Proposition 4.1) and thus there exists $x \in H_p$ and $y \in H_q$ such that $H_p = \langle x \rangle$ and $H_= \langle y \rangle$. Define

$$\varphi\colon H_p \to \operatorname{Aut}(H_q)$$

by $x \mapsto \varphi_x$ with $\varphi_x\colon H_q \to H_q$ being conjugation by x, an automorphism. Since H_q is cyclic, any automorphism must send a generator to another generator, i.e. $y \mapsto y^k$. Thus, $k \in (\mathbb{Z}/q\mathbb{Z})^\times$. Then

$$\operatorname{Aut}(H_q) \cong (\mathbb{Z}/q\mathbb{Z})^\times$$

which implies that $|\operatorname{Aut}(H_q)| = q - 1$. Then

$$\begin{aligned} \ker\varphi &= \{x \in H_p\colon \varphi_x(y) = y,\ \forall\, y \in H_q\} \\ &= \{x \in H_p\colon xy = yx,\ \forall\, y \in H_q\} \triangleleft H_p \end{aligned}$$

By Lagrange's Theorem, $|\ker\varphi| \Big| |H_p|$. Hence $|\ker\varphi| = 1$ or p. If $|\ker\varphi| = 1$, by the First Isomorphism Theorem, we have

$$H_p/\ker\varphi \cong \varphi(H_p) \leqslant \operatorname{Aut}(H_q)$$

and thus $|H_p/\ker\varphi| = |H_p| = p = |\varphi(H_p)|$. Then, by Lagrange's Theorem, $|\varphi(H_p)| \Big| |\operatorname{Aut}(H_q)|$, but by assumption, $q \not\equiv 1 \mod p$. Hence we can not have $|\ker\varphi| = 1$. Thus, $|\ker\varphi| = p$ and therefore by the First Isomorphism Theorem, $|H_p/\ker\varphi| = 1 = |\varphi(H_p)|$. Thus, $\varphi(H_p) = \{\text{id.}\}$. Since φ is trivial, every element of H_p commutes with every element of H_q. Thus, H_pH_q is abelian. Let $z \in H_p \cap H_q$. Then $z \in H_p$ and $z \in H_q$. Hence, $z = x^k$ and $z = y^l$ for some $k \in (\mathbb{Z}/p\mathbb{Z})^\times$ and $l \in (\mathbb{Z}/q\mathbb{Z})^\times$. Then

$$\begin{aligned} x^k &= y^l \\ (x^k)^p &= (y^l)^p \\ e &= y^{lp} \end{aligned}$$

Thus, $q \mid lp$. By Euclid's Lemma, since q is a prime and $q \nmid p$, we have $q \mid l$ which implies that $l \equiv 0 \mod q$ and $y^l = e$. Then $x^k = y^l = e \Rightarrow k \equiv 0 \mod p$ and therefore $z = e$. Hence $H_p \cap H_q = \{e\}$. Thus,

$$|H_pH_q| = \frac{|H_p| \cdot |H_q|}{|H_p \cap H_q|} = \frac{pq}{1} = pq = |G|$$

Then $G = H_pH_q$. Since H_p, H_q are cyclic with $(p, q) = 1$, H_p, H_q abelian, and $H_p \cap H_q = \{e\}$, we have $H_pH_q = \langle xy \rangle$ with $|xy| = \operatorname{lcm}(|x|, |y|) = \operatorname{lcm}(p, q) = pq = |G|$. Therefore, $\langle xy \rangle$ has order pq and thus $G = H_pH_q = \langle xy \rangle$ is cyclic.

b) Let $|G| = 15 = 3 \cdot 5$. Since $5 \not\equiv 1 \mod 3$, by part (a), G is cyclic.

□

Exercise 0.18. (Exercise 28) Let p, q be distinct primes. Prove that a group of order p^2q is solvable, and that one of its Sylow subgroups is normal.

Proof. Let n_p and n_q denote the number of p-Sylow and q-Sylow subgroups G, respectively. By Sylow's Theorems,

$$n_p \Big| q, \quad n_p \equiv 1 \pmod p, \quad n_q \Big| p^2, \quad n_q \equiv 1 \pmod q$$

Hence,

$$n_p \in \{1, q\}, \quad n_q \in \{1, p, p^2\}$$

<u>Case 1: $(p < q)$</u> If $n_q = p$, then $q \Big| p - 1$ which is impossible since $q > p - 1$. If $n_q = p^2$, then $q \Big| (p-1)(p+1)$ which is impossible since $q > p + 1$ (as well). Thus, $n_q = 1$, so the q-Sylow subgroup $Q \triangleleft G$ is normal. <u>Case 2: $(p > q)$</u> If $n_p = q$, then $p \Big| q - 1$ which is impossible since $p > q - 1$. Thus, $n_p = 1$, so the p-Sylow subgroups $P \triangleleft G$ is normal. In both cases, G has a normal Sylow subgroup N (either $|N| = q$ or $|N| = p^2$). Now consider the quotient G/N.

- $|N|$ is a prime power and therefore is solvable.
- $|G/N|$ is either p^2 or pq and in either case, both are known to be solvable.

Since both N and G/N are solvable, it follows that G is solvable. [The fact that $n_p, n_q = 1 \Rightarrow P, Q$ are normal is since all Sylow groups are conjugate and if there is only one Sylow subgroup, then it is fixed by conjugation by G, hence normal.]

□

Exercise 0.19. (Exercise 32) Let S_n be the permutation group on n elements. Determine the p-Sylow subgroups of S_3, S_4, S_5 for $p = 2$ and $p = 3$.

Proof. We utilize Thm 6.2 throughout the following derivations to justify existence. We will find the Sylow subgroups up to isomorphism. <u>(S_3; order $3! = 6 = 2 \cdot 3$)</u> Let $\mathbf{p = 2}$. The highest power of 2 dividing 6 is 2. Hence, let $P_2 \leqslant S_3$ be a 2-Sylow subgroups and hence $|P_2| = 2$. Thus, $P_2 \cong C_2$ (P_2 is the subgroup generated by any of the three transpositions of S_3). Let $\mathbf{p = 3}$. The highest power of 3 dividing 6 is 3. Hence, let $P_3 \leqslant S_3$ be a 3-Sylow subgroup, $|P_3| = 3$. Observe $(S_3 \colon P_3) = 2$, thus $P_3 \triangleleft S_3$ and the only nontrivial normal subgroup of S_3 is A_3. Hence, $P_3 = A_3 = \langle (123) \rangle \cong C_3$.
<u>S_4; order $4! = 24 = 2^3 \cdot 3$</u> Let $\mathbf{p = 2}$. A 2-Sylow subgroup P_2 has order $2^3 = 8$. Note that all elements of a 2-Sylow subgroup have an order of a power of 2. For example:

- Transpositions have order 2.
- The product of disjoint transpositions have order 2.
- 4-cycles have order 4.

Thus, there are no 3-cycles. We will construct a 2-Sylow subgroup. Start with disjoint the disjoint transpositions $(12), (34)$. These cycles generate the Klein 4-group,

$$V_4 = \{(1), (12), (34), (12)(34)\}$$

Thus far, we have order 4. Now consider the permutation $(13)(24)$. Conjugation by this element yields,

$$(13)(24)(12)(13)(24) = (43) = (34)$$

Hence, $(13)(24)$ swaps the two generators of V_4. Now the subgroup

$$P_2 = \langle (12), (34), (13)(24) \rangle$$

contains V_4 as a normal subgroup and an element acting by symmetry. Furthermore, $|P_2| = 8$. A group of order 8 containing a normal Klein 4-subgroup, plus an element that swaps its generators, is exactly the Dihedral group, D_4. Hence,

$$P_2 \cong D_4$$

Thus, all 2-Sylow subgroups of S_4 are conjugate and isomorphic to D_4. Let $\mathbf{p = 3}$. A 3-Sylow subgroup of S_4 has order 3 and thus is generated by any 3-cycle. Thus, $P_3 \cong C_3$. <u>S_5; order $5! = 120 = 2^3 \cdot 3 \cdot 5$</u> Since the 2-,3-Sylow subgroups of S_5 will have the same order as the 2-,3-Sylow subgroups of S_4, we have $P_2 \cong D_4$ and $P_3 \cong C_3$.

□

Abelian Groups

Exercise 0.20. (Exercise 42)
Viewing $\mathbb{Z}, \mathbb{Q}$ as additive groups, show that $\mathbb{Q}/\mathbb{Z}$ is a torsion group, which has one and only one subgroup of order n for each integer $n \geq 1$, and that this subgroup is cyclic.

Proof. We have

$$\mathbb{Q}/\mathbb{Z} = \{x + \mathbb{Z} \colon x \in \mathbb{Q}\}.$$

Let $x + \mathbb{Z} \in \mathbb{Q}/\mathbb{Z}$ be an arbitrary nonzero element. Let $x = \frac{a}{b} \in \mathbb{Q}$. Then

$$b(x + \mathbb{Z}) = b\left(\frac{a}{b} + \mathbb{Z}\right) = a + \mathbb{Z} = 0 + \mathbb{Z} = 0.$$

Therefore, $x + \mathbb{Z}$ has finite order. Then, since $x + \mathbb{Z} \in \mathbb{Q}/\mathbb{Z}$ was arbitrary, every element of $\mathbb{Q}/\mathbb{Z}$ is a torsion element and, therefore, $\mathbb{Q}/\mathbb{Z}$ is a torsion group. For each integer $n \geq 1$, define

$$H_n \left\langle \frac{1}{n} + \mathbb{Z} \right\rangle.$$

Then $n\left(\frac{1}{n} + \mathbb{Z}\right) = 1 + \mathbb{Z} = 0$. Thus, $\frac{1}{n} + \mathbb{Z} \in \mathbb{Q}/\mathbb{Z}$ has order dividing n. If $0 < k < n$, then

$$k\left(\frac{1}{n} + \mathbb{Z}\right) = \frac{k}{n} + \mathbb{Z}$$

and $\frac{k}{n} \notin \mathbb{Z}$. Hence, $\frac{1}{n} + \mathbb{Z}$ has order exactly n. Thus,

$$H_n = \left\langle \frac{1}{n} + \mathbb{Z} \right\rangle \leqslant \mathbb{Q}/\mathbb{Z}$$

is a cyclic subgroup of order n. Suppose $K_n \leqslant \mathbb{Q}/\mathbb{Z}$ is a subgroup of order n. Thus, let $x + \mathbb{Z}$ be a generator for K_n. Write $x = \frac{a}{b} \in \mathbb{Q}$ in lowest terms. Then $x + \mathbb{Z}$ has order $b = n$. Thus, every generator of a subgroup of order n has the form

$$\frac{a}{n} + \mathbb{Z}$$

with $(a, n) = 1$. Then a is invertible modulo n and thus there exists some $c \in \mathbb{Z}$ such that

$$ac \equiv 1 \mod n$$

Thus $ac = 1 + nk$ with $k \in \mathbb{Z}$. Hence,

$$c\left(\frac{a}{n} + \mathbb{Z}\right) = \frac{ca}{n} + \mathbb{Z} = \frac{1 + nk}{n} + \mathbb{Z} = \frac{1}{n} + \mathbb{Z}.$$

Therefore,

$$\left\langle \frac{a}{n} + \mathbb{Z} \right\rangle = \left\langle \frac{1}{n} + \mathbb{Z} \right\rangle = H_n$$

and thus $\mathbb{Q}/\mathbb{Z}$ has one subgroup of order n, namely H_n, which is cyclic.

□