

Homework # 6

MATH 3160 – Complex Variables
Miguel Gomez

Completed: October 20, 2025

Problem 1

Find parameterized representations $z(t)$ of the following contours in the plane including t -ranges.

1. A straight line from point $(1 + 2i)$ to point $(i + 2)$
 2. A line from $(0, 0)$ to point $(1 + \sqrt{3}i)$
 3. A half-ellipse from point 2 to -2 passing through i centered at the origin. Recall that such an ellipse is defined by an equation of the form $\left(\frac{x}{a}\right)^2 + \left(\frac{y}{b}\right)^2 = 1$ in the xy -plane (for some real constants $a, b > 0$). *Hint: First find the suitable values of a and b defining the said ellipse. Then try parametrizing it similar to how $(\cos(t), \sin(t))$ parametrizes the unit circle.*
-

(a)

A straight line from point $(1 + 2i)$ to point $(i + 2)$


This one will require the expression for a line between points: $P + t(Q - P)$ where P and Q are the points and t runs from $0 \leq t \leq 1$.

$$P = (1 + 2i)$$

$$Q = (i + 2)$$

$$Q - P = (i + 2) - (1 + 2i) = 1 - i$$


$$\gamma(t) = 1 + 2i + t(1 - i)$$



(b)

A line from $(0, 0)$ to point $(1 + \sqrt{3}i)$.

This one is quite simple as we only have to multiply the point by t as the first point P is the origin and that handles moving from the origin to the point $(1 + \sqrt{3}i)$ as it moves from $0 \leq t \leq 1$.




(c)

A half-ellipse from point 2 to -2 passing through i centered at the origin.

$$x = a \cos(t) = 2 \cos(t)$$

$$y = b \sin(t) = \sin(t)$$

$$z(t) = x + iy = 2 \cos(t) + i \sin(t)$$



Problem 2

Evaluate the following integrals:

1. $\int_1^2 (\frac{1}{t} - i)^2 dt$
2. $\int_0^{\pi/6} e^{i2t} dt$
3. $\int_0^\infty e^{izt} dt$ where $\text{Im}(z) > 0$

(a)

$$\begin{aligned}
 \int_1^2 (\frac{1}{t} - i)^2 dt &= \int_1^2 (\frac{1}{t^2} - 2i\frac{1}{t} - i^2) dt \\
 &= \int_1^2 \frac{1}{t^2} dt - 2i \int_1^2 \frac{1}{t} dt + \int_1^2 dt \\
 &= -\frac{1}{3} \int_1^2 -3t^{-2} dt - 2i \ln(t) \Big|_1^2 + t \Big|_1^2 \\
 &= -\frac{1}{3} t^{-3} \Big|_1^2 + -2i \ln(t) \Big|_1^2 + t \Big|_1^2 \\
 &= \left(\frac{1}{3} - \frac{1}{3 * 2^3} \right) - 2i(\ln(2) - \ln(1)) + 1 \\
 &= \left(\frac{1}{3} - \frac{1}{3 * 2^3} + 1 \right) - 2i(\ln(2) - \ln(1))
 \end{aligned}$$

(b)

$$\begin{aligned}
 \int_0^{\pi/6} e^{i2t} dt &= \frac{1}{2i} \int_0^{\pi/6} 2ie^{i2t} dt \\
 &= \frac{1}{2i} e^{i2t} \Big|_0^{\pi/6} = -\frac{1}{2} i (e^{i\pi/3} - 1)
 \end{aligned}$$

(c)

Problem 3

Sketch the oriented curve defined by the following four contours and compute $\int_C f(z)dz$ where $f(z) = z - 1$:

1. C_1 : A semicircle $z = 2e^{i\theta}$ for $\theta \in [\pi, 2\pi]$.
 2. C_2 : A full circle $z = 2e^{i\theta}$ for $\theta \in [0, 2\pi]$.
 3. C_3 : A line on the real axis from 2 to -2 .
 4. $C_4 = C_1 + C_3$ where $+$ denotes concatenation.
-