

# Worksheet 2

MATH 3160  
Miguel Gomez

Completed: August 24, 2025

## Problem 1

Reduce each of these to a real number

---

(a)  $\frac{1+2i}{3-4i} + \frac{2-i}{5i}$

$$\begin{aligned}
 & \frac{1+2i}{3-4i} + \frac{2-i}{5i} = \\
 & \frac{(1+2i)(3+4i)}{(3-4i)(3+4i)} + \frac{(2-i)(-5i)}{(5i)(-5i)} = \\
 & \frac{(1+2i)(3+4i)}{9-16i^2} + \frac{(2-i)(-5i)}{-25i^2} = \\
 & \frac{(3+4i+6i+8i^2)^{-1}}{9-16i^2^{-1}} + \frac{(-10i+5i^2)^{-1}}{-25i^2^{-1}} = \\
 & \frac{(3+4i+6i-8)}{25} + \frac{(-10i-5)}{25} = \\
 & \frac{(3+4i+6i-8)}{25} + \frac{(-10i-5)}{25} = \\
 & \frac{(-5+10i)}{25} + \frac{(-10i-5)}{25} = \\
 & \frac{(-5-5+10i-10i)}{25} = \boxed{-\frac{2}{5}}
 \end{aligned}$$

(b)  $\frac{5i}{(1-i)(2-i)(3-i)}$

$$\begin{aligned}
 & \frac{5i}{(1-i)(2-i)(3-i)} = \frac{5i}{(2-i-2i+i^2)(3-i)} = \\
 & \frac{5i}{(1-3i)(3-i)} = \frac{5i}{(1-3i)(3-i)} = \\
 & \frac{5i}{-10i} = \boxed{-\frac{1}{2}}
 \end{aligned}$$

## Problem 2

Find the principal argument  $\text{Arg } z$  when..  
 preliminary necessary expressions:

$$\arg(z) = \text{Arg}(z) + 2 \cdot \pi \cdot k \quad ; k \in \mathbb{Z}$$

$$e^{i\theta} = \cos \theta + i \sin \theta$$

$$-\pi < \theta \leq \pi$$

(a)  $z = \frac{-2}{1+\sqrt{3}i}$

$$\begin{aligned} \frac{-2}{1+\sqrt{3}i} &= \frac{-2}{1+\sqrt{3}i} \frac{1-\sqrt{3}i}{1-\sqrt{3}i} = \\ \frac{-2(1-\sqrt{3}i)}{1-\sqrt{3}i+\sqrt{3}i+\sqrt{3}^2 i^2} &= \frac{-2+2\sqrt{3}i}{1-3} = \\ \frac{-2+2\sqrt{3}i}{-2} &= 1-\sqrt{3}i \\ 2 \left( -\frac{1}{2} + \frac{\sqrt{3}}{2}i \right) &= 2e^{i\frac{2\pi}{3}} \\ \therefore \text{Arg } z &= \boxed{\frac{2\pi}{3}} \end{aligned}$$

(b)  $z = \frac{2i}{i-1}$

$$\begin{aligned} \frac{2i}{i-1} &= \frac{2i(-i-1)}{(i-1)(-i-1)} = \frac{2i(-1-i)}{(-1+i)(-1-i)} = \\ \frac{(-2i-2i^2)}{1+i-i-i^2} &= \frac{(-2i-2i^2)}{1+i-i-i^2} = \frac{2-2i}{2} = 1-i = \sqrt{2}e^{-i\frac{\pi}{4}} \\ \therefore \text{Arg}(z) &= \boxed{-\frac{\pi}{4}} \end{aligned}$$

(c)  $z = (\sqrt{3}-i)^6$

For this one, we need to first include a factor of  $2^6$  and divide by it as well to bring a half into the parentheses.

$$\begin{aligned} (\sqrt{3}-i)^6 &= 2^6 \left( \frac{\sqrt{3}}{2} - \frac{1}{2}i \right)^6 \\ 2^6 (e^{-i\frac{\pi}{6}})^6 &= 2^6 e^{-i\pi} \\ \therefore \text{Arg}(z) &= \boxed{\pi} \end{aligned}$$

### Problem 3

For the next few questions write the individual factors on the left in exponential form, perform the needed operations on complex numbers, and finally change back to rectangular coordinates *Show that*:

(a)  $i(1 - \sqrt{3}i)(\sqrt{3} + i) = 2(1 + \sqrt{3}i)$

To convert into exponential form, we can add factors to get us the correct exponentials in normalized form:

$$\begin{aligned} i &= e^{i\frac{\pi}{2}} \\ (1 - \sqrt{3}i) &= 2 \left( \frac{1}{2} - \frac{\sqrt{3}}{2}i \right) = 2e^{-i\frac{\pi}{3}} \\ (\sqrt{3} + i) &= 2 \left( \frac{\sqrt{3}}{2} + \frac{1}{2}i \right) = 2e^{i\frac{\pi}{6}} \end{aligned}$$

Note,  $(\sqrt{3} + i)$  is  $90^\circ$  rotated from  $(1 - \sqrt{3}i)$ . This is easily verified by adding  $\pi/2$  to  $\text{Arg}(1 - \sqrt{3}i)$ .

$$\begin{aligned} \therefore i(1 - \sqrt{3}i) &= e^{i\frac{\pi}{2}} 2e^{-i\frac{\pi}{3}} = 2e^{i(\frac{\pi}{2} - \frac{\pi}{3})} \\ 2e^{i(\frac{3\pi}{6} - \frac{2\pi}{6})} &= 2e^{i\frac{\pi}{6}} \end{aligned}$$

Including the next factor gives:

$$2e^{i\frac{\pi}{6}} 2e^{i\frac{\pi}{6}} = 4e^{i\frac{2\pi}{6}} = 4e^{i\frac{\pi}{3}} = \boxed{2(1 + \sqrt{3}i)}$$

(b)  $(\sqrt{3} + i)^6 = -64$

This is the same as problem 2c with the angle being  $\pi/6$  instead of the negative angle. Same work shows the result:

$$\begin{aligned} (\sqrt{3} + i)^6 &= 2^6 \left( \frac{\sqrt{3}}{2} + \frac{1}{2}i \right)^6 \\ 2^6 (e^{i\frac{\pi}{6}})^6 &= 2^6 e^{i\pi} = 2^6 (-1) = \boxed{-64} \end{aligned}$$

(c)  $(1 + \sqrt{3}i)^{-10} = 2^{-11}(-1 + \sqrt{3}i)$

Here we can first rationalize the fraction

$$\begin{aligned} (1 + \sqrt{3}i)^{-10} &= \left( \frac{1}{1 + \sqrt{3}i} \right)^{10} = \\ \left( \frac{1 - \sqrt{3}i}{(1 + \sqrt{3}i)(1 - \sqrt{3}i)} \right)^{10} &= \left( \frac{1 - \sqrt{3}i}{1 - \sqrt{3}i + \sqrt{3}i - \sqrt{3}^2 i^2} \right)^{10} = \\ \left( \frac{1 - \sqrt{3}i}{4} \right)^{10} &= 2^{-11} \left( \frac{1 - \sqrt{3}i}{2} \right)^{10} = \\ 2^{-11} (e^{-i\frac{\pi}{3}})^{10} &= 2^{-11} e^{-i\frac{10\pi}{3}} = 2^{-11} e^{-i(2\pi + \frac{4\pi}{3})} \\ 2^{-11} e^{-i(\pi + \frac{1\pi}{3})} &= 2^{-11} e^{-i(\frac{-2\pi}{3})} = \boxed{2^{-11}(-1 + \sqrt{3}i)} \end{aligned}$$

## Problem 4

Use exponential form to find  $(1 - i)^5$

$$\begin{aligned}(1 - i)^5 &= \left(\frac{2}{\sqrt{2}}\right)^5 \left(\frac{\sqrt{2}}{2} - \frac{\sqrt{2}}{2}i\right)^5 = \left(\frac{2}{\sqrt{2}}\right)^5 (e^{-i\frac{\pi}{4}})^5 = \\ &\text{angle } \frac{-5\pi}{4} \text{ corresponds to angle } \frac{3\pi}{4} \\ &\left(\frac{2}{\sqrt{2}}\right)^5 e^{-i\frac{5\pi}{4}} = \left(\frac{2}{\sqrt{2}}\right)^5 e^{i\frac{3\pi}{4}} = \\ &\left(\frac{2}{\sqrt{2}}\right)^5 \left(-\frac{\sqrt{2}}{2} + \frac{\sqrt{2}}{2}i\right) = \left(\frac{2}{\sqrt{2}}\right)^4 (-1 + i) \\ &\text{recall } \frac{2}{\sqrt{2}} = \frac{2\sqrt{2}}{\sqrt{2}^2} = \sqrt{2} \\ &\therefore \left(\frac{2}{\sqrt{2}}\right)^4 = \sqrt{2}^4 = 2^2 = 4 \\ &\therefore (1 - i)^5 = \boxed{4(-1 + i)}\end{aligned}$$