


# ORNIS SCANDINAVICA

INTERNATIONAL JOURNAL OF ORNITHOLOGY

Hans Källander & Olof Rydén:

Inter-observer differences in studies of visible  
migration at Falsterbo



VOL. 5, NO. 1, 1974

UNIVERSITETSFÖRLAGET

# Inter-observer differences in studies of visible migration at Falsterbo

HANS KÄLLANDER & OLOF RYDÉN

Källander, H. & Rydén, O. Inter-observer differences in studies of visible migration at Falsterbo. *Ornis Scand.* 5, 53–62, 1974.

Inter-observer differences were studied at Falsterbo on four occasions. Sixteen experienced observers, four on each occasion, simultaneously and independently recorded the migration. No single observer could record the entire migration even under moderate conditions. Observers were relatively efficient in recording raptors and abundant and easily detected species. Observers were compared with regard to the number of species and individuals recorded. Any two observers both recorded 50–80 % of all species in their pooled records. For species occurring in sufficient numbers for analysis, the inter-observer agreement, i. e. the percentage of individuals two observers both recorded related to the highest number recorded by either of them, ranged between 26 and 88 %; for five species that were well represented on more than one occasion the corresponding proportions varied between 59 and 80 %.

Hans Källander, Lund University, Dept. of Animal Ecology, Ecology Building, S-223 62 Lund, Sweden.

Olof Rydén, Lund University, Psychological Laboratory, S-223 50 Lund, Sweden.

## INTRODUCTION

Much information on the diurnal migration of birds has been collected by systematic observation in localities where heavy concentrations of migrating birds are regularly found. Our knowledge of the effects of weather on migration and of the seasonal distribution of migration of many species is primarily obtained from such studies. But there are difficulties: it is known, for instance, that a substantial and varying part of the total migration takes place at heights where it is impossible to see the birds from the ground (Eastwood 1967). Furthermore, meteorological conditions, especially the speed and direction of the wind, influence the exact migration path. The varying accessibility of the migration to observation thus reduces the reliability of visual counts, and

comparisons between years and days must be made with great caution.

Even if only highly competent ornithologists record visible migration, their ability to detect and record passing birds differs. As a consequence, an observed difference in numbers between two seasons for a given species is to an unknown extent attributable to differences in the capacity of the observers. The magnitude of this source of error is poorly known. Enemar (1964) concluded that skilled ornithologists overlook, on average, every second individual and four out of ten species. Källander et al. (1972) found that when recording sea-bird migration through Kalmar Sound in south-east Sweden, competent observers agreed relatively well with regard to the numbers of the conspicuous and dominating Eider *Somateria mollissima*, whereas inter-observer differences in-

creased dramatically for less abundant species. Sea bird migration is relatively easily observed at least in this locality, since few species are involved, and observation from the land can be concentrated in one direction.

To estimate the magnitude of inter-observer differences in recording visible migration we decided to conduct a series of experiments at Falsterbo (13°50'E, 55°20'N), where land birds make up most of the migration. Because of the larger number of species and individuals involved, as well as the complexity of the migration itself, it seemed to us that this would be an ideal locality.

## METHOD

Our procedure was partly adopted from Enemar (1964). Four observers independently recorded the migration. They were separated by screens at 8 m intervals and equipped with similar binoculars. They were instructed to record all migrating birds. Another group of three or four ornithologists (control group) simultaneously observed the migration. In the present paper the observers' records have not generally been related to those of the control group. Observations were carried out on four separate days. The observation period on each occasion was about three hours, giving a total period of observation of 12.5 hours. The observers were experienced ornithologists, who were well acquainted with the locality. In all 16 observers took part.

## OBSERVATION PERIODS

For a general description of the topography of the Falsterbo peninsula and for an account of its bird migration, the reader is referred to Rudebeck (1950). All observations were made in September and October when migration is at its maximum at Falsterbo. Weather conditions and the main features of migration on the different occasions are briefly described below.

### 19 October 1968

Observations took place between 0740 and 0940

and between 1010 and 1200 hrs. There was a weak northwesterly wind and the temperature was about 3–4° C. Initially the sky was completely overcast, but during the period it cleared and the temperature rose. Visibility was good.

Several features of the migration complicated observation. Firstly, many species were involved with different patterns of migration. Secondly, certain groups such as pigeons *Columba*, corvids and Starlings *Sturnus vulgaris* not only passed in relatively large numbers but also on a broad front. A large proportion of the buzzards *Buteo* returned after attempting migration, and sometimes buzzards were simultaneously flying in two or more directions, which meant that the observers had to spend much time and effort to establish which individuals actually left the peninsula.

### 14 October 1969

Observations took place between 0720 and 0910 and between 0925 and 0955 hrs. The sky was overcast, but visibility was fairly good. There was a weak southwesterly wind and a temperature of about 8°C.

Starlings, *Fringilla* finches, and cardueline finches dominated. The flocks passed as a comparatively narrow stream above the observers, which facilitated detection, but also at different levels at great height. This, together with the fact that migration was fairly intense, rendered observation less easy.

### 19 September 1970

Observations took place between 0610 and 0910 and between 1010 and 1105 hrs. The sky was overcast, clearing up towards the end of the period. There was a weak to moderate, gradually waning wind from the southwest, and visibility was good. The temperature was 13°C at the start, thereafter rising.

The migration was intense and dominated by a continuous passage of Chaffinches *Fringilla coelebs* and some Siskins *Carduelis spinus* at a relatively great height. Siskins also passed low in flocks. Loose assemblages of Meadow

Pipits *Anthus pratensis*, Yellow *Motacilla flava* and White *Motacilla alba* Wagtails passed near the ground. Tree Pipits *Anthus trivialis* were often heard from a great height but were difficult to see. Hirundines, mostly Swallows *Hirundo rustica*, migrated on a broad, low front.

The complicated character of the migration made it highly unlikely that a single observer could record simultaneously the perpetual stream of Chaffinches and the various species passing low around him.

#### 25 October 1970

Observations took place between 0750 and 1010 hrs. The sky was overcast at the start and at the end, with a clear period in between. There was a weak to moderate south south-westerly wind and the temperature was about 10°C.

The migration was exceptionally uncomplicated. Few species were involved; Starlings dominated and passed close to the observers in flocks of tens or a couple of hundreds. Meadow Pipits, Siskins, Linnets *Carduelis cannabina*, and Greenfinches *Carduelis chloris* were also easily detected. Some flocks of Linnets, Greenfinches and Sparrowhawks *Accipiter nisus* presented some difficulties to the observers, since they tended to return after attempted migration.

### RESULTS AND DISCUSSION

Records from all observers and the control group are shown in Tables I–IV. Sea-birds and waders are excluded since observation conditions for some of them were not quite equal for all observers.

#### 19 October 1968 (Table I)

Considerable differences were found for corvids. A, C, and D agreed fairly well on the number of Jackdaws *Corvus monedula*, while B reported a much lower number. The variation for the Rook *Corvus frugilegus* was 58–272 and for the Hooded Crow *Corvus corone* 209–833. Some of these discrepancies may be explained by the fact that adverse light made

separation of Rooks and Crows at a distance rather difficult. The figures for the Starling range from 2,735 for A to only 471 for B.

Considering less abundant species, only B noticed the migrating finches to any extent and was outstanding in sorting out the Roughlegged Buzzards *Buteo lagopus*. In general, however, B's capacity was somewhat lower than that of the others.

Up to 200 Magpies *Pica pica* repeatedly attempted migration thereby creating a very confusing situation. This resulted in 51 and 95 individuals being erroneously recorded as migrants by two observers.

#### 14 October 1969 (Table II)

Starlings, *Fringilla*, and cardueline finches dominated. There is a relatively low variation in the figures for the Starling – from 7,044 to 8,875 recorded individuals. While there is high consistency in the numbers of cardueline finches as a group, the totals for the different species show considerable variation. This is partly a consequence of a varying number of birds not identified as to species. While two of the observers agreed closely with respect to the number of Greenfinches and Siskins recorded, 719 against 738 and 713 against 775 respectively, the greatest difference between any two observers for these species is much larger, 89 against 738 and 713 against 2,422 respectively. The *Fringilla* finches, which constituted around 50% of total migration, were estimated to 4,507 by F, while H recorded more than twice as many, 10,652.

#### 19 September 1970 (Table III)

Relative to the control group, a few species, viz. the Chaffinch and the Siskin, were generally overlooked, probably partly due to the high mean intensity of the migration (640 birds/min) and partly because they passed rather high. Regarding the Chaffinch, three observers showed a remarkable agreement, recording 16,800, 17,090, and 17,980 individuals respectively, while the fourth observer reported 43,610. The picture is similar for the Siskin;

Table I. Recorded migration on 19 October 1968. Co = control group, A, B, C, and D = the individual observers. Minus sign denotes return migration

	Co	A	B	C	D
<i>Buteo buteo</i>	774	246	141	465	581
<i>B. lagopus</i>	10	3	15	2	7
<i>Buteo sp.</i>	264	587	554	0	48
<i>Accipiter nisus</i>	154	69	72	51	74
<i>Milvus milvus</i>	1	0	0	1	1
<i>Haliaetus albicilla</i>	1	0	0	0	0
<i>Circus cyaneus</i>	1	0	0	0	0
<i>Circus sp.</i>	0	0	1	0	0
<i>Falco columbarius</i>	0	1	0	0	0
<i>F. tinnunculus</i>	2	0	2	0	0
<i>Columba oenas</i>	34	16	5	41	56
<i>C. palumbus</i>	6,163	8,871	1,289	8,957	6,103
<i>C. livia 'domestica'</i>	5	0	0	2	7
<i>Streptopelia decaocto</i>	1	0	0	0	6
<i>Streptopelia sp.</i>	0	0	0	0	2
<i>Dendrocopos major</i>	0	0	0	3	5
<i>Lullula arborea</i>	30	3	0	0	8
<i>Alauda arvensis</i>	23	24	22	0	17
<i>Eremophila alpestris</i>	0	2	0	0	0
<i>Hirundo rustica</i>	44	15	16	6	27
<i>Corvus corone</i>	783	833	209	330	329
<i>C. frugilegus</i>	240	79	272	58	148
<i>C. monedula</i>	2,126	690	362	951	790
<i>Corvus sp.</i>	95	177	540	130	0
<i>Pica pica</i>	0	0	51	95	-9
<i>Nucifraga caryocatactes</i>	2	0	2	2	2
<i>Garrulus glandarius</i>	2	-1	0	0	0
<i>Parus major</i>	0	0	0	2	0
<i>P. caeruleus</i>	46	0	0	5	2
<i>Turdus pilaris</i>	61	5	3	31	12
<i>T. philomelos</i>	1	0	1	0	0
<i>T. musicus</i>	24	0	40	-2	0
<i>Turdus sp.</i>	50	0	934	0	13
<i>Prunella modularis</i>	1	2	6	1	16
<i>Anthus pratensis</i>	0	1	1	4	6
<i>A. spinoletta</i>	0	1	0	0	0
<i>Lanius excubitor</i>	0	0	1	0	0
<i>Sturnus vulgaris</i>	4,387	2,735	471	2,125	1,635
<i>Coccothraustes coccothraustes</i>	0	0	0	0	2
<i>Carduelis chloris</i>	8	0	0	0	0
<i>C. carduelis</i>	8	14	6	-16	7
<i>C. spinus</i>	5	2	5	2	0
<i>C. cannabina</i>	3	0	0	6	0
<i>Pyrrhula pyrrhula</i>	19	9	0	0	2
<i>Loxia sp.</i>	10	0	0	0	0
<i>Fringilla coelebs</i>	54	20	104	0	0
<i>F. montifringilla</i>	35	2	30	5	0
<i>Fringilla sp.</i>	310	5	273	20	35
<i>Emberiza citrinella</i>	43	21	0	-6	5
<i>E. schoeniclus</i>	1	3	3	1	0
<i>Plectrophenax nivalis</i>	1	0	0	0	0
<i>Passer sp.</i>	0	0	0	0	15
<i>Passeriformes non det.</i>	10	25	0	0	94
Total	19,252	14,461-1	7834	13,333-24	10,055-24
Number of identified species	35	26	25	27	26

Table II. Recorded migration on 14 October 1969. Co = control group, E, F, G, and H = the individual observers. Minus sign denotes return migration

	Co	E	F	G	H
<i>Columba oenas</i>	0	40	30	0	16
<i>C. palumbus</i>	45	55	15	0	15
<i>Columba sp.</i>	-29	0	40	115	0
<i>Alauda arvensis</i>	114	29	43	44	20
<i>Hirundo rustica</i>	26	9	16	2	13
<i>Corvus corone</i>	4	3	6	5	1
<i>C. frugilegus</i>	3	4	0	0	7
<i>C. monedula</i>	13	16	16	7	7
<i>Corvus sp.</i>	0	0	0	7	0
<i>Turdus musicus</i>	0	0	0	0	2
<i>Anthus pratensis</i>	151	120	168	100	38
<i>A. trivialis</i>	1	0	0	6	1
<i>A. spinoletta</i>	0	0	1	0	1
<i>Anthus sp.</i>	6	0	0	2	36
<i>Motacilla alba</i>	2	0	5	0	7
<i>Lanius excubitor</i>	1	1	0	0	0
<i>Sturnus vulgaris</i>	9,178	7,044	8,081	7,250	8,875
<i>Carduelis chloris</i>	908	719	738	89	343
<i>C. carduelis</i>	8	14	3	0	0
<i>C. spinus</i>	1,800	2,422	1,442	713	775
<i>C. cannabina</i>	437	263	361	404	262
<i>C. flavirostris</i>	10	0	3	0	0
<i>Carduelis sp.</i>	254	220	25	1,548	480
<i>Fringilla coelebs</i>	105	27	0	29	23
<i>Fringilla montifringilla</i>	0	0	6	0	12
<i>Fringilla sp.</i>	11,724	7,073	4,501	5,073	10,617
<i>Emberiza citrinella</i>	12	10	4	9	3
<i>E. schoeniclus</i>	27	20	49	13	18
<i>Passer montanus</i>	0	1	0	0	1
<i>Passer sp.</i>	0	0	0	2	0
Passeriformes non det.	76	0	0	343	0
Total	24,905-29	18,090	15,553	15,761	21,573
Number of identified species	19	18	18	13	21

one observer reported only 40 individuals, the others 365, 536, and 584 respectively.

The buzzards received everyone's attention; the variation in the total number recorded is moderate – from 294 to 407 – and corresponds fairly well with the estimate of the control group (444). It is interesting to note, however, how the identification as to species differs between observers. The highest number of Common Buzzards *Buteo buteo* recorded is 321, the lowest 6, while the corresponding figures for the Honey Buzzard *Pernis apivorus* are 172 and 65 (cf. Enemar 1964).

For the Sparrowhawk, Swallow, and White Wagtail, three observers agreed reasonably well, while the remaining observer recorded a much

higher or lower total. Records for the Meadow Pipit, Sand Martin *Riparia riparia*, and Starling show good overall agreement.

#### 25 October 1970

Compared with the preceding experiment this one was less demanding – one single species, the Starling, dominated. Since the Starlings passed very close to the observers and were easy to detect, the variation in totals, from 3,770 to 6,715, is probably largely an effect of differing flock-size estimates. With a few exceptions, there is a remarkable agreement for most other species. Thus one of the observers deviated markedly from the others in recording

Table III. Recorded migration on 19 September 1970. Co = control group, I, K, L, and M = the individual observers

	Co	I	K	L	M
<i>Buteo buteo</i>	45	321	6	96	135
<i>Pernis apivorus</i>	291	65	172	97	72
<i>Buteo/Pernis</i>	108	21	116	198	145
<i>Accipiter nisus</i>	76	69	43	49	48
<i>Milvus milvus</i>	0	1	0	0	0
<i>Pandion haliaetus</i>	1	1	1	1	1
<i>Falco columbarius</i>	0	1	0	0	1
<i>F. tinnunculus</i>	0	2	1	1	1
<i>Falco sp.</i>	0	0	1	0	0
<i>Columba oenas</i>	34	14	21	7	2
<i>C. palumbus</i>	4	4	0	4	12
<i>C. livia 'domestica'</i>	5	0	4	0	0
<i>Hirundo rustica</i>	2,016	1,121	639	1,121	1,037
<i>Delichon urbica</i>	22	4	9	2	10
<i>Riparia riparia</i>	112	25	37	31	30
Hirundinidae	180	0	209	0	0
<i>Corvus corone</i>	1	0	1	1	0
<i>Turdus philomelos</i>	1	0	0	0	0
<i>Oenanthe oenanthe</i>	1	0	1	0	0
<i>Prunella modularis</i>	3	0	1	0	1
<i>Anthus pratensis</i>	1,113	962	734	945	842
<i>A. campestris</i>	0	0	0	0	1
<i>A. trivialis</i>	47	96	195	311	343
<i>A. cervinus</i>	7	6	4	2	5
<i>A. spinoletta</i>	0	6	0	0	0
<i>Anthus sp.</i>	1	0	0	0	0
<i>Motacilla alba</i>	72	63	114	510	126
<i>M. flava</i>	59	70	123	105	104
<i>Sturnus vulgaris</i>	1,099	550	474	555	522
<i>Carduelis chloris</i>	6	0	13	26	0
<i>C. spinus</i>	1,577	365	584	40	536
<i>C. cannabina</i>	6	244	0	1	27
<i>Fringilla coelebs</i>	143,358	16,800	43,610	17,090	17,980
<i>Emberiza schoeniclus</i>	4	3	5	0	2
Passeriformes non det.	3	200	0	140	0
Total	150,252	21,014	47,118	21,333	21,983
Number of identified species	25	23	23	21	23

more Meadow Pipits and Greenfinches and was the only one to report Bramblings *Fringilla montifringilla*. On the other hand, this observer recorded no Linnets.

#### *Recording efficiency for different categories of the migrating birds*

One way to compare the observers is to analyse the way their relative efficiency is distributed among different aspects of the total migration. This was done for each period of observation by establishing categories of birds

which differ in their mode of migration, e. g. solitary vs. migrating in flocks, scattered vs. defined in space, fast vs. slow, etc. Each observer's effectiveness, relative to the control group, was then ranked, and the rankings of all observers compared. The correlation between the rankings for all observers is expressed by the Kendall coefficient of concordance ( $W$ ).

In the first period of observation the following six categories were established: Sparrowhawk, buzzards, pigeons, fringillids, Starling, and corvids. The test failed to show any correlation between the observers ( $W=0.35$ , n. s.),

Table IV. Recorded migration on 25 October 1970. Co = control group, N, O, P, and R = the individual observers

	Co	N	O	P	R
<i>Ardea cinerea</i>	0	1	0	0	1
<i>Buteo buteo</i>	0	0	0	0	1
<i>B. lagopus</i>	2	2	1	2	0
<i>Buteo sp.</i>	0	0	0	0	1
<i>Accipiter nisus</i>	20	35	23	23	20
<i>Columba livia</i> 'domestica'	1	1	1	0	1
<i>Alauda arvensis</i>	17	30	17	13	9
<i>Corvus corone</i>	19	20	9	27	16
<i>C. frugilegus</i>	5	1	26	13	11
<i>C. monedula</i>	17	11	20	19	22
<i>Corvus sp.</i>	0	25	32	0	25
<i>Turdus pilaris</i>	1	2	30	1	0
<i>Anthus pratensis</i>	54	39	82	40	45
<i>A. spinoletta</i>	0	0	1	0	1
<i>Bombycilla garrulus</i>	0	16	0	0	0
<i>Sturnus vulgaris</i>	7,779	6,103	5,405	3,770	6,715
<i>Carduelis chloris</i>	182	152	220	135	138
<i>C. carduelis</i>	4	4	2	5	4
<i>C. spinus</i>	100	73	67	74	88
<i>C. cannabina</i>	97	33	0	29	61
<i>C. flavirostris</i>	10	37	16	14	25
<i>Carduelis sp.</i>	60	0	0	26	0
<i>Loxia sp.</i>	5	5	5	14	5
<i>Fringilla montifringilla</i>	0	0	141	0	0
<i>Fringilla sp.</i>	24	0	0	0	0
Fringillidae	0	0	52	0	0
<i>Emberiza citrinella</i>	29	14	16	14	9
Passeriformes non det.	0	114	0	0	203
Total	8,427	6,721	6,166	4,219	7,402
Number of identified species	16	18	17	15	17

which means that their attention, or ability, was unevenly distributed on these categories of birds. Although no consistent pattern was found, buzzards and pigeons usually ranked highest.

On the second occasion, the same test was applied to the categories Skylark *Alauda arvensis*, Meadow Pipit, Starling, cardueline finches, and *Fringilla* finches. A significant agreement between the observers was found ( $W = 0.64$ ,  $P < 0.05$ ). Compared with the control group the observers generally concentrated on the Starlings and cardueline finches.

There is a high correlation between the observers in their relative effectiveness for different categories on the third occasion ( $W = 0.91$ ,  $P < 0.01$ ; categories: buzzards, Sparrowhawk, hirundines, Meadow Pipit, Starling, and

Chaffinch plus Siskin). The raptors and Meadow Pipit obtained the highest rankings.

Probably as a consequence of the low number of species that took part in the migration, there is a high correlation in the distribution of the effectiveness of the observers on the last occasion ( $W = 0.81$ ,  $P < 0.01$ , calculated on the categories Sparrowhawk, Siskin plus Twite *Carduelis flavirostris*, Greenfinch plus Linnet, and Starling). The Sparrowhawk ranked highest followed by Siskin plus Twite.

The only occasion when no correlation was established, i. e. the first one, was characterized by more species than the others. It seems likely that with an increased number of species the chance becomes less that the observers make the same species preferences. When the same preferences are made they apparently favour



raptors and species that are easy to detect (provided that the recording efficiency of the control group is reasonably evenly distributed among the different categories of birds).

#### *Comparisons of the number of species recorded*

To measure their ability to record the qualitative aspects of migration, the observers were compared in couples, the number of species detected by both observers being expressed as a percentage of that observed by either or both. The percentage therefore indicates to what extent two observers agreed upon which




Fig. 1. Comparison of the number of species recorded. The number of species recorded by both observers is expressed as a percentage of that observed by either or both. The values are arranged in decreasing magnitude.

species took part in migration (Fig. 1). This percentage varies from 50 to 84 and corresponds with the W-values presented above (56 % – 0.35, 64 % – 0.64, 73 % – 0.91, and 75 % – 0.81). Naturally, when the observers direct their attention to different aspects of the migration in a similar way they tend to detect and record the same species.

#### *Comparisons of the number of individuals recorded*

The number of individuals recorded for each species often varies considerably between observers (Tables I–IV). The most notable of these differences have already been pointed out. An attempt was made to quantify the differences by comparing all observers on each occasion two at a time. The lowest number of individuals recorded for each species presented as a percentage of the highest in each pair expresses the degree of agreement between the observers. For this analysis, species were selected that migrated in sufficient numbers and in a typical pattern to make comparisons meaningful. The results are shown in Table V. There is a concentration of values between 50 and 80 % with mean correspondences around 60 %.

When inter-observer agreement on species content (Fig. 1) and that on number of individuals (Table V) are compared, a similar distribution of means is found on the four occasions, indicating a positive correlation between these two measurements.

In Table VI five species have been selected that were well represented on 2–4 occasions, making 12–24 pairwise comparisons available. Means and standard error (calculated separately for each occasion and averaged) were computed to get a more generalized measurement of inter-observer agreement.

Inter-observer differences are much more difficult to evaluate for species that are represented by only a few individuals. Nevertheless, some of these are interesting for two reasons: they normally occur in small numbers each season and are considered easily detected. As examples in our observation pe-

Table V. Percentage agreement between pairs of observers for selected species. The percentage is calculated on the basis of the higher of the two records of each pair

## First occasion, 19 Oct. 1968

Pair of observers	A-B	A-C	A-D	B-C	B-D	C-D	Mean
<i>Accipiter nisus</i>	96	74	93	71	97	69	85
<i>Buteo buteo</i>	57	53	42	30	24	80	48
<i>Columba palumbus</i>	15	99	69	14	21	68	48
<i>Sturnus vulgaris</i>	17	78	60	22	29	77	47
<i>Corvus monedula</i>	52	73	87	38	46	83	62
<i>C. frugilegus</i>	29	73	53	21	86	39	50
<i>C. corone</i>	25	40	39	63	64	100	55
Mean	42	70	63	37	52	74	

## Second occasion, 14 Oct. 1969

Pair of observers	E-F	E-G	E-H	F-G	F-H	G-H	Mean
<i>Alauda arvensis</i>	67	66	69	98	47	45	65
<i>Anthus pratensis</i>	71	83	32	60	23	38	51
<i>Sturnus vulgaris</i>	87	97	79	90	91	82	88
<i>Carduelis chloris</i>	97	12	48	12	46	26	40
<i>C. spinus</i>	60	29	32	49	54	92	53
<i>C. cannabina</i>	73	65	100	89	73	65	78
<i>Emberiza schoeniclus</i>	41	65	90	27	38	72	56
Mean	71	60	64	61	53	60	

## Third occasion, 19 Sept. 1970

Pair of observers	I-K	I-L	I-M	K-L	K-M	L-M	Mean
<i>Buteo buteo</i>	2	30	42	6	4	71	26
<i>Pernis apivorus</i>	38	67	90	56	42	74	61
<i>Accipiter nisus</i>	62	71	70	88	90	98	80
<i>Hirundo rustica</i>	57	100	93	57	62	93	77
<i>Riparia riparia</i>	68	81	83	84	81	97	82
<i>Anthus pratensis</i>	76	98	88	77	87	89	86
<i>A. trivialis</i>	49	31	28	63	57	91	53
<i>Motacilla alba</i>	55	12	50	22	90	25	42
<i>M. flava</i>	57	67	67	85	85	100	77
<i>Sturnus vulgaris</i>	86	99	95	85	91	94	92
<i>Carduelis spinus</i>	63	11	68	7	92	7	41
<i>Fringilla coelebs</i>	39	98	93	39	41	95	68
Mean	54	64	72	56	69	78	

## Fourth occasion, 25 Oct. 1970

Pair of observers	N-O	N-P	N-R	O-P	O-R	P-R	Mean
<i>Accipiter nisus</i>	66	66	57	100	87	87	77
<i>Alauda arvensis</i>	57	43	30	76	69	69	49
<i>Corvus frugilegus</i>	4	8	9	50	42	85	33
<i>Anthus pratensis</i>	48	98	87	49	55	89	71
<i>Sturnus vulgaris</i>	89	62	91	70	80	56	75
<i>Carduelis chloris</i>	69	89	91	61	63	98	79
<i>C. spinus</i>	92	99	83	91	76	84	88
Mean	61	66	64	71	67	81	

riods can be mentioned Heron *Ardea cinerea*, Sea Eagle *Haliaeetus albicilla*, Red Kite *Milvus milvus*, Osprey *Pandion haliaetus*, and Hen Harrier *Circus cyaneus* (all of which passed close to the observers; Tables I, III, IV). It was found that 9 records were made of these birds out of a total of 24 possible. Thus, when recording the entire migration, experienced ornithologists can overlook even these birds.

## CONCLUSIONS

Any conclusions to be drawn from these experiments must be based on a consideration of the generality of the results. Although experienced ornithologists, the participating observers did not have day-to-day experience, which no doubt increases the capability of a regular observer. The effectiveness of a regular observer, however, necessarily varies which is probably an unavoidable result of day-long, laborious observation. On the other hand, our observers knew that their records were going to be compared, which almost certainly motivated them to do their best. Since the observation periods lasted for only a few hours, their vigilance was not affected by tiredness or boredom. The experimental situation apparently compensated, at least partly, for the less day-to-day experience of our observers.

Inter-observer differences occur due to discrepancies in flock size estimates, species identification, and, undoubtedly most important, varying ability to detect the birds. The average amount of error is difficult to estimate, because it varies according to characteristics of the migration, weather conditions, and differential effectiveness of the observer. Only for a few species have sufficient data been obtained to

allow such an estimate. If the data of Table VI are accepted as representative of the species involved, one can conclude that if the records of two experienced observers are compared, one would on average expect the following proportions of obtained differences to be attributable to inter-observer differences: Sparrowhawk 10–30 %, Meadow Pipit 15–45 %, Starling 15–35 %, Greenfinch 20–60 %, and Siskin 25–55 %. The ranges represent approximately the two standard error limits.

If one wants to apply our results to data collected at Falsterbo one must bear in mind that inter-observer differences can either accumulate or cancel each other out, and that the chances of cancelling out would increase with more observers involved. If one deals with a trend, the trend would, of course, become less and less influenced by inter-observer differences the more years that are included. One might add that other sources of error affect observation data, with weather conditions probably being the most important.

## ACKNOWLEDGEMENTS

We thank all who participated in our observations, both observers and members of the control group: Thomas Alerstam, Nils-Ake Andersson, Carl-Axel Bauer, Jens Bruun, Rune Gerell, Bodil and Krister Hjalte, Niels Kjellén, Gunnar Lid, Henning Noer, Jørgen Rabøl, Torben Rasmussen, Gunnar Roos, Gustaf Rudebeck, Bo Svensson, Magnus Sylvén, Stafan Ulfstrand, Krister Wadén, and Gerd Østerbye.

## REFERENCES

- Eastwood, E. 1967. *Radar Ornithology*. Methuen, London.
- Enemar, A. 1964. A preliminary estimation of the reliability of the registration work of four ornithologists watching autumn bird migration. (Swedish with English summary.) *Vår Fågelvärld* 23, 1–25.
- Källander, H., Rydén, O. & Weikert, C. 1972. Unterschiede in der Beobachtungs-Effektivität bei der Registrierung von Küsten-Seevogelzug. *Vogelwarte* 26, 303–310.

Received April 1973, revised February 1974  
Published April 1974

Table VI Mean percentage agreement between pairs of observers for five well represented species

Species	Number of possible pair comparisons	Mean $\pm$ S. E.
<i>Accipiter nisus</i>	18	80 $\pm$ 5.5
<i>Anthus pratensis</i>	18	69 $\pm$ 6.8
<i>Sturnus vulgaris</i>	24	75 $\pm$ 4.5
<i>Carduelis chloris</i>	12	59 $\pm$ 9.0
<i>C. spinus</i>	18	61 $\pm$ 8.5