

Adv Exp Med Biol. Author manuscript; available in PMC 2013 August 12.

Published in final edited form as:

Adv Exp Med Biol. 2012; 740: 1145-1174. doi:10.1007/978-94-007-2888-2_52.

Remodeling of Calcium Handling in Human Heart Failure

Qing Lou, Ph.D., Ajit Janardhan, M.D., Ph.D., and Igor R. Efimov, Ph.D. Department of Biomedical Engineering, Washington University in St. Louis, MO 63130, USA. (Q.L., I.R.E.). Department of Medicine, Division of Cardiology, Washington University School of Medicine, St. Louis, MO 63110, USA (A.J., I.R.E.)

Abstract

Heart failure (HF) is an increasing public health problem accelerated by a rapidly aging global population. Despite considerable progress in managing the disease, the development of new therapies for effective treatment of HF remains a challenge. To identify targets for early diagnosis and therapeutic intervention, it is essential to understand the molecular and cellular basis of calcium handling and the signaling pathways governing the functional remodeling associated with HF in humans. Calcium (Ca²⁺) cycling is an essential mediator of cardiac contractile function, and remodeling of calcium handling is thought to be one of the major factors contributing to the mechanical and electrical dysfunction observed in HF. Active research in this field aims to bridge the gap between basic research and effective clinical treatments of HF. This chapter reviews the most relevant studies of calcium remodeling in failing human hearts and discusses their connections to current and emerging clinical therapies for HF patients.

1. Introduction

HF is a rising public health problem, with a prevalence of over 5.8 million in the USA, over 23 million worldwide, and continues to increase. ^{1, 2} The contractile dysfunction and arrhythmogenesis associated with HF is closely related to the remodeling of calcium handling, ³ which, in turn, is partially controlled by several signaling pathways in which Ca²⁺ has a prominent role. ⁴ Deriving a mechanistic understanding of alterations in calcium handling and calcium signaling is a critical step towards the development and improvement of physiology-based treatments for HF.

2. Overview of Cardiac Calcium Signaling

 ${\rm Ca^{2+}}$ plays a central part in regulating excitation-contraction (EC) coupling and in modulating systolic and diastolic function in the heart as shown in Figure 1. ${\rm Ca^{2+}}$ signal transduction in EC coupling comprises four steps. $^{4-6}$ Firstly, the trigger ${\rm Ca^{2+}}$ current ($I_{\rm Ca}$) is generated by the L-type ${\rm Ca^{2+}}$ channels expressed in the transverse tubules (T-tubules) and is initiated by membrane depolarization. Secondly, the ${\rm Ca^{2+}}$ ions diffuse across the narrow junctional zone to activate ryanodine receptors (RyR) and generate ${\rm Ca^{2+}}$ sparks, which considerably amplifies the original trigger ${\rm Ca^{2+}}$ signal. This process is known as ${\rm Ca^{2+}}$ -induced ${\rm Ca^{2+}}$ release (CICR). Thirdly, the ${\rm Ca^{2+}}$ efflux from the sarcoplasmic reticulum (SR) then diffuses in the cytoplasm to activate contraction by ${\rm Ca^{2+}}$ binding to troponin-C. Lastly, ${\rm Ca^{2+}}$ is transported back to the SR by SR ${\rm Ca^{2+-}ATPases}$ (SERCA) and out of cell via ${\rm Na^{+/}}$ ${\rm Ca^{2+}}$ exchangers (NCX). Abnormal handling of intracellular ${\rm Ca^{2+}}$ at any of these steps can cause cardiac dysfunction.

Intracellular Ca²⁺ homeostasis of cardiac myocytes is regulated by the phosphorylation and dephosphorylation of several key Ca²⁺-handling proteins. One important regulatory kinase is cAMP-dependant protein kinase (PKA), which has been shown to regulate L-type Ca²⁺ channels, RyR and phospholamban (PLN). Despite the fact that global PKA activity is not changed in the failing human heart,^{8, 9} its activity in the RyR macromolecular signaling complex might be locally increased.^{10, 11}

Another important regulatory kinase is the Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII). ¹² CAMKII is a protein kinase that modulates several intracellular Ca^{2+} -handling proteins such as RyR, PLN, L-type Ca^{2+} channels as well as Na^+ channels. ¹² CAMKII is associated directly with the RyR and modulates the activity of RyR. ^{13–15} Phosphorylation of PLN via CAMKII or PKA enhances the SR Ca^{2+} uptake via increased SERCA activity. The activity of CAMKII was shown to be significantly increased in the failing human heart and is correlated with the impaired ejection fraction. ^{9,16} Both PKA and CAMKII can be activated by β -adrenergic stimulation.

Finally, multiple isoforms of protein kinase C (PKC) might also play a role in regulating the Ca²⁺ handling. PKC- α is the dominant isoform of PKC in the human heart¹⁷ and its activity is triggered by the activation of $G_{\alpha q}$ coupled receptors (angiotensin II receptor, endothelin-1 receptor, and the α -adrenergic receptor).¹⁸ PKC- α can phosphorylate protein phosphatase inhibitor 1 (I-1), consequently increasing the activity of protein phosphatase 1 (PP1), and leading to dephosphorylation of PLN and thus decreasing the activity of SERCA.¹⁹ The level of PKC- α is increased in human HF.^{20–22} The role of other isoforms of PKC in regulating calcium handling remains to be elucidated.

3. Alteration in Intracellular Ca²⁺ and Functional Abnormalities in Human Heart Failure

3.1 Abnormal Ca2+ Handling and Mechanical Dysfunction in Human Heart Failure

The amount of Ca^{2+} delivered to the cytoplasm and the rate of Ca^{2+} removal from the cytoplasm are the two of the major factors determining the rate, intensity and duration of myocyte contraction.²³ Understanding of alterations in the intracellular Ca^{2+} concentration ($[Ca^{2+}]_i$) and their causal roles in contractile dysfunction in the failing human heart has been greatly advanced by the use of fluorescent $[Ca^{2+}]_i$ indicators,^{5, 24–32} which reflect changes in the free $[Ca^{2+}]_i$ necessary for the activation of contractile proteins.^{33, 34}

In isolated cells and tissues from failing human hearts, decreased amplitude of Ca^{2+} transient measured by the fluorescent intracellular Ca^{2+} indicators implies reduced Ca^{2+} release from $SR.^{28}, ^{30}, ^{35}$ This is correlated with decreased peak stretch amplitude, a measure of myocardial contraction. The reduced amplitude of Ca^{2+} transient is associated with decreased EC coupling gain 37 and decreased SR Ca^{2+} content. $^{30,38-42}$

Moreover, the Ca^{2+} transient from failing human hearts exhibits a reduced rate of Ca^{2+} removal. $^{24,\ 28,\ 30}$ The slowed rate of recovery of Ca^{2+} transient recovery is associated with a marked delay in tension relaxation in the failing human heart. 24 Finally, failing human hearts exhibits increased resting intracellular Ca^{2+} level, leading to diastolic dysfunction. $^{27,\ 28}$

Altered $[Ca^{2+}]_i$ is frequency-dependent and most obvious at high heart rate. ⁴³ Normally, the amplitude of the $[Ca^{2+}]_i$ transient is larger at higher stimulation frequencies. ³³ In human HF, however, the amplitude of Ca^{2+} transient was decreased at faster stimulation rates, leading to reduced tension development at higher frequencies. ^{27, 29} Increased resting $[Ca^{2+}]_i$ and a fusion of Ca^{2+} transient at fast frequencies may also occur, leading to an increase of end-

diastolic tension and a decrease of active tension generation associated with incomplete relaxation and twitch fusion. 27 The blunted or negative force frequency relationship (FFR) observed in both *in vivo* and *in vitro* studies of failing human hearts is in contrast with the positive FFR in non-failing human hearts, 27 , $^{44-46}$ and is associated with altered Ca²⁺ and Na⁺ homeostasis as well as an inability to increase the SR Ca²⁺ content at increasing stimulation frequencies. 39 , 42

The alteration of Ca²⁺ transient in failing human hearts is also region-dependent. We recently demonstrated the transmural hetergeneous remodeling of Ca²⁺ handing in the coronary-perfused left ventricular wedge preparations from failing and non-failing human hearts. 31 The sequence of Ca²⁺ transient relaxation is from epicardium to endocardium in both failing and non-failing human hearts at a slow heart rate (e.g., 0.67Hz/40 BPM) during endocardial pacing, because the difference of Ca²⁺ transient duration between subendocardium and subepicardium (or duration difference) is larger than the conduction time from subendocardium to subepicardium (Figure 2). Interestingly, this sequence is reversed at a fast heart rate (e.g., 1.67Hz/100 BPM) in the failing human heart due to a significant decrease of this duration difference (Figure 2C and 2D). In contrast, this sequence is not reversed in the non-failing human heart because the duration difference is not significantly changed at faster frequencies (Figure 2B and 2D). We hypothesize that this reversed sequence of relaxation at fast heart rates could contribute to the end-systolic dysfunction⁴⁴ observed in the failing human heart. The maintenance of the normal relaxation sequence at slow heart rates in the failing hearts provides another mechanism for the beneficial effects of the heart-rate reduction in the patients with HF.⁴⁷

3.2 Abnormal Ca²⁺ Handling and Arrhythmia in Human Heart Failure

While approximately fifty percent HF patients die from progressive pump failure, the remainder die suddenly, mostly due to tachyarrhythmias.⁴⁸ The relevance of arrhythmia to HF is evident from the significant survival benefit conferred by implantable cardioverter-defibrillators (ICD) on patients with advanced HF.⁴⁹ Among the multiple mechanisms contributing to the development of arrhythmia in HF patients,⁵⁰ changes in Ca²⁺ handling provide both triggers and substrate for the induction of arrhythmia.

Alterated Ca²⁺ handling in HF contributes to triggered activity arising from delayed after-depolarization (DAD) or early after-depolarization (EAD). DADs result from elevated intracellular Ca²⁺ load and spontaneous SR Ca²⁺ release, which leads to activation of transient inward current presumably carried by NCX.⁵¹ In a rabbit model of HF, enhanced NCX in HF increased the frequency of DADs.⁵¹ In a canine model of HF, increased RyR-mediated SR Ca²⁺ leak caused proarrhythmic DADs.⁵² EAD could occur in the setting of increasingly labile repolarization observed in HF,⁵³ possibly the result of synergistic activation of L-type Ca²⁺ current and NCX during phase 2 and phase 3 of the AP.⁵⁴ The cellular mechanisms of triggered activity have been reviewed elsewhere.⁵⁵

In contrast to the amount of work in animal models, our knowledge of triggered activity directly from human studies remains rather limited. Three-dimensional intraoperative mapping revealed that arrhythmias were maintained by focal mechanisms in end-stage HF patients with idiopathic dilated cardiomyopathy. ⁵⁶ However, whether these focal activations were triggered activities was not directly demonstrated in this study. Both DADs and EADs were not observed when the ventricular trabeculae of failing human hearts were perfused with normal Tyrodes' solution. Rather, DADs were only induced upon perfusion with a modified Tyrode's solution that mimicked the extracellular milieu of patients with severe HF. ⁵⁷ EADs were not observed in the modified solution. ⁵⁷ In contrast, EADs could be induced in 30% of failing rabbit trabeculae. ⁵⁷ This species difference signals the caution that

must be used when interpreting animal models and highlights the importance in conducting mechanistic studies directly in failing human hearts.

Abnormal Ca²⁺ handling may also produce the substrate for arrhythmia by facilitating AP duration alternans (APD alternans, or repolarization alternans),^{58, 59} the relevance of which was shown in the predictive power of T-wave alternans in predicting arrhythmic events in HF patients.⁶⁰ Spatially discordant APD alternans increases dispersion of repolarization and promotes unidirectional block and subsequent induction of arrhythmia.⁶¹ In a dog model of HF, Wilson et al.⁵⁹ showed that HF significantly lowered heart-rate threshold for calcium transient alternans. They also demonstrated that enhanced calcium transient alternans enhancement was independent of HF-associated changes in repolarization and appeared to be responsible for the enhanced APD alternans.⁵⁹ Finally, improvement of Ca²⁺ cycling through targeted SERCA2a gene expression was shown to retard the development of APD alternans.⁶² It remains to be demonstrated how the altered Ca²⁺ cycling leads to cellular alternans in the failing human heart.

4. Molecular and Cellular Basis of Abnormal Calcium Handling and Signaling in Human HF

Alteration in the $[Ca^{2+}]_i$ is attributed to the abnormal calcium handling in the EC coupling process, which is operated by the sarcolemma and SR, including L-type Ca^{2+} channels, RyR, NCX and SERCA2a. Altered EC coupling in HF have been reviewed in detail elsewhere. ^{3,18,43} Here we mainly focus on reviewing the results regarding the failing human heart.

4.1 Calcium-Induced Calcium Release (CICR)

Triggering of CICR (i.e., I_{Ca}) in the failing human heart is mostly unchanged, 26,28,30,63 though inhibition of I_{Ca} was observed at higher frequencies. 64 Thus, the smaller Ca^{2+} transient observed in HF is mainly due to a reduced capability of I_{Ca} to trigger Ca^{2+} release from the SR (or a reduced EC coupling gain).

The reduced EC coupling gain may result from the hyperphosphorylation of RyRs in the failing human heart. 10 PKA hyperphosphorylation of RyRs leads to the dissociation of the FKBP12.6 regulatory subunit, which inhibits the coupled gating of arrays of RyR channels and thus could result in a loss of EC coupling gain. 10 This is supported by the reduced amplitude and changed properties of Ca^{2+} sparks measured from isolated ventricular myocytes from failing human hearts. 65 More discussion of RyRs can be found in the section 4.2.

Rapid activation of RyRs by I_{Ca} is facilitated by the close proximity of the L-type Ca²⁺ channels and RyRs. Reduced EC coupling gain in HF could thus also originate from the geometric derangement of RyRs and L-type Ca²⁺ channels, as suggested by the spontaneous hypertensive rat with HF (SHR-HF).³⁷ Disorganization of T-tubules and a decrease in the colocalization of L-type Ca²⁺ channels and RyRs have been demonstrated in fixed ventricular samples from failing human hearts.⁶⁶ The actual loss of T-tubules in isolated myocytes from failing human heart was reported in one study ⁶⁷ but not in another,⁶⁸ findings that might be explained by the large spatial variations in T-tubule remodeling in human HF.⁶⁶

The ultrastructural defects in the T-tubule system were demonstrated to cause the dyssynchronous Ca^{2+} release (or defective EC coupling) by confocal line scanning techniques in the isolated ventricular myocytes from SHR-HF, with Ca^{2+} release being delayed in certain regions of a myocyte compared to the other normally coupled areas.⁶⁹

Louch and coworkers showed modest dyssynchrony of Ca²⁺ release in the isolated myocytes from failing human hearts. 70 While the local delayed SR Ca²⁺ release was confirmed in the whole heart level in SHR-HF, 71 no study has been done so far to demonstrate the dyssynchronous Ca²⁺ release at the tissue level from the failing human hearts. However, the morphological changes of Ca²⁺ transient observed in our recent left ventricular wedge preparations from failing human hearts imply the potentially important role of dyssynchronous Ca²⁺. ³¹ We observed two components in the rising portion of Ca²⁺ transient, with a slow rising component following an initial fast rising component (Figure 3A and 3B).³¹ It is possible that the first fast-rising component corresponds to the normally triggered Ca²⁺ release and the second slow-rising component corresponds to the delayed Ca²⁺ release, which has been shown in SHR-HF.⁷¹ Interestingly, this morphological change of Ca²⁺ was only observed at the subendocardium in 60% of the studied failing human hearts. ³¹ This regional difference might result from the higher susceptibility to ischemia by the endocardium compared with epicardium, ^{72,73} and suggests that the extrapolation of results from one region (e.g., epicardium) to another region (e.g., epicardium) in human studies should be done with caution. This delayed Ca²⁺ release might be also underlie the slower recovery of intracellular Ca²⁺ relative to the recovery of the action potential observed in our study (Figure 3A and 3C). That is, the Ca²⁺ transient outlasts the action potential at the subendocardium of the failing human heart, which might lead to phase three early afterdepolarization.⁷⁴

The reduced EC coupling gain could also result from decreased SR Ca^{2+} content, which has been extensively demonstrated in human HF. 30,38,39,41,42 The success of molecular therapies aimed at restoring SR Ca^{2+} content further underscores the importance of SR Ca^{2+} content. 75,76 Reduced SR Ca^{2+} content in the failing human heart could result from leaky RyRs, reduced SR Ca^{2+} uptake via SERCA2a, and increased Ca^{2+} extrusion via NCX, which are reviewed in the sections below (4.2-4.3).

4.2 RyR

Most studies showed no change in the protein expression of RyR in human HF. $^{3, 43}$ However, the characteristics of Ca $^{2+}$ sparks are altered in isolated myocytes from failing human hearts. 65 Furthermore, RyRs in human HF are "leaky," 10 contributing to a reduction of SR Ca $^{2+}$ content. 77 The SR Ca $^{2+}$ leak occurred despite reduced SR Ca $^{2+}$ loading in a canine model of HF. 77

Leaky RyR is thought to result from hyperphosphorylation of RyR by PKA or CAMKII. Increased Ca²⁺ sensitivity and higher probability of open RyR in failing human hearts was first observed by Marx and coworkers. ¹⁰ They concluded that the increased local PKA-mediated phosphorylation of RyR in HF leads to the disassociation of FKBP12.6 from RyRs, leading to higher open probability at rest. ¹⁰ They also observed decreased association of phosphatases (PP1, protein phosphatase 2A [PP2A]) to RyR, which may exacerbate PKA-mediated hyperphosporylation of RyR. ¹⁰ The hyperphosphoryation of RyRs could also result from the deficiency of phosphodiesterase 4D (PDE4D), which resides in the RyR macromolecular signaling complex and regulates the local concentration of cAMP that activates PKA. ¹¹ The impact of PKA phosphorylation in reducing the RyR/FKBP12.6 association remains controversial. ^{78,79}

Increased SR Ca²⁺ leak in isolated myocytes from failing rabbit hearts was shown to relate to the hyperphosphoryation of RyR by CAMKII. ⁸⁰ Ca²⁺ leak was reduced by the inhibition of CAMKII but not altered by PKA inhibition, ⁸⁰ suggesting the potential role of CAMKII inhibition in improving the Ca²⁺ handling in HF. This hypothesis is further supported by a recent study by Sossalla et al., ¹⁶ who showed a significant increase in the expression the CAMKII in both left and right ventricles of the failing human heart, and observed that

inhibition of CAMKII reduced the SR Ca²⁺ leak and increased the Ca²⁺ content. Importantly, they also showed that inhibition of CAMKII improved contractility in isolated ventricular trabeculae. ¹⁶ They reported that CAMKII inhibition restored the positive FFR. ¹⁶ This is in sharp contrast to the study by Kushiner et al., ⁸¹ which showed that CAMKII inhibition completely abolished the positive FFR in mouse heart. The result from the latter study is consistent with the hypothesis that CAMKII is responsible for sensing the frequency of Ca²⁺ oscillation ⁸² as well as for causing the positive FFR via increased phosphorylation of RyR and PLN at increasing frequencies. ^{13, 83} Kushiner and coworkers also showed that RyR phosphorylation by CAMKII was decreased in failing human hearts in despite of the global increase of CAMKII, ⁸¹ and suggested that the impaired RyR phosphorylation by CAMKII plays a role in blunted FFR in human HF. Further studies are needed to resolve these areas of controversy and clarify the molecular mechanism and the promise of CAMKII inhibition in improving the Ca²⁺ handling in human HF.

While much evidence supports altered regulation and function of the RyR leading to abnormal Ca^{2+} handling in failing human heart, there are studies indicating the opposite. Recordings of currents through the RyR from failing human hearts did not reveal any significant alterations at a single channel level. ⁸⁴ Jiang et al. observed neither structural nor functional change of RyRs from the failing human heart but did report a significant reduction in SERCA2a expression, suggesting that abnormal Ca^{2+} uptake may contribute more to the altered Ca^{2+} handling in human HF. ⁷⁸

4.3 SERCA2a, PLN and NCX

SR Ca²⁺ uptake was reduced in the failing human heart. ^{30, 46, 85, 86} This might be due to depressed protein expression of SERCA2a. Hasenfuss et al. observed downregulation of SERCA2a expression as well as a significant correlation between SERCA protein levels and SR Ca²⁺ uptake in failing human hearts. ⁴⁶ Overexpression of SERCA2a has been shown to restore the Ca²⁺ handling and the contractile function with positive FFR in isolated failing human myocytes. ^{87, 88} While some studies observed the downregulation of protein expression of SERCA2a in the failing human heart, others did not find any change in the protein expression of SERCA2a. ⁸⁹ This inconsistency might be explained by our recent findings. ³¹ We observed down-regulation of SERCA2a expression in samples from the subendocardium of failing human hearts with ischemic cardiomyopathy but not in samples from epicardium or from failing hearts with dilated cardiomyopathy (Figure 4A), suggesting that the alteration of SERCA2a expression might be region-dependent as well as HF etiology-dependent. ³¹

Besides the potential decrease in protein expression, the decreased activity of SERCA2a in HF might also result from altered regulation. This is supported by the findings that decreased SR $\rm Ca^{2+}$ uptake was observed in despite unchanged protein levels of SERCA2a. 85, 86, 90

SERCA2a is directly regulated by PLN which is mainly phosphorylated by PKA and CAMKII.⁸⁹ PLN inhibits SERCA2a activity when it is not phosphorylated, while its phosphorylated form disassociates from SERCA2a. In the failing human heart, majority of the studies indicate no change in the protein expression of PLN,^{31, 89} which is consistent with our recent study (Figure 4B).³¹ However, the phosphorylation state of PLN was decreased in the failing human heart,^{85, 90, 91} suggesting increased inhibition of SERCA2a by PLN in the failing human heart. PLN is mainly phosphorylated by PKA at serine-16 and by CAMKII at threonine-17. Phosphorylation at threonine-17 is decreased due to increased dephosphorylation by calcineurin in failing human hearts with dilated cardiomyopathy.⁹² PLN phosphorylation at serine-16 is decreased presumably due to increased level of PP1 in the failing human heart,^{90, 93} which might be a result of increased PKC-a.^{19, 20}

Interventions to attenuate the inhibitory effect of PLN on SERCA2a have been tested in animal models. Minamisawa et al. found that knockout of PLN significantly increased SR Ca²⁺ content and completely rescued the spectrum of heart-failure phenotype in a mouse model of HF. P4 Decreased PLN expression via adenoviral gene transfer of antisense of PLN was shown to improve both contraction and relaxation in isolated myocytes from failing human hearts. Inhibition of PKC- α was shown to increase the SR Ca²⁺ load and protect the mouse from HF. P5. The importance of PKC- α and other isoforms of PKC in the Ca²⁺ handling in human HF remains to be determined.

While protein expression of NCX was found upregulated in most animal models of HF,³ it is less consistent in the failing human heart, with most studies finding either increased or unchanged protein expression of NCX.⁴³ In contrast to reduced SR Ca²⁺ uptake, the NCX current density as a function of $[Ca^{2+}]$ was not changed in the failing human heart.³⁰ However, the contribution of NCX to the $[Ca^{2+}]_i$ relaxation was increased due to the depressed SR Ca²⁺ uptake.³⁰ Furthermore, the preference of NCX current direction during the action potential plateau shifted from inward direction $(Ca^{2+}$ efflux) to outward direction $(Ca^{2+}$ influx) due to a reduced submembrane $[Ca^{2+}]_i$ and increased $[Na^+]_i$ in the failing human heart.⁹⁶ The reversed-mode NCX during AP plateau could contribute to a slow decay of $[Ca^{2+}]_i$ transient,^{41, 96} which may facilitate contraction at slow heart rates but may also lead to diastolic dysfunction at faster heart rates.⁴²

4.4 Loss of metabolic capacity

 Ca^{2+} handling and energy homeostasis are interdependent. 97 Ca^{2+} homeostasis relies on efficient energy-driven ionic fluxes, i.e., through SERCA2a and Na⁺-K⁺ ATPase, while $[Ca^{2+}]_i$ in turn determines energy consumption through contraction and Ca^{2+} transport as well as energy production via the regulation of ATP generation in mitochondria. $^{97,\,98}$ Disturbance of the finely tuned balance between the two could be responsible for the abnormal Ca^{2+} handling and diminished contractility that are hallmarks of HF.

HF is associated with defects in energy metabolism, with decreased energy production as well as impaired energy transfer and utilization. These impaired cardiac energetics may represent the thermodynamic limit for Ca²⁺ handling. Reduced local ATP/ADP ratio, due to a local lack of creatinine kinase, could affect the kinetic and thermodynamic efficiency of SERCA in HF, Providing another mechanism for impaired SR Ca²⁺ uptake. Indeed, ATP was reported to protect SERCA2a from being denatured by hydroxyl radicals, in implying that energy starvation might render SERCA2a unprotected from increased oxidative stress in human HF.

Improving the myocardial energetics has been shown to normalize the Ca^{2+} cycling in isolated failing human myocytes. 101 β -blockers, which decrease the energy demand and thus ameliorate the mismatch between energy production and consumption, has been shown to normalize the function and regulation of key Ca^{2+} handling proteins in failing human hearts. 102 Similarly, left ventricular assist devices (LVADs), which unload the heart and support the circulation, impart improved Ca^{2+} handling in human HF. $^{10,\ 103}$ Finally, hemodynamic improvement by cardiac resynchronization therapy (CRT) is correlated with improved Ca^{2+} handling in the subset of HF patients who respond to this therapy. 104 On the other hand, restoration of Ca^{2+} homeostasis may result in improved cardiac energetics. 105

5. Correcting Abnormal Calcium Handling in HF

While the causes of HF may differ, there is a common theme underlying the progression from normal to failing heart. An initial cardiac insult, which, in the United States is most commonly inadequate myocardial flow (myocardial infarction), prolonged pressure overload

(hypertension), or abnormal flow through the heart valves (valvular stenosis or insufficiency) causes the heart to alter the shape of its principal pumping chamber, the left ventricle, and the surrounding organs to activate multiple hormonal systems in an attempt to maintain cardiac output. These compensatory changes are initially helpful in sustaining adequate cardiac output and blood pressure. 106 Over time, however, these same changes become maladaptive. The remodeled ventricle becomes increasingly dilated and hypertrophied, resulting in suboptimal pump geometry. In parallel, continuous activation of the sympathetic nervous system and the renin-angiotensin-aldosterone (RAA) axis 107, 108 leads to increased oxygen consumption, increased metabolism and molecular changes that ultimately impair the contractile function of myocytes. In this manner, an initial cardiac insult initiates a cascade of events leading to reduced ejection fraction, reduced myocardial contractility and poor cardiac output. Current clinical therapies aim to halt or reverse these maladaptive events. Myocardial revascularization by coronary artery bypass grafting (CABG) and percutaneous coronary intervention (PCI) restores myocardial blood flow to ensure that adequate oxygen and metabolites are supplied to the myocardium. On the other hand, most medical therapies aim to block the deleterious effects of prolonged hormonal stimulation and the subsequent molecular changes that ultimately impair myocardial contractility and promote fatal cardiac arrhythmias. Importantly, reduced cardiac contractility and diminished contractile reserve are hallmarks of HF, and central to these pathologies is defective intracellular Ca²⁺ handling.³⁹ Current and future medical strategies to restore the function of the key calcium handling molecules are discussed below.

5.1 Molecular Changes Due to Activation of the Sympathetic Nervous System

After an initial cardiac insult, the sympathetic nervous system is activated to maintain adequate cardiac output. This stress response, also known as the "fight or flight" response, is highly conserved evolutionarily and crucial to increasing cardiac stroke volume and heart rate, ¹⁰⁹ the product of which is cardiac output. Catecholamines such as epinephrine and norepinephrine (adrenaline and noradrenaline, respectively) are released into the bloodstream where they effect changes on target organs, including the heart.

In response to sympathetic stimulation, both heart rate and myocyte contractility are increased, the latter of which is directly related to changes in Ca^{2+} handling. Epinephrine and norepinephrine bind to the β -adrenoreceptor, leading to activation of several signaling pathways that ultimately increase the amount of intracellular Ca^{2+} released by the RyR per amount of trigger Ca^{2+} entering the cell through L-type Ca^{2+} channels, thereby increasing EC coupling gain. 37 β -adrenoreceptor binding also increases contractile force and allows more rapid release and reuptake of Ca^{2+} , allowing more time for diastolic filling, increasing stroke volume and therefore increasing cardiac output. 109 , 110

Downstream effectors activated through these pathways include PKA via G protein activation of adenylate cyclase 106 and increased CAMKII activity, 13 while activation of both the RAA axis or β_1 -adrenoreceptor leads to activation of PKC. $^{19},\,^{111}$ PKA and CAMKII phosphorylate L-type Ca^{2^+} channels, RyR and PLN. Many of these signaling events inherent to the "fight or flight" response are seen physiologically and in fact, are crucial for survival during isolated periods of stress or exercise. However, the continuous activation of these pathways that occurs in HF ultimately leads to defective Ca^{2^+} handling and diminished contractility, $^{112},\,^{113}$ and are therefore attractive targets of current and future medical therapies.

Chronic activation of G protein coupled β -adrenoreceptors by sympathetic hormones leads to several maladaptive changes including decreased expression and function of adenylate cyclase, increased expression of inhibitory proteins G protein G_i and β -adrenoreceptor kinase, and even decreased expression and coupling of the β -adrenoreceptor

itself. $^{106,\ 114-116}$ Downstream of the β -adrenoreceptor, additional changes occur. PKA hyperphosphorylates L-type Ca^{2+} channels, RyR and NCX, while simultaneously, the protein levels of L-type Ca^{2+} channels, RyR and SERCA2a are altered. $^{10,\ 41,\ 117,\ 118}$ In summary, chronic β -adrenoreceptor activation induces numerous changes that alter the function of proteins critical to Ca^{2+} handling. Determining which of these changes are causal for the diminished cardiac contractility seen in HF has proven far more challenging. Nonetheless, a number of pathways and targets have been elucidated.

5.2 Combating Chronic β-adrenoreceptor Stimulation

Blockade of β -adrenoreceptors using a class of drugs known as β -blockers restores cardiac function and significantly increase the survival of patients with HF. 119-121122, 123 The effectiveness of this therapy is somewhat counterintuitive since β-blockers depress contractility and heart rate, two major determinants of cardiac output. Indeed, physicians treating acute HF often temporarily discontinue β-blockers to improve cardiac output and restore fluid balance. After the acute volume overload has been corrected, however, treatment with β -blockers is reinstituted. The reasoning behind this strategy is that chronic activation of β-adrenoreceptors is ultimately maladaptive to the heart, and leads to the progression of HF. β-blockers combat this chronic sympathetic stimulation. Mechanistically, they decrease intracellular cAMP concentration and thus decrease the activity of PKA, restoring physiologic function and expression levels of downstream effectors. $\hat{102}$, 124, 125, 118, 126–128 Treatment with β -blockers also reverses the hyperphosphorylation of RyR to improve binding of FKBP12.6, thereby restoring physiologic RyR function. ^{102, 125, 128, 129} Diminished SERCA2a protein levels are also restored by $\beta\text{-blocker}$ therapy, allowing appropriate reuptake of Ca^{2+} to allow adequate diastolic function and filling time. 129 Thus, β -blocker therapy may halt or reverse disease progression and reduce mortality by reversing the diminished Ca²⁺ transient amplitude in systole, while simultaneously reversing increased intracellular Ca²⁺ concentration and slowed rate of Ca²⁺ transient decay in diastole. ^{28, 130} It is important to point out that despite numerous studies showing improvement in HF with the use of β-blockers, the precise mechanisms responsible for their well established beneficial effects remain controversial. And while effects on Ca²⁺ handling have been demonstrated, improved organ level structural changes have also been observed. 106

5.3 Inhibiting the Effect of Angiotensin II

In addition to the sympathetic nervous system, another hormonal axis is chronically elevated in patients with HF. The renin-angiotensin-aldosterone (RAA) axis is central to maintaining blood pressure. In response to low blood pressure, renin is released from the kidney and converts the pre-hormone angiotensinogen to angiotensin I. Angiotensin I is converted to Angiotensin II (ATII) in the lungs by angiotensin converting enzyme (ACE). ATII potently increase blood pressure by binding ATI receptors in the vasculature and causing constriction of blood vessels. ATI receptors are also located on myocytes, where they are acted upon by ATII. Two major categories of drugs designed to lower blood pressure by inhibiting this pathway are angiotensin converting enzyme inhibitors (ACEI), which block the conversion of ATI to ATII, and angiotensin receptor blockers (ARB), which inhibit the binding of ATII to ATI-receptors.

HF is a state of low cardiac output, which results in diminished blood pressure. However, low blood pressure is typically seen only at the last stages of HF. This is likely due to the maintenance of blood pressure by chronically elevated levels of ATII seen in HF patients. $^{107,\ 108}$ Importantly, the use of ACEIs and ARBs has been shown to further significantly reduce mortality and fatal cardiac arrhythmias in patients with HF. 122 Hence, in addition to β -blockers, ACEIs or ARBs are recommended first line agents for the treatment

of HF. ¹⁰⁸ ACEIs and ARBs likely reduce mortality by lowering blood pressure and reducing myocardial fibrosis in patients with HF. More recently, however, they have been shown to alter signaling cascades that are involved in Ca²⁺ handling.

ATII binding to ATI receptors on myocytes activates the G protein Gq, which has multiple downstream effectors that impact the progression of HF. One of these effectors is PLC, which, in turn, activates PKC α , the predominant cardiac PKC isoform. ¹³¹ PKC- α activates I-1, which increases the activity of PP1 to dephosphorylate PLN. ¹⁹ In a mouse model of diastolic HF, ACE was overexpressed to increase levels of ATII, resulting in dephosphorylation of PLN and diastolic HF. ¹³² Importantly, dephosphorylated PLN inhibits the ability of SERCA2a to pump Ca²⁺ back into the SR, which may impair heart function by increasing the Ca²⁺ transient duration and decreasing the SR Ca²⁺ content. This alteration likely exacerbates the reduced diastolic filling time, stroke volume and contractility that are seen in heart failure.

ATII mediated Gq activation also causes activation of the downstream effectors calcineurin 133, 134 and mitogen-activated protein (MAP) kinases such as extracellular signal regulated kinases 1 and 2 (ERK1/2), 118, 135 leading to pathological hypertrophic ventricular remodeling. Calcineurin, a Ca²⁺-dependent phosphatase that regulates hypertrophic gene transcription by desphosphorylating transcription factors such as nuclear factor of activated T-cells (NFAT), has been identified as a key enzyme involved in the induction of pathological cardiac hypertrophy. NFAT activity was upregulated in mouse models of pressure overload and HF but not in mice with exercise-induced physiological hypertrophy. Supporting this notion, inhibition of calcineurin prevented cardiac hypertrophy in rodent models of cardiomyopathy and pressure overload. ACEIs and ARBs may thus impart some of their beneficial effects on failing hearts by reversing maladaptive effects of ATII on Ca²⁺ handling, which, in turn, may also ameliorate ventricular remodeling via the Ca²⁺ dependent calcineurin pathway. Interestingly, all of these effects appear to occur through activation of major phosphatase, rather than kinase effectors.

Patients with HF exhibit elevated levels of reactive oxygen species (ROS). ATII has been shown to be a key player in producing destructive ROS that are involved in the progression of HF. 139, 140 ATII induces the production of ROS in the heart, 141 leading to damage to cellular components that are critical to calcium handling in myocytes such as the outer and inner membranes of organelles such as the SR and protein involved in Ca²⁺ signaling and handling. 142–145 ROS alter the function of NCX, decrease L-type Ca²⁺ channel currents and depress the activity of SERCA2a. 146–148 Thus, blockade of ATII through ACEIs and ARBs provides clinicians a tool to reduce oxidative stress in failing myocytes and move closer to the goal of restoring calcium handling.

Interestingly, blocking ATII may also directly inhibit β -adrenoreceptor overstimulation. The ARB valsartan binds presynaptic ATI receptors to inhibit the release of norepinephrine while also stimulating its reuptake, ¹⁴⁹ thereby reducing the amount of norepinephrine released into the bloodstream. Whether this holds true for other ARBs is unclear.

5.4 Calcium Channel Blockade

One of the disappointments of clinicians and researchers alike is the lack of benefit and even detrimental effect of drugs that block L-type Ca²⁺ channels (dihydropyridine receptors). These drugs, used primarily to treat blood pressure, have dilatory effects on blood vessels inside and outside the heart. The vasodilatory action of Ca²⁺ channel blockers was theorized by many to improve cardiac performance and reduce myocardial ischemia. However, these

theoretical advantages have not been translated into clinical benefits in controlled clinical HF trials, discussed in more detail below.

Ca²⁺ channel blocking drugs have not improved symptoms of HF or enhanced exercise tolerance, and signficantly, both short- and long-term treatment with these drugs have increased the risks of worsening HF and death. $^{150-156}$ L-type Ca²⁺ channel blockade is also ineffective in mild HF, 157 and non-dihydropyridine type Ca²⁺ antagonists, including T-type Ca²⁺ channel blockers, have also proven ineffective. $^{158,\ 159}$ Moreover, Ca²⁺ channel blockers can have life threatening interactions with β -blockers and ACEIs, by inducing severe bradycardia (low heart rate) and hypotension (low blood pressure). $^{160,\ 161}$ As a result, most Ca²⁺ channel blockers are avoided in patients with HF, even when used for the treatment of chest pain or hypertension. Of available agents, only the dihydropyridine drug amlodipine has been shown not to adversely affect survival, although experience with the drug exists largely in patients who are not taking β -blockers. 162 In fact, current clinical HF guidelines state that Ca²⁺ channel blocking drugs are contraindicated in the treatment for patients with current or prior symptoms of HF and reduced left ventricular ejection fraction. 108

5.5 Experimental Therapies Targeting Abnormal Calcium Handling

The limited medical therapies available for the treatment of HF and the significant abnormalities in Ca^{2+} handling observed in failing hearts has prompted significant interest in developing therapies that directly target and correct these altered molecular pathways. A subset of some promising therapies and strategies are discussed below.

- **5.5.1. Sensitizing β-adrenoreceptor Function**—Restoring sensitization of β-adrenoreceptors via new agents represents an attractive drug therapy. The desensitization of β-adrenoreceptors is mediated in part by G-protein coupled receptor kinase (GRK, also known as β-ARK). Activation of GRK desensitizes β-adrenoreceptors in the heart. ¹⁶³ In contrast, inhibition of GRK with concomitant β-blocker therapy improves survival in a mouse model of HF. ^{164, 165} Currently, no drug has been developed to inhibit GRK pharmacologically, but a dominant-negative GRK expressed in a mouse model of HF has been shown to prevent disease progression. ¹⁶⁵ In the future, such a multi-pronged approach to combat chronic β-adrenoreceptor stimulation may provide significant further benefit in the treatment of HF.
- **5.5.2.** Improving RyR Function—Hyperphosphorylation of RyR by PKA occurs due to chronic β -adrenergic stimulation and leads to Ca²⁺ leak through the RyR, ¹⁰¹⁶⁶ which contributes to contractile dysfunction and fatal ventricular arrhythmias. ^{109, 167, 168} These arrhythmias are a major cause of death in patients with advanced HF. ¹⁰⁸ This knowledge has prompted efforts to design a small molecule that will improve RyR function. JTV519 is an experimental drug that enhances the binding of FKBP12.6 to RyR and reduces Ca²⁺ leak from the SR. Encouragingly, this agent has been shown to suppress ventricular arrhythmias in a mouse model ¹⁶⁸ and improve contractile function in a canine model of HF. ¹⁶⁹ Finally, Inhibition of CAMKII, which also hyperphosphorylates RyR, represents another strategy of reducing the RyR Ca²⁺ leak seen in HF. Chemical or genetic inhibition of CAMKII prevented cardiac remodeling in a murine model of isoproterenol-induced cardiomyopathy. ¹⁷⁰
- **5.5.3 Enhancing SERCA2a Reuptake of Calcium**—As discussed above, diminished Ca²⁺ reuptake in failing hearts is likely due to a combination of decreased levels of SERCA2a expression^{4329, 17131} and hypophosphorylation of PLN. ^{172, 173} Understandably,

methods to increase SERCA2a expression and PLN phosphorylation are being actively pursued by investigators.

Attempts to overexpress SERCA2a in humans with advanced HF are currently underway.⁷⁶ In this study, a SERCA2a expression vector packaged in an Adenovirus-Associated Virus (AAV) envelope is administered by direct intracoronary injection.⁷⁶ Of the 9 patients treated, 5 demonstrated improvements from baseline to month six across a number of parameters important in HF, including symptoms, 6-minute walk test, oxygen consumption, and ejection fraction.⁷⁶ Notably, 2 patients who failed to improve had preexisting anti-AAV neutralizing antibodies, underscoring a limitation of such an approach.⁷⁶

Selectively enhancing PLN phosphorylation is another avenue being actively pursued. One such line of experiments involves the delivery of a pseudo-phosphorylated mutant of PLN into failing hearts using a viral vector. Interestingly, this mutant was demonstrated to suppress HF progression in hamsters ¹⁷⁴ and post-myocardial infarction rats, ¹⁷⁵ and reversed HF in sheep with chronic pacing induced HF. ¹⁷⁶ Decreasing the activity of PP1 (which dephosphorylates PLN) is another possible approach. Inhibitor-2 (INH-2) is an endogenous phosphatase that inhibits PP1 that selectively decreases SR microvesicle-associated PP1 activity. Moreover, gene delivery of INH-2 was shown to increase PLN phosphorylation and increase survival in a hamster model HF. ¹⁷⁷ Despite these encouraging studies, no small molecules that can selectively increase PLN phosphorylation have been developed to date.

5.5.4 Retarding ROS Effects—Established experimental studies showing that ROS are detrimental have prompted efforts to combat oxidative damage in patients with HF. The β -blocker carvedilol has been proven effective in reducing the progression of HF, and notably, may exert some of its beneficial effects through anti-oxidant activity. ¹¹⁹ Xanthine oxidase (X0), an enzyme involved in the synthesis of DNA precursors, has been shown to generate ROS, and has attracted the interest of researchers as a molecule to target with inhibitors. ^{178, 179} Encouragingly, inhibition of XO with the drug allopurinol improved myocardial efficiency in patients with HF, ¹⁸⁰ and prompted the initiation of a large-scale clinical trial testing its safety and efficacy in patients with advanced HF. ¹⁸¹ While this study did not show a reduction in morbidity or mortality overall, post-hoc analysis revealed benefit in a subset of patients with elevated levels of the XO product uric acid. ¹⁸² And while it will be helpful to determine the specific changes in Ca²⁺ handling pathways that were affected in this subset of patient, this study also suggests that more robust improvements in HF may be elicited in future trials by more specifically targeting the subset of HF patients with elevated ROS activity.

5.6 Calcium Abnormalities in Human Ischemic versus Non-ischemic Cardiomyopathy

End-stage HF that is the result of myocardial infarction is known as ischemic cardiomyopathy (ICM). HF in patients with no history of myocardial infarction is referred to as non-ischemic cardiomyopathy (NICM). Interestingly, distinct abnormalities in calcium handling have been demonstrated depending upon the etiology of HF. For example, mRNA expression of RyR is significantly decreased in the hearts of patients with end-stage ICM, but increased (albeit nonsignificantly) in patients with NICM. ¹⁸³ Similarly, ICM is associated with a decreased rate of calcium uptake into the SR, while NICM is associated with a decreased rate calcium release from the SR. ¹⁸⁴ The authors of the latter study concluded that abnormal SR calcium uptake may explain the contractile dysfunction seen in ICM, while abnormal SR calcium release may be the primary disruption in NICM. Thus, different insults leading to cardiomyopathy may do so by disrupting calcium handling via different mechanisms.

While different etiologies of HF exhibit different profiles of calcium handling abnormalities, there are no therapies to date that have been shown to improve calcium handling in an etiology-specific manner. Recently, Sossalla and colleagues demonstrated that inhibition of CaMKII using two different small molecules significantly increased contractility of isolated preparations of human failing heart tissue. ¹⁶ The increase in contractility was equal in HF of ischemic and non-ischemic etiologies. These studies suggest that in addition to the need to use human tissue to analyze signaling pathways and small molecules for improved calcium handling, separating observations with respect to etiology will be important.

6. Conclusion

It has been well recognized that abnormal Ca²⁺ handling is a key pathophysiological mechanism in human HF. However, our understanding of the underlying molecular and cellular mechanisms for the altered calcium handling in the failing human heart remains incomplete.

This is partly due to the complexity of the system, which involves the interplay between a number of signaling pathways that regulates the Ca²⁺ homeostasis at different time scales. ^{75, 185} That is, while interrupting or augmenting one of pathways in the cascade might lead to expected beneficial therapeutic effects, it might also produce unexpected deleterious effects. ⁷⁵ Nevertheless, the overall structure of this complex system is continually being revealed by ongoing basic and clinical research, which carries the hope of facilitating the development of effective diagnostic and treatment modalities for HF. The progress is also slowed by limited data from human studies. While many mechanistic hypothesis and potential therapeutic intervention for the abnormal Ca²⁺ handling in HF are being proposed and tested in animal models of HF, the examination of these hypothesis and therapies using functional studies of isolated cells or tissues from the failing human heart are rather limited. Basic understanding and clinical translation can be greatly facilitated by testing these hypotheses in explanted human heart and human heart tissue donated for research by patients. ¹⁸⁶

Gaining a clearer understanding of the causative mechanisms of abnormal Ca²⁺ handling is crucial to developing promising new therapies to treat HF. Despite our best efforts, there are currently only two major medical pharmacological approaches available to the clinician for the treatment of patients with HF: blockade of the β-adrenoreceptor and inhibition of the RAA axis. These therapies are used to treat non-ischemic (the majority of which are idiopathic), ischemic and valvular cardiomyopathies, even though we recognize fundamental differences in the insults that cause these separate conditions. Such blanket approaches demonstrate the limits of our current knowledge, and the need for further observation and testing before new therapies can be delivered to the patient. Moreover, it is clear that many pathways involving Ca²⁺ handling converge on and act through a few key molecules. Thus, the complex biological processes leading to HF must be further dissected with respect to specific isoforms, subcellular locations and etiology of HF. Similarly, it is important to realize that individual drugs effects must be categorized based on the species and type of animal model used. Finally, we must recognize that the road to developing a human therapeutic agent, i.e., going from the bench to the bedside, is a time consuming and expensive one, and littered with failures. These complexities may explain why after years of research, the clinical armamentarium for reversing HF remains rather limited. Despite these drawbacks, it is encouraging that many promising new therapies to ameliorate abnormal calcium handling are visible on the horizon, based on findings in animal models of HF. Increased research on functional human heart tissue would facilitate translation of these findings to the clinic.

References

1. Lloyd-Jones D, Adams RJ, Brown TM, Carnethon M, Dai S, De Simone G, Ferguson TB, Ford E, Furie K, Gillespie C. Heart disease and stroke statistics--2010 update: a report from the American Heart Association. Circulation. 2010; 121(7):e46. [PubMed: 20019324]

- 2. Bui AL, Horwich TB, Fonarow GC. Epidemiology and risk profile of heart failure. Nat Rev Cardiol. 2011; 8(1):30–41. [PubMed: 21060326]
- Bers, DM. Excitation-contraction coupling and cardiac contractile force. Springer Netherlands: 2001.
- 4. Berridge MJ, Bootman MD, Roderick HL. Calcium signalling: dynamics, homeostasis and remodelling. Nat Rev Mol Cell Biol. 2003; 4(7):517–529. [PubMed: 12838335]
- 5. Morgan JP. Abnormal intracellular modulation of calcium as a major cause of cardiac contractile dysfunction. N Engl J Med. 1991; 325(9):625–632. [PubMed: 1861696]
- Hasenfuss G, Pieske B, Holubarsch C, Alpert NR, Just H. Excitation-contraction coupling and contractile protein function in failing and nonfailing human myocardium. Adv Exp Med Biol. 1993; 346:91–100. [PubMed: 8184784]
- 7. Katz AM. Cardiomyopathy of overload. A major determinant of prognosis in congestive heart failure. N Engl J Med. 1990; 322(2):100–110. [PubMed: 2403651]
- 8. Bohm M, Reiger B, Schwinger RH, Erdmann E. cAMP concentrations, cAMP dependent protein kinase activity, and phospholamban in non-failing and failing myocardium. Cardiovasc Res. 1994; 28(11):1713–1719. [PubMed: 7842467]
- 9. Kirchhefer U, Schmitz W, Scholz H, Neumann J. Activity of cAMP-dependent protein kinase and Ca2+/calmodulin-dependent protein kinase in failing and nonfailing human hearts. Cardiovasc Res. 1999; 42(1):254–261. [PubMed: 10435018]
- Marx SO, Reiken S, Hisamatsu Y, Jayaraman T, Burkhoff D, Rosemblit N, Marks AR. PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. Cell. 2000; 101(4):365–376. [PubMed: 10830164]
- Lehnart SE, Wehrens XH, Reiken S, Warrier S, Belevych AE, Harvey RD, Richter W, Jin SL, Conti M, Marks AR. Phosphodiesterase 4D deficiency in the ryanodine-receptor complex promotes heart failure and arrhythmias. Cell. 2005; 123(1):25–35. [PubMed: 16213210]
- 12. Maier LS, Bers DM. Role of Ca2+/calmodulin-dependent protein kinase (CaMK) in excitation-contraction coupling in the heart. Cardiovasc Res. 2007; 73(4):631–640. [PubMed: 17157285]
- 13. Wehrens XH, Lehnart SE, Reiken SR, Marks AR. Ca2+/calmodulin-dependent protein kinase II phosphorylation regulates the cardiac ryanodine receptor. Circ Res. 2004; 94(6):e61–e70. [PubMed: 15016728]
- 14. Hain J, Onoue H, Mayrleitner M, Fleischer S, Schindler H. Phosphorylation modulates the function of the calcium release channel of sarcoplasmic reticulum from cardiac muscle. J Biol Chem. 1995; 270(5):2074–2081. [PubMed: 7836435]
- Currie S, Loughrey CM, Craig MA, Smith GL. Calcium/calmodulin-dependent protein kinase IIdelta associates with the ryanodine receptor complex and regulates channel function in rabbit heart. Biochem J. 2004; 377(Pt 2):357–366. [PubMed: 14556649]
- Sossalla S, Fluschnik N, Schotola H, Ort KR, Neef S, Schulte T, Wittkopper K, Renner A, Schmitto JD, Gummert J, El-Armouche A, Hasenfuss G, Maier LS. Inhibition of elevated Ca2+/ calmodulin-dependent protein kinase II improves contractility in human failing myocardium. Circ Res. 2010; 107(9):1150–1161. [PubMed: 20814023]
- 17. Hambleton M, Hahn H, Pleger ST, Kuhn MC, Klevitsky R, Carr AN, Kimball TF, Hewett TE, Dorn GW 2nd, Koch WJ, Molkentin JD. Pharmacological- and gene therapy-based inhibition of protein kinase Calpha/beta enhances cardiac contractility and attenuates heart failure. Circulation. 2006; 114(6):574–582. [PubMed: 16880328]
- 18. Yano M, Ikeda Y, Matsuzaki M. Altered intracellular Ca2+ handling in heart failure. J Clin Invest. 2005; 115(3):556–564. [PubMed: 15765137]
- Braz JC, Gregory K, Pathak A, Zhao W, Sahin B, Klevitsky R, Kimball TF, Lorenz JN, Nairn AC, Liggett SB, Bodi I, Wang S, Schwartz A, Lakatta EG, DePaoli-Roach AA, Robbins J, Hewett TE,

- Bibb JA, Westfall MV, Kranias EG, Molkentin JD. PKC-alpha regulates cardiac contractility and propensity toward heart failure. Nat Med. 2004; 10(3):248–254. [PubMed: 14966518]
- 20. Bowling N, Walsh RA, Song G, Estridge T, Sandusky GE, Fouts RL, Mintze K, Pickard T, Roden R, Bristow MR, Sabbah HN, Mizrahi JL, Gromo G, King GL, Vlahos CJ. Increased protein kinase C activity and expression of Ca2+-sensitive isoforms in the failing human heart. Circulation. 1999; 99(3):384–391. [PubMed: 9918525]
- Wang J, Liu X, Sentex E, Takeda N, Dhalla NS. Increased expression of protein kinase C isoforms in heart failure due to myocardial infarction. Am J Physiol Heart Circ Physiol. 2003; 284(6):H2277–H2287. [PubMed: 12742831]
- 22. Simonis G, Briem SK, Schoen SP, Bock M, Marquetant R, Strasser RH. Protein kinase C in the human heart: differential regulation of the isoforms in aortic stenosis or dilated cardiomyopathy. Mol Cell Biochem. 2007; 305(1–2):103–111. [PubMed: 17594058]
- 23. Yue DT, Marban E, Wier WG. Relationship between force and intracellular [Ca2+] in tetanized mammalian heart muscle. J Gen Physiol. 1986; 87(2):223–242. [PubMed: 2419483]
- 24. Gwathmey JK, Copelas L, MacKinnon R, Schoen FJ, Feldman MD, Grossman W, Morgan JP. Abnormal intracellular calcium handling in myocardium from patients with end-stage heart failure. Circ Res. 1987; 61(1):70–76. [PubMed: 3608112]
- 25. Gwathmey JK, Slawsky MT, Hajjar RJ, Briggs GM, Morgan JP. Role of intracellular calcium handling in force-interval relationships of human ventricular myocardium. J Clin Invest. 1990; 85(5):1599–1613. [PubMed: 2332508]
- Beuckelmann DJ, Nabauer M, Erdmann E. Characteristics of calcium-current in isolated human ventricular myocytes from patients with terminal heart failure. J Mol Cell Cardiol. 1991; 23(8): 929–937. [PubMed: 1658345]
- 27. Gwathmey JK, Warren SE, Briggs GM, Copelas L, Feldman MD, Phillips PJ, Callahan M Jr, Schoen FJ, Grossman W, Morgan JP. Diastolic dysfunction in hypertrophic cardiomyopathy. Effect on active force generation during systole. J Clin Invest. 1991; 87(3):1023–1031. [PubMed: 1999483]
- 28. Beuckelmann DJ, Nabauer M, Erdmann E. Intracellular calcium handling in isolated ventricular myocytes from patients with terminal heart failure. Circulation. 1992; 85(3):1046–1055. [PubMed: 1311223]
- Pieske B, Kretschmann B, Meyer M, Holubarsch C, Weirich J, Posival H, Minami K, Just H, Hasenfuss G. Alterations in intracellular calcium handling associated with the inverse force-frequency relation in human dilated cardiomyopathy. Circulation. 1995; 92(5):1169–1178.
 [PubMed: 7648662]
- 30. Piacentino V 3rd, Weber CR, Chen X, Weisser-Thomas J, Margulies KB, Bers DM, Houser SR. Cellular basis of abnormal calcium transients of failing human ventricular myocytes. Circ Res. 2003; 92(6):651–658. [PubMed: 12600875]
- 31. Lou Q, Fedorov VV, Glukhov AV, Moazami N, Fast VG, Efimov IR. Transmural heterogeneity and remodeling of ventricular excitation-contraction coupling in human heart failure. Circulation. 2011; 123:1881–1890. [PubMed: 21502574]
- 32. Vahl CF, Bonz A, Timek T, Hagl S. Intracellular calcium transient of working human myocardium of seven patients transplanted for congestive heart failure. Circ Res. 1994; 74(5):952–958. [PubMed: 8156642]
- 33. Allen DG, Blinks JR. Calcium transients in aequorin-injected frog cardiac muscle. Nature. 1978; 273(5663):509–513. [PubMed: 307184]
- 34. Blinks JR, Wier WG, Hess P, Prendergast FG. Measurement of Ca2+ concentrations in living cells. Prog Biophys Mol Biol. 1982; 40(1–2):1–114. [PubMed: 6758036]
- 35. Kubo H, Margulies KB, Piacentino V 3rd, Gaughan JP, Houser SR. Patients with end-stage congestive heart failure treated with beta-adrenergic receptor antagonists have improved ventricular myocyte calcium regulatory protein abundance. Circulation. 2001; 104(9):1012–1018. [PubMed: 11524394]
- 36. Hasenfuss G, Mulieri LA, Leavitt BJ, Allen PD, Haeberle JR, Alpert NR. Alteration of contractile function and excitation-contraction coupling in dilated cardiomyopathy. Circ Res. 1992; 70(6): 1225–1232. [PubMed: 1576741]

37. Gomez AM, Valdivia HH, Cheng H, Lederer MR, Santana LF, Cannell MB, McCune SA, Altschuld RA, Lederer WJ. Defective excitation-contraction coupling in experimental cardiac hypertrophy and heart failure. Science. 1997; 276(5313):800–806. [PubMed: 9115206]

- 38. Lindner M, Erdmann E, Beuckelmann DJ. Calcium content of the sarcoplasmic reticulum in isolated ventricular myocytes from patients with terminal heart failure. J Mol Cell Cardiol. 1998; 30(4):743–749. [PubMed: 9602423]
- 39. Pieske B, Maier LS, Bers DM, Hasenfuss G. Ca2+ handling and sarcoplasmic reticulum Ca2+ content in isolated failing and nonfailing human myocardium. Circ Res. 1999; 85(1):38–46. [PubMed: 10400909]
- Hobai IA, O'Rourke B. Decreased sarcoplasmic reticulum calcium content is responsible for defective excitation-contraction coupling in canine heart failure. Circulation. 2001; 103(11):1577– 1584. [PubMed: 11257088]
- 41. Dipla K, Mattiello JA, Margulies KB, Jeevanandam V, Houser SR. The sarcoplasmic reticulum and the Na+/Ca2+ exchanger both contribute to the Ca2+ transient of failing human ventricular myocytes. Circ Res. 1999; 84(4):435–444. [PubMed: 10066678]
- 42. Pieske B, Maier LS, Piacentino V 3rd, Weisser J, Hasenfuss G, Houser S. Rate dependence of [Na +]i and contractility in nonfailing and failing human myocardium. Circulation. 2002; 106(4):447–453. [PubMed: 12135944]
- 43. Hasenfuss G, Pieske B. Calcium cycling in congestive heart failure. J Mol Cell Cardiol. 2002; 34(8):951–969. [PubMed: 12234765]
- 44. Feldman MD, Alderman JD, Aroesty JM, Royal HD, Ferguson JJ, Owen RM, Grossman W, McKay RG. Depression of systolic and diastolic myocardial reserve during atrial pacing tachycardia in patients with dilated cardiomyopathy. J Clin Invest. 1988; 82(5):1661–1669. [PubMed: 3183060]
- 45. Mulieri LA, Hasenfuss G, Leavitt B, Allen PD, Alpert NR. Altered myocardial force-frequency relation in human heart failure. Circulation. 1992; 85(5):1743–1750. [PubMed: 1572031]
- 46. Hasenfuss G, Reinecke H, Studer R, Meyer M, Pieske B, Holtz J, Holubarsch C, Posival H, Just H, Drexler H. Relation between myocardial function and expression of sarcoplasmic reticulum Ca(2+)-ATPase in failing and nonfailing human myocardium. Circ Res. 1994; 75(3):434–442. [PubMed: 8062417]
- 47. Swedberg K, Komajda M, Bohm M, Borer JS, Ford I, Dubost-Brama A, Lerebours G, Tavazzi L. Ivabradine and outcomes in chronic heart failure (SHIFT): a randomised placebo-controlled study. Lancet. 2010; 376(9744):875–885. [PubMed: 20801500]
- 48. Ørn S, Dickstein K. How do heart failure patients die? European Heart Journal Supplements. 2002; 4(suppl D):D59.
- 49. Bardy GH, Lee KL, Mark DB, Poole JE, Packer DL, Boineau R, Domanski M, Troutman C, Anderson J, Johnson G, McNulty SE, Clapp-Channing N, Davidson-Ray LD, Fraulo ES, Fishbein DP, Luceri RM, Ip JH. Amiodarone or an implantable cardioverter-defibrillator for congestive heart failure. N Engl J Med. 2005; 352(3):225–237. [PubMed: 15659722]
- 50. Tomaselli GF, Marban E. Electrophysiological remodeling in hypertrophy and heart failure. Cardiovasc Res. 1999; 42(2):270–283. [PubMed: 10533566]
- Pogwizd SM, Schlotthauer K, Li L, Yuan W, Bers DM. Arrhythmogenesis and contractile dysfunction in heart failure: Roles of sodium-calcium exchange, inward rectifier potassium current, and residual beta-adrenergic responsiveness. Circ Res. 2001; 88(11):1159–1167. [PubMed: 11397782]
- 52. Belevych AE, Terentyev D, Terentyeva R, Nishijima Y, Sridhar A, Hamlin RL, Carnes CA, Gyorke S. The relationship between arrhythmogenesis and impaired contractility in heart failure: role of altered ryanodine receptor function. Cardiovasc Res. 2011; 90(3):493–502. [PubMed: 21273243]
- 53. Tomaselli GF, Beuckelmann DJ, Calkins HG, Berger RD, Kessler PD, Lawrence JH, Kass D, Feldman AM, Marban E. Sudden cardiac death in heart failure. The role of abnormal repolarization. Circulation. 1994; 90(5):2534–2539. [PubMed: 7955213]
- 54. Weiss JN, Garfinkel A, Karagueuzian HS, Chen PS, Qu Z. Early afterdepolarizations and cardiac arrhythmias. Heart Rhythm. 2010; 7(12):1891–1899. [PubMed: 20868774]

55. Pogwizd SM, Bers DM. Cellular basis of triggered arrhythmias in heart failure. Trends Cardiovasc Med. 2004; 14(2):61–66. [PubMed: 15030791]

- Pogwizd SM, McKenzie JP, Cain ME. Mechanisms underlying spontaneous and induced ventricular arrhythmias in patients with idiopathic dilated cardiomyopathy. Circulation. 1998; 98(22):2404–2414. [PubMed: 9832485]
- 57. Vermeulen JT, McGuire MA, Opthof T, Coronel R, de Bakker JM, Klopping C, Janse MJ. Triggered activity and automaticity in ventricular trabeculae of failing human and rabbit hearts. Cardiovasc Res. 1994; 28(10):1547–1554. [PubMed: 8001044]
- 58. Pruvot EJ, Katra RP, Rosenbaum DS, Laurita KR. Role of calcium cycling versus restitution in the mechanism of repolarization alternans. Circ Res. 2004; 94(8):1083–1090. [PubMed: 15016735]
- 59. Wilson LD, Jeyaraj D, Wan X, Hoeker GS, Said TH, Gittinger M, Laurita KR, Rosenbaum DS. Heart failure enhances susceptibility to arrhythmogenic cardiac alternans. Heart Rhythm. 2009; 6(2):251–259. [PubMed: 19187920]
- 60. Klingenheben T, Zabel M, D'Agostino RB, Cohen RJ, Hohnloser SH. Predictive value of T-wave alternans for arrhythmic events in patients with congestive heart failure. Lancet. 2000; 356(9230): 651–652. [PubMed: 10968440]
- Pastore JM, Girouard SD, Laurita KR, Akar FG, Rosenbaum DS. Mechanism linking T-wave alternans to the genesis of cardiac fibrillation. Circulation. 1999; 99(10):1385–1394. [PubMed: 10077525]
- 62. Cutler MJ, Wan X, Laurita KR, Hajjar RJ, Rosenbaum DS. Targeted SERCA2a gene expression identifies molecular mechanism and therapeutic target for arrhythmogenic cardiac alternans. Circ Arrhythm Electrophysiol. 2009; 2(6):686–694. [PubMed: 19948504]
- 63. Mewes T, Ravens U. L-type calcium currents of human myocytes from ventricle of non-failing and failing hearts and from atrium. J Mol Cell Cardiol. 1994; 26(10):1307–1320. [PubMed: 7869391]
- 64. Sipido KR, Stankovicova T, Flameng W, Vanhaecke J, Verdonck F. Frequency dependence of Ca2+ release from the sarcoplasmic reticulum in human ventricular myocytes from end-stage heart failure. Cardiovasc Res. 1998; 37(2):478–488. [PubMed: 9614502]
- 65. Lindner M, Brandt MC, Sauer H, Hescheler J, Bohle T, Beuckelmann DJ. Calcium sparks in human ventricular cardiomyocytes from patients with terminal heart failure. Cell Calcium. 2002; 31(4):175–182. [PubMed: 12027382]
- 66. Crossman DJ, Ruygrok PR, Soeller C, Cannell MB. Changes in the organization of excitation-contraction coupling structures in failing human heart. PLoS One. 2011; 6(3):e17901. [PubMed: 21408028]
- 67. Lyon AR, MacLeod KT, Zhang Y, Garcia E, Kanda GK, Lab MJ, Korchev YE, Harding SE, Gorelik J. Loss of T-tubules and other changes to surface topography in ventricular myocytes from failing human and rat heart. Proc Natl Acad Sci USA. 2009; 106(16):6854–6859. [PubMed: 19342485]
- 68. Ohler A, Weisser-Thomas J, Piacentino V, Houser SR, Tomaselli GF, O'Rourke B. Two-photon laser scanning microscopy of the transverse-axial tubule system in ventricular cardiomyocytes from failing and non-failing human hearts. Cardiol Res Pract. 2009; 2009:802373. [PubMed: 20224636]
- 69. Song LS, Sobie EA, McCulle S, Lederer WJ, Balke CW, Cheng H. Orphaned ryanodine receptors in the failing heart. Proc Natl Acad Sci USA. 2006; 103(11):4305–4310. [PubMed: 16537526]
- 70. Louch WE, Bito V, Heinzel FR, Macianskiene R, Vanhaecke J, Flameng W, Mubagwa K, Sipido KR. Reduced synchrony of Ca2+ release with loss of T-tubules-a comparison to Ca2+ release in human failing cardiomyocytes. Cardiovasc Res. 2004; 62(1):63–73. [PubMed: 15023553]
- Wasserstrom JA, Sharma R, Kapur S, Kelly JE, Kadish AH, Balke CW, Aistrup GL. Multiple defects in intracellular calcium cycling in whole failing rat heart. Circ Heart Fail. 2009; 2(3):223– 232. [PubMed: 19808344]
- 72. Jantunen E, Collan Y. Transmural differences in ischaemic heart disease: a quantitative histologic study. Appl Pathol. 1989; 7(3):179–187. [PubMed: 2765296]
- 73. Figueredo VM, Brandes R, Weiner MW, Massie BM, Camacho SA. Endocardial versus epicardial differences of intracellular free calcium under normal and ischemic conditions in perfused rat hearts. Circ Res. 1993; 72(5):1082–1090. [PubMed: 8477520]

74. Burashnikov A, Antzelevitch C. Reinduction of atrial fibrillation immediately after termination of the arrhythmia is mediated by late phase 3 early afterdepolarization-induced triggered activity. Circulation. 2003; 107(18):2355–2360. [PubMed: 12695296]

- 75. Sobie EA, Guatimosim S, Song LS, Lederer WJ. The challenge of molecular medicine: complexity versus Occam's razor. J Clin Invest. 2003; 111(6):801–803. [PubMed: 12639985]
- 76. Jaski BE, Jessup ML, Mancini DM, Cappola TP, Pauly DF, Greenberg B, Borow K, Dittrich H, Zsebo KM, Hajjar RJ. Calcium upregulation by percutaneous administration of gene therapy in cardiac disease (CUPID Trial), a first-in-human phase 1/2 clinical trial. J Card Fail. 2009; 15(3): 171–181. [PubMed: 19327618]
- 77. Kubalova Z, Terentyev D, Viatchenko-Karpinski S, Nishijima Y, Gyorke I, Terentyeva R, da Cunha DN, Sridhar A, Feldman DS, Hamlin RL, Carnes CA, Gyorke S. Abnormal intrastore calcium signaling in chronic heart failure. Proc Natl Acad Sci USA. 2005; 102(39):14104–14109. [PubMed: 16172392]
- 78. Jiang MT, Lokuta AJ, Farrell EF, Wolff MR, Haworth RA, Valdivia HH. Abnormal Ca2+ release, but normal ryanodine receptors, in canine and human heart failure. Circ Res. 2002; 91(11):1015–1022. [PubMed: 12456487]
- Li Y, Kranias EG, Mignery GA, Bers DM. Protein kinase A phosphorylation of the ryanodine receptor does not affect calcium sparks in mouse ventricular myocytes. Circ Res. 2002; 90(3):309– 316. [PubMed: 11861420]
- 80. Ai X, Curran JW, Shannon TR, Bers DM, Pogwizd SM. Ca2+/calmodulin-dependent protein kinase modulates cardiac ryanodine receptor phosphorylation and sarcoplasmic reticulum Ca2+ leak in heart failure. Circ Res. 2005; 97(12):1314–1322. [PubMed: 16269653]
- 81. Kushnir A, Shan J, Betzenhauser MJ, Reiken S, Marks AR. Role of CaMKIIdelta phosphorylation of the cardiac ryanodine receptor in the force frequency relationship and heart failure. Proc Natl Acad Sci USA. 2010; 107(22):10274–10279. [PubMed: 20479242]
- 82. De Koninck P, Schulman H. Sensitivity of CaM kinase II to the frequency of Ca2+ oscillations. Science. 1998; 279(5348):227–230. [PubMed: 9422695]
- 83. Zhao W, Uehara Y, Chu G, Song Q, Qian J, Young K, Kranias EG. Threonine-17 phosphorylation of phospholamban: a key determinant of frequency-dependent increase of cardiac contractility. J Mol Cell Cardiol. 2004; 37(2):607–612. [PubMed: 15276030]
- 84. Holmberg SR, Williams AJ. Single channel recordings from human cardiac sarcoplasmic reticulum. Circ Res. 1989; 65(5):1445–1449. [PubMed: 2478313]
- 85. Schmidt U, Hajjar RJ, Kim CS, Lebeche D, Doye AA, Gwathmey JK. Human heart failure: cAMP stimulation of SR Ca(2+)-ATPase activity and phosphorylation level of phospholamban. Am J Physiol. 1999; 277(2 Pt 2):H474–H480. [PubMed: 10444471]
- 86. Schwinger RH, Bohm M, Schmidt U, Karczewski P, Bavendiek U, Flesch M, Krause EG, Erdmann E. Unchanged protein levels of SERCA II and phospholamban but reduced Ca2+ uptake and Ca(2+)-ATPase activity of cardiac sarcoplasmic reticulum from dilated cardiomyopathy patients compared with patients with nonfailing hearts. Circulation. 1995; 92(11):3220–3228. [PubMed: 7586307]
- 87. del Monte F, Harding SE, Schmidt U, Matsui T, Kang ZB, Dec GW, Gwathmey JK, Rosenzweig A, Hajjar RJ. Restoration of contractile function in isolated cardiomyocytes from failing human hearts by gene transfer of SERCA2a. Circulation. 1999; 100(23):2308–2311. [PubMed: 10587333]
- 88. del Monte F, Harding SE, Dec GW, Gwathmey JK, Hajjar RJ. Targeting phospholamban by gene transfer in human heart failure. Circulation. 2002; 105(8):904–907. [PubMed: 11864915]
- 89. Frank KF, Bolck B, Brixius K, Kranias EG, Schwinger RH. Modulation of SERCA: implications for the failing human heart. Basic Res Cardiol. 2002; 97(Suppl 1):I72–I78. [PubMed: 12479238]
- 90. Schwinger RH, Munch G, Bolck B, Karczewski P, Krause EG, Erdmann E. Reduced Ca(2+)-sensitivity of SERCA 2a in failing human myocardium due to reduced serin-16 phospholamban phosphorylation. J Mol Cell Cardiol. 1999; 31(3):479–491. [PubMed: 10198180]
- 91. Dash R, Frank KF, Carr AN, Moravec CS, Kranias EG. Gender influences on sarcoplasmic reticulum Ca2+-handling in failing human myocardium. J Mol Cell Cardiol. 2001; 33(7):1345–1353. [PubMed: 11437540]

92. Munch G, Bolck B, Karczewski P, Schwinger RH. Evidence for calcineurin-mediated regulation of SERCA 2a activity in human myocardium. J Mol Cell Cardiol. 2002; 34(3):321–334. [PubMed: 11945024]

- 93. Mishra S, Gupta RC, Tiwari N, Sharov VG, Sabbah HN. Molecular mechanisms of reduced sarcoplasmic reticulum Ca(2+) uptake in human failing left ventricular myocardium. J Heart Lung Transplant. 2002; 21(3):366–373. [PubMed: 11897526]
- 94. Minamisawa S, Hoshijima M, Chu G, Ward CA, Frank K, Gu Y, Martone ME, Wang Y, Ross J Jr, Kranias EG, Giles WR, Chien KR. Chronic phospholamban-sarcoplasmic reticulum calcium ATPase interaction is the critical calcium cycling defect in dilated cardiomyopathy. Cell. 1999; 99(3):313–322. [PubMed: 10555147]
- 95. Liu Q, Chen X, Macdonnell SM, Kranias EG, Lorenz JN, Leitges M, Houser SR, Molkentin JD. Protein kinase C{alpha}, but not PKC{beta} or PKC{gamma}, regulates contractility and heart failure susceptibility: implications for ruboxistaurin as a novel therapeutic approach. Circ Res. 2009; 105(2):194–200. [PubMed: 19556521]
- 96. Weber CR, Piacentino V 3rd, Houser SR, Bers DM. Dynamic regulation of sodium/calcium exchange function in human heart failure. Circulation. 2003; 108(18):2224–2229. [PubMed: 14557358]
- 97. Ventura-Clapier R, Garnier A, Veksler V. Energy metabolism in heart failure. J Physiol. 2004; 555(Pt 1):1–13. [PubMed: 14660709]
- 98. Bers DM. Calcium cycling and signaling in cardiac myocytes. Annu Rev Physiol. 2008; 70:23–49. [PubMed: 17988210]
- De Sousa E, Veksler V, Minajeva A, Kaasik A, Mateo P, Mayoux E, Hoerter J, Bigard X, Serrurier B, Ventura-Clapier R. Subcellular creatine kinase alterations. Implications in heart failure. Circ Res. 1999; 85(1):68–76. [PubMed: 10400912]
- 100. Xu KY, Zweier JL, Becker LC. Hydroxyl radical inhibits sarcoplasmic reticulum Ca(2+)-ATPase function by direct attack on the ATP binding site. Circ Res. 1997; 80(1):76–81. [PubMed: 8978325]
- 101. Hasenfuss G, Maier LS, Hermann HP, Luers C, Hunlich M, Zeitz O, Janssen PM, Pieske B. Influence of pyruvate on contractile performance and Ca(2+) cycling in isolated failing human myocardium. Circulation. 2002; 105(2):194–199. [PubMed: 11790700]
- 102. Reiken S, Wehrens XH, Vest JA, Barbone A, Klotz S, Mancini D, Burkhoff D, Marks AR. Betablockers restore calcium release channel function and improve cardiac muscle performance in human heart failure. Circulation. 2003; 107(19):2459–2466. [PubMed: 12743001]
- 103. Chaudhary KW, Rossman EI, Piacentino V 3rd, Kenessey A, Weber C, Gaughan JP, Ojamaa K, Klein I, Bers DM, Houser SR, Margulies KB. Altered myocardial Ca2+ cycling after left ventricular assist device support in the failing human heart. J Am Coll Cardiol. 2004; 44(4):837–845. [PubMed: 15312868]
- 104. Vanderheyden M, Mullens W, Delrue L, Goethals M, de Bruyne B, Wijns W, Geelen P, Verstreken S, Wellens F, Bartunek J. Myocardial gene expression in heart failure patients treated with cardiac resynchronization therapy responders versus nonresponders. J Am Coll Cardiol. 2008; 51(2):129–136. [PubMed: 18191736]
- 105. del Monte F, Williams E, Lebeche D, Schmidt U, Rosenzweig A, Gwathmey JK, Lewandowski ED, Hajjar RJ. Improvement in survival and cardiac metabolism after gene transfer of sarcoplasmic reticulum Ca(2+)-ATPase in a rat model of heart failure. Circulation. 2001; 104(12):1424–1429. [PubMed: 11560860]
- 106. Rockman HA, Koch WJ, Lefkowitz RJ. Seven-transmembrane-spanning receptors and heart function. Nature. 2002; 415(6868):206–212. [PubMed: 11805844]
- 107. Cambronero F, Marin F, Roldan V, Hernandez-Romero D, Valdes M, Lip GY. Biomarkers of pathophysiology in hypertrophic cardiomyopathy: implications for clinical management and prognosis. Eur Heart J. 2009; 30(2):139–151. [PubMed: 19136482]
- 108. Hunt SA, Abraham WT, Chin MH, Feldman AM, Francis GS, Ganiats TG, Jessup M, Konstam MA, Mancini DM, Michl K, Oates JA, Rahko PS, Silver MA, Stevenson LW, Yancy CW. 2009 Focused update incorporated into the ACC/AHA 2005 Guidelines for the Diagnosis and Management of Heart Failure in Adults A Report of the American College of Cardiology

- Foundation/American Heart Association Task Force on Practice Guidelines Developed in Collaboration With the International Society for Heart and Lung Transplantation. Journal of the American College of Cardiology. 2009; 53(15):e1–e90. [PubMed: 19358937]
- 109. Wehrens XH, Lehnart SE, Huang F, Vest JA, Reiken SR, Mohler PJ, Sun J, Guatimosim S, Song LS, Rosemblit N, D'Armiento JM, Napolitano C, Memmi M, Priori SG, Lederer WJ, Marks AR. FKBP12.6 deficiency and defective calcium release channel (ryanodine receptor) function linked to exercise-induced sudden cardiac death. Cell. 2003; 113(7):829–840. [PubMed: 12837242]
- 110. DeSantiago J, Maier LS, Bers DM. Frequency-dependent acceleration of relaxation in the heart depends on CaMKII, but not phospholamban. Journal of molecular and cellular cardiology. 2002; 34(8):975–984. [PubMed: 12234767]
- 111. Sugden PH, Bogoyevitch MA. Intracellular signalling through protein kinases in the heart. Cardiovascular research. 1995; 30(4):478–492. [PubMed: 8574996]
- 112. Kluger J, Cody RJ, Laragh JH. The contributions of sympathetic tone and the renin-angiotensin system to severe chronic congestive heart failure: response to specific inhibitors (prazosin and captopril). Am J Cardiol. 1982; 49(7):1667–1674. [PubMed: 7044087]
- 113. Cohn JN, Levine TB, Olivari MT, Garberg V, Lura D, Francis GS, Simon AB, Rector T. Plasma norepinephrine as a guide to prognosis in patients with chronic congestive heart failure. The New England journal of medicine. 1984; 311(13):819–823. [PubMed: 6382011]
- 114. Ungerer M, Bohm M, Elce JS, Erdmann E, Lohse MJ. Altered expression of beta-adrenergic receptor kinase and beta 1-adrenergic receptors in the failing human heart. Circulation. 1993; 87(2):454–463. [PubMed: 8381058]
- 115. Daaka Y, Luttrell LM, Lefkowitz RJ. Switching of the coupling of the beta2-adrenergic receptor to different G proteins by protein kinase A. Nature. 1997; 390(6655):88–91. [PubMed: 9363896]
- 116. Bristow MR, Minobe WA, Raynolds MV, Port JD, Rasmussen R, Ray PE, Feldman AM. Reduced beta 1 receptor messenger RNA abundance in the failing human heart. The Journal of clinical investigation. 1993; 92(6):2737–2745. [PubMed: 8254027]
- 117. Wei SK, Ruknudin A, Hanlon SU, McCurley JM, Schulze DH, Haigney MC. Protein kinase A hyperphosphorylation increases basal current but decreases beta-adrenergic responsiveness of the sarcolemmal Na+-Ca2+ exchanger in failing pig myocytes. Circulation research. 2003; 92(8): 897–903. [PubMed: 12676818]
- 118. A trial of the beta-blocker bucindolol in patients with advanced chronic heart failure. The New England journal of medicine. 2001; 344(22):1659–1667. [PubMed: 11386264]
- 119. Packer M, Bristow MR, Cohn JN, Colucci WS, Fowler MB, Gilbert EM, Shusterman NH. The effect of carvedilol on morbidity and mortality in patients with chronic heart failure. U.S. Carvedilol Heart Failure Study Group. N Engl J Med. 1996; 334(21):1349–1355. [PubMed: 8614419]
- 120. MERIT-HFTrial. Effect of metoprolol CR/XL in chronic heart failure: Metoprolol CR/XL Randomised Intervention Trial in Congestive Heart Failure (MERIT-HF). Lancet. 1999; 353(9169):2001–2007. [PubMed: 10376614]
- 121. CIBIS-IITrial. The Cardiac Insufficiency Bisoprolol Study II (CIBIS-II): a randomised trial. Lancet. 1999; 353(9146):9–13. [PubMed: 10023943]
- 122. Yancy CW. Comprehensive treatment of heart failure: state-of-the-art medical therapy. Rev Cardiovasc Med. 2005; 6(Suppl 2):S43–S57. [PubMed: 15891702]
- 123. Bristow MR. beta-adrenergic receptor blockade in chronic heart failure. Circulation. 2000; 101(5):558–569. [PubMed: 10662755]
- 124. Lowes BD, Gilbert EM, Abraham WT, Minobe WA, Larrabee P, Ferguson D, Wolfel EE, Lindenfeld J, Tsvetkova T, Robertson AD, Quaife RA, Bristow MR. Myocardial gene expression in dilated cardiomyopathy treated with beta-blocking agents. N Engl J Med. 2002; 346(18):1357– 1365. [PubMed: 11986409]
- 125. Reiken S, Gaburjakova M, Gaburjakova J, He Kl KL, Prieto A, Becker E, Yi Gh GH, Wang J, Burkhoff D, Marks AR. beta-adrenergic receptor blockers restore cardiac calcium release channel (ryanodine receptor) structure and function in heart failure. Circulation. 2001; 104(23): 2843–2848. [PubMed: 11733405]

126. Reiken S, Gaburjakova M, Guatimosim S, Gomez AM, D'Armiento J, Burkhoff D, Wang J, Vassort G, Lederer WJ, Marks AR. Protein kinase A phosphorylation of the cardiac calcium release channel (ryanodine receptor) in normal failing hearts. Role of phosphatases and response to isoproterenol. The Journal of biological chemistry. 2003; 278(1):444–453. [PubMed: 12401811]

- 127. Yano M, Ono K, Ohkusa T, Suetsugu M, Kohno M, Hisaoka T, Kobayashi S, Hisamatsu Y, Yamamoto T, Noguchi N, Takasawa S, Okamoto H, Matsuzaki M. Altered stoichiometry of FKBP12.6 versus ryanodine receptor as a cause of abnormal Ca(2+) leak through ryanodine receptor in heart failure. Circulation. 2000; 102(17):2131–2136. [PubMed: 11044432]
- 128. Doi M, Yano M, Kobayashi S, Kohno M, Tokuhisa T, Okuda S, Suetsugu M, Hisamatsu Y, Ohkusa T, Matsuzaki M. Propranolol prevents the development of heart failure by restoring FKBP12.6-mediated stabilization of ryanodine receptor. Circulation. 2002; 105(11):1374–1379. [PubMed: 11901051]
- 129. George I, Sabbah HN, Xu K, Wang N, Wang J. {beta}-Adrenergic Receptor Blockade Reduces Endoplasmic Reticulum Stress and Normalizes Calcium Handling in a Coronary Embolization Model of Heart Failure in Canines. Cardiovascular research. 2011
- 130. Beuckelmann DJ, Nabauer M, Kruger C, Erdmann E. Altered diastolic [Ca2+]i handling in human ventricular myocytes from patients with terminal heart failure. Am Heart J. 1995; 129(4): 684–689. [PubMed: 7900618]
- 131. Pass JM, Gao J, Jones WK, Wead WB, Wu X, Zhang J, Baines CP, Bolli R, Zheng YT, Joshua IG, Ping P. Enhanced PKC beta II translocation and PKC beta II-RACK1 interactions in PKC epsilon-induced heart failure: a role for RACK1. American journal of physiology. Heart and circulatory physiology. 2001; 281(6):H2500–H2510. [PubMed: 11709417]
- 132. Silberman GA, Fan TH, Liu H, Jiao Z, Xiao HD, Lovelock JD, Boulden BM, Widder J, Fredd S, Bernstein KE, Wolska BM, Dikalov S, Harrison DG, Dudley SC Jr. Uncoupled cardiac nitric oxide synthase mediates diastolic dysfunction. Circulation. 2010; 121(4):519–528. [PubMed: 20083682]
- 133. Mende U, Kagen A, Cohen A, Aramburu J, Schoen FJ, Neer EJ. Transient cardiac expression of constitutively active Galphaq leads to hypertrophy and dilated cardiomyopathy by calcineurindependent and independent pathways. Proceedings of the National Academy of Sciences of the United States of America. 1998; 95(23):13893–13898. [PubMed: 9811897]
- 134. Taigen T, De Windt LJ, Lim HW, Molkentin JD. Targeted inhibition of calcineurin prevents agonist-induced cardiomyocyte hypertrophy. Proceedings of the National Academy of Sciences of the United States of America. 2000; 97(3):1196–1201. [PubMed: 10655507]
- 135. Thomas WG, Brandenburger Y, Autelitano DJ, Pham T, Qian H, Hannan RD. Adenoviral-directed expression of the type 1A angiotensin receptor promotes cardiomyocyte hypertrophy via transactivation of the epidermal growth factor receptor. Circulation research. 2002; 90(2):135–142. [PubMed: 11834705]
- 136. Wilkins BJ, Molkentin JD. Calcium-calcineurin signaling in the regulation of cardiac hypertrophy. Biochem Biophys Res Commun. 2004; 322(4):1178–1191. [PubMed: 15336966]
- 137. Sussman MA, Lim HW, Gude N, Taigen T, Olson EN, Robbins J, Colbert MC, Gualberto A, Wieczorek DF, Molkentin JD. Prevention of cardiac hypertrophy in mice by calcineurin inhibition. Science. 1998; 281(5383):1690–1693. [PubMed: 9733519]
- 138. Maack C, Kartes T, Kilter H, Schafers HJ, Nickenig G, Bohm M, Laufs U. Oxygen free radical release in human failing myocardium is associated with increased activity of rac1-GTPase and represents a target for statin treatment. Circulation. 2003; 108(13):1567–1574. [PubMed: 12963641]
- 139. Giordano FJ. Oxygen, oxidative stress, hypoxia, and heart failure. The Journal of clinical investigation. 2005; 115(3):500–508. [PubMed: 15765131]
- 140. Sawyer DB, Siwik DA, Xiao L, Pimentel DR, Singh K, Colucci WS. Role of oxidative stress in myocardial hypertrophy and failure. Journal of molecular and cellular cardiology. 2002; 34(4): 379–388. [PubMed: 11991728]
- 141. Seshiah PN, Weber DS, Rocic P, Valppu L, Taniyama Y, Griendling KK. Angiotensin II stimulation of NAD(P)H oxidase activity: upstream mediators. Circulation research. 2002; 91(5): 406–413. [PubMed: 12215489]

142. Rathore N, John S, Kale M, Bhatnagar D. Lipid peroxidation and antioxidant enzymes in isoproterenol induced oxidative stress in rat tissues. Pharmacol Res. 1998; 38(4):297–303. [PubMed: 9774493]

- 143. Thollon C, Iliou JP, Cambarrat C, Robin F, Vilaine JP. Nature of the cardiomyocyte injury induced by lipid hydroperoxides. Cardiovascular research. 1995; 30(5):648–655. [PubMed: 8595608]
- 144. Suematsu N, Tsutsui H, Wen J, Kang D, Ikeuchi M, Ide T, Hayashidani S, Shiomi T, Kubota T, Hamasaki N, Takeshita A. Oxidative stress mediates tumor necrosis factor-alpha-induced mitochondrial DNA damage and dysfunction in cardiac myocytes. Circulation. 2003; 107(10): 1418–1423. [PubMed: 12642364]
- 145. Nishida M, Maruyama Y, Tanaka R, Kontani K, Nagao T, Kurose H. G alpha(i) and G alpha(o) are target proteins of reactive oxygen species. Nature. 2000; 408(6811):492–495. [PubMed: 11100733]
- 146. Guerra L, Cerbai E, Gessi S, Borea PA, Mugelli A. The effect of oxygen free radicals on calcium current and dihydropyridine binding sites in guinea-pig ventricular myocytes. Br J Pharmacol. 1996; 118(5):1278–1284. [PubMed: 8818354]
- 147. Kaplan P, Babusikova E, Lehotsky J, Dobrota D. Free radical-induced protein modification and inhibition of Ca2+-ATPase of cardiac sarcoplasmic reticulum. Molecular and cellular biochemistry. 2003; 248(1–2):41–47. [PubMed: 12870653]
- 148. Goldhaber JI. Free radicals enhance Na+/Ca2+ exchange in ventricular myocytes. The American journal of physiology. 1996; 271(3 Pt 2):H823–H833. [PubMed: 8853314]
- 149. Okuda S, Yano M, Doi M, Oda T, Tokuhisa T, Kohno M, Kobayashi S, Yamamoto T, Ohkusa T, Matsuzaki M. Valsartan restores sarcoplasmic reticulum function with no appreciable effect on resting cardiac function in pacing-induced heart failure. Circulation. 2004; 109(7):911–919. [PubMed: 14757694]
- 150. Agostoni PG, De Cesare N, Doria E, Polese A, Tamborini G, Guazzi MD. Afterload reduction: a comparison of captopril and nifedipine in dilated cardiomyopathy. Br Heart J. 1986; 55(4):391–399. [PubMed: 3516187]
- 151. Tan LB, Murray RG, Littler WA. Felodipine in patients with chronic heart failure: discrepant haemodynamic and clinical effects. Br Heart J. 1987; 58(2):122–128. [PubMed: 3304372]
- 152. Elkayam U, Amin J, Mehra A, Vasquez J, Weber L, Rahimtoola SH. A prospective, randomized, double-blind, crossover study to compare the efficacy and safety of chronic nifedipine therapy with that of isosorbide dinitrate and their combination in the treatment of chronic congestive heart failure. Circulation. 1990; 82(6):1954–1961. [PubMed: 2242521]
- 153. Cohn JN, Ziesche S, Smith R, Anand I, Dunkman WB, Loeb H, Cintron G, Boden W, Baruch L, Rochin P, Loss L. Effect of the calcium antagonist felodipine as supplementary vasodilator therapy in patients with chronic heart failure treated with enalapril: V-HeFT III. Vasodilator-Heart Failure Trial (V-HeFT) Study Group. Circulation. 1997; 96(3):856–863. [PubMed: 9264493]
- 154. Elkayam U, Parikh K, Torkan B, Weber L, Cohen JL, Rahimtoola SH. Effect of diltiazem on renal clearance and serum concentration of digoxin in patients with cardiac disease. Am J Cardiol. 1985; 55(11):1393–1395. [PubMed: 3993576]
- 155. Barjon JN, Rouleau JL, Bichet D, Juneau C, De Champlain J. Chronic renal and neurohumoral effects of the calcium entry blocker nisoldipine in patients with congestive heart failure. Journal of the American College of Cardiology. 1987; 9(3):622–630. [PubMed: 2880884]
- 156. Goldstein RE, Boccuzzi SJ, Cruess D, Nattel S. Diltiazem increases late-onset congestive heart failure in postinfarction patients with early reduction in ejection fraction. The Adverse Experience Committee; and the Multicenter Diltiazem Postinfarction Research Group. Circulation. 1991; 83(1):52–60. [PubMed: 1984898]
- 157. Littler WA, Sheridan DJ. Placebo controlled trial of felodipine in patients with mild to moderate heart failure. UK Study Group. Br Heart J. 1995; 73(5):428–433. [PubMed: 7786657]
- 158. Mohindra SK, Udeani GO. Long-acting verapamil and heart failure. Jama. 1989; 261(7):994. [PubMed: 2915424]

159. Levine TB, Bernink PJ, Caspi A, Elkayam U, Geltman EM, Greenberg B, McKenna WJ, Ghali JK, Giles TD, Marmor A, Reisin LH, Ammon S, Lindberg E. Effect of mibefradil, a T-type calcium channel blocker, on morbidity and mortality in moderate to severe congestive heart failure: the MACH-1 study. Mortality Assessment in Congestive Heart Failure Trial. Circulation. 2000; 101(7):758–764. [PubMed: 10683349]

- 160. Mullins ME, Horowitz BZ, Linden DH, Smith GW, Norton RL, Stump J. Life-threatening interaction of mibefradil and beta-blockers with dihydropyridine calcium channel blockers. Jama. 1998; 280(2):157–158. [PubMed: 9669789]
- 161. Benatar D, Hall V, Reddy S, Gheorghiade M. Clinical and neurohormonal effects of nicardipine hydrochloride in patients with severe chronic heart failure receiving angiotensin-converting enzyme inhibitor therapy. Am J Ther. 1998; 5(1):25–32. [PubMed: 10099034]
- 162. Packer M, O'Connor CM, Ghali JK, Pressler ML, Carson PE, Belkin RN, Miller AB, Neuberg GW, Frid D, Wertheimer JH, Cropp AB, DeMets DL. Effect of amlodipine on morbidity and mortality in severe chronic heart failure. Prospective Randomized Amlodipine Survival Evaluation Study Group. The New England journal of medicine. 1996; 335(15):1107–1114. [PubMed: 8813041]
- 163. Freedman NJ, Liggett SB, Drachman DE, Pei G, Caron MG, Lefkowitz RJ. Phosphorylation and desensitization of the human beta 1-adrenergic receptor. Involvement of G protein-coupled receptor kinases and cAMP-dependent protein kinase. The Journal of biological chemistry. 1995; 270(30):17953–17961. [PubMed: 7629102]
- 164. Harding VB, Jones LR, Lefkowitz RJ, Koch WJ, Rockman HA. Cardiac beta ARK1 inhibition prolongs survival and augments beta blocker therapy in a mouse model of severe heart failure. Proceedings of the National Academy of Sciences of the United States of America. 2001; 98(10): 5809–5814. [PubMed: 11331748]
- 165. Rockman HA, Chien KR, Choi DJ, Iaccarino G, Hunter JJ, Ross J Jr, Lefkowitz RJ, Koch WJ. Expression of a beta-adrenergic receptor kinase 1 inhibitor prevents the development of myocardial failure in gene-targeted mice. Proceedings of the National Academy of Sciences of the United States of America. 1998; 95(12):7000–7005. [PubMed: 9618528]
- 166. Marx SO, Reiken S, Hisamatsu Y, Gaburjakova M, Gaburjakova J, Yang YM, Rosemblit N, Marks AR. Phosphorylation-dependent regulation of ryanodine receptors: a novel role for leucine/isoleucine zippers. J Cell Biol. 2001; 153(4):699–708. [PubMed: 11352932]
- 167. Most P, Koch WJ. Sealing the leak, healing the heart. Nature medicine. 2003; 9(8):993–994.
- 168. Wehrens XH, Lehnart SE, Reiken SR, Deng SX, Vest JA, Cervantes D, Coromilas J, Landry DW, Marks AR. Protection from cardiac arrhythmia through ryanodine receptor-stabilizing protein calstabin2. Science. 2004; 304(5668):292–296. [PubMed: 15073377]
- 169. Yano M, Kobayashi S, Kohno M, Doi M, Tokuhisa T, Okuda S, Suetsugu M, Hisaoka T, Obayashi M, Ohkusa T, Matsuzaki M. FKBP12.6-mediated stabilization of calcium-release channel (ryanodine receptor) as a novel therapeutic strategy against heart failure. Circulation. 2003; 107(3):477–484. [PubMed: 12551874]
- 170. Zhang R, Khoo MS, Wu Y, Yang Y, Grueter CE, Ni G, Price EE Jr, Thiel W, Guatimosim S, Song LS, Madu EC, Shah AN, Vishnivetskaya TA, Atkinson JB, Gurevich VV, Salama G, Lederer WJ, Colbran RJ, Anderson ME. Calmodulin kinase II inhibition protects against structural heart disease. Nature medicine. 2005; 11(4):409–417.
- 171. Limas CJ, Olivari MT, Goldenberg IF, Levine TB, Benditt DG, Simon A. Calcium uptake by cardiac sarcoplasmic reticulum in human dilated cardiomyopathy. Cardiovascular research. 1987; 21(8):601–605. [PubMed: 3328648]
- 172. Schmitt JP, Kamisago M, Asahi M, Li GH, Ahmad F, Mende U, Kranias EG, MacLennan DH, Seidman JG, Seidman CE. Dilated cardiomyopathy and heart failure caused by a mutation in phospholamban. Science. 2003; 299(5611):1410–1413. [PubMed: 12610310]
- 173. Huang B, Wang S, Qin D, Boutjdir M, El-Sherif N. Diminished basal phosphorylation level of phospholamban in the postinfarction remodeled rat ventricle: role of beta-adrenergic pathway, G(i) protein, phosphodiesterase, and phosphatases. Circulation research. 1999; 85(9):848–855. [PubMed: 10532953]
- 174. Hoshijima M, Ikeda Y, Iwanaga Y, Minamisawa S, Date MO, Gu Y, Iwatate M, Li M, Wang L, Wilson JM, Wang Y, Ross J Jr, Chien KR. Chronic suppression of heart-failure progression by a

- pseudophosphorylated mutant of phospholamban via in vivo cardiac rAAV gene delivery. Nature medicine. 2002; 8(8):864–871.
- 175. Iwanaga Y, Hoshijima M, Gu Y, Iwatate M, Dieterle T, Ikeda Y, Date MO, Chrast J, Matsuzaki M, Peterson KL, Chien KR, Ross J Jr. Chronic phospholamban inhibition prevents progressive cardiac dysfunction and pathological remodeling after infarction in rats. The Journal of clinical investigation. 2004; 113(5):727–736. [PubMed: 14991071]
- 176. Kaye DM, Preovolos A, Marshall T, Byrne M, Hoshijima M, Hajjar R, Mariani JA, Pepe S, Chien KR, Power JM. Percutaneous cardiac recirculation-mediated gene transfer of an inhibitory phospholamban peptide reverses advanced heart failure in large animals. Journal of the American College of Cardiology. 2007; 50(3):253–260. [PubMed: 17631218]
- 177. Yamada M, Ikeda Y, Yano M, Yoshimura K, Nishino S, Aoyama H, Wang L, Aoki H, Matsuzaki M. Inhibition of protein phosphatase 1 by inhibitor-2 gene delivery ameliorates heart failure progression in genetic cardiomyopathy. Faseb J. 2006; 20(8):1197–1199. [PubMed: 16627625]
- 178. Dawson J, Walters M. Uric acid and xanthine oxidase: future therapeutic targets in the prevention of cardiovascular disease? Br J Clin Pharmacol. 2006
- 179. Spector T. Oxypurinol as an inhibitor of xanthine oxidase-catalyzed production of superoxide radical. Biochem Pharmacol. 1988; 37(2):349–352. [PubMed: 2829916]
- 180. Cappola TP, Kass DA, Nelson GS, Berger RD, Rosas GO, Kobeissi ZA, Marban E, Hare JM. Allopurinol improves myocardial efficiency in patients with idiopathic dilated cardiomyopathy. Circulation. 2001; 104(20):2407–2411. [PubMed: 11705816]
- 181. Freudenberger RS, Schwarz RP Jr, Brown J, Moore A, Mann D, Givertz MM, Colucci WS, Hare JM. Rationale, design and organisation of an efficacy and safety study of oxypurinol added to standard therapy in patients with NYHA class III IV congestive heart failure. Expert Opin Investig Drugs. 2004; 13(11):1509–1516.
- 182. Hare JM, Mangal B, Brown J, Fisher C Jr, Freudenberger R, Colucci WS, Mann DL, Liu P, Givertz MM, Schwarz RP. Impact of oxypurinol in patients with symptomatic heart failure. Results of the OPT-CHF study. Journal of the American College of Cardiology. 2008; 51(24): 2301–2309. [PubMed: 18549913]
- 183. Brillantes AM, Allen P, Takahashi T, Izumo S, Marks AR. Differences in cardiac calcium release channel (ryanodine receptor) expression in myocardium from patients with end-stage heart failure caused by ischemic versus dilated cardiomyopathy. Circ Res. 1992; 71(1):18–26. [PubMed: 1318794]
- 184. Sen L, Cui G, Fonarow GC, Laks H. Differences in mechanisms of SR dysfunction in ischemic vs. idiopathic dilated cardiomyopathy. Am J Physiol Heart Circ Physiol. 2000; 279(2):H709– H718. [PubMed: 10924070]
- 185. Houser SR. Reduced abundance of transverse tubules and L-type calcium channels: another cause of defective contractility in failing ventricular myocytes. Cardiovasc Res. 2001; 49(2):253–256. [PubMed: 11164835]
- 186. Efimov IR, Fedorov VV, Glukhov A, Lou Q, Ambrosi C, Janks D, Hucker WJ, Kurian T, Schuessler RB, Moazami N. Multiscale imaging of the human heart: Building the foundation for human systems physiology and translational medicine. Conf Proc IEEE Eng Med Biol Soc. 2010; 2010:5177–5180. [PubMed: 21095821]




Figure 1. Intracellular Ca^{2+} cycling and regulation by signaling pathways. After the activation of sarcolemma (including T-tubules), Ca^{2+} enters cytoplasm through L-type Ca^{2+} channel. The entering Ca^{2+} then induces a much larger Ca^{2+} release from the sarcoplasmic reticulum (SR) via the ryanodine receptor (RyR). The released Ca^{2+} binds with Troponin-C to activate contraction. Relaxation starts when Ca^{2+} is returned by sarcoplasmic reticulum Ca^{2+} ATPase (SERCA) to SR and via the Na^+/Ca^{2+} exchanger (NCX) to the extracellular domain. Some Ca^{2+} enters mitochondria to stimulate the production of ATP which is utilized for contraction and transcription. SERCA is inhibited by the dephosphorylated phospholamban (PLN). PLN can be phosphorylated by protein kinase A (PKA) and $Ca^{2+}/calmodulin$

dependent kinase (CAMKII), both of which can be activated by β -adrenergic stimulation. PLN can be dephosphorylated by phosphotase 1 (PP1), which can be activated through Gaq-coupled receptors (angiotensin II receptor, endothelin 1 receptor, or α -adrenergic receptor). α , G-protein subunit α ; β , G-protein subunit β ; γ , G-protein subunit γ ; AC, adenylate cyclase; cAMP, cyclic adenosine monophosphate.




Figure 2.Region-dependent and cycle-length-dependent calcium transient duration (CaTD) in failing human heart. (A) CaTD was quantified at 80% relaxation (CaTD80). (B) CaTD80 at nonfailing human hearts (n=6) at subendocardium (sub-ENDO), midmyocardium (MID), and subepicardium (sub-EPI). (C) CaTD80 at failing human hearts (n=5). (D) The difference of CaTD80 between sub-ENDO and sub-EPI. It can be seen that this difference is significantly reduced in failing heart at faster heart rate (cycle length at 600ms).³¹ These data are obtained from Ca²⁺ transient measured using Rhod-2AM from the coronary-perfused wedge preparations from both failing and non-failing human hearts.




Figure 3.

Morphological changes of calcium transient (CaT) and its relation to action potential (AP).

(A) Simultaneous recordings of AP and CaT at one site at subendocardium from a failing human heart (F, top) and a nonfailing human heart (NF, bottom). (B) The two CaTs from panel A are overlapped for easy comparison. Compared to the CaT from NF, there is a distinct second rising component (labeled by "II") in the CaT from failing human heart. Note that this second component was only observed at the sub-endocardium in 60% of the studied failing human hearts. (C) CaT duration at 80% relaxation (CaTD80) minus AP duration at 80% (APD80), summarized from five nonfailing hearts and five failing hearts (two ischemic cardiomyopathy and three idiopathic cardiomyopathy). It can be seen that this duration difference is significantly longer at the subendocardium in the failing human heart compared with the non-failing human heart, which is reflected in the example shown in panel A.³¹




Figure 4.Protein expressions of sarcoplasmic reticulum Ca²⁺ ATPase 2a (SERCA2a) and phospholamban (PLN). Representative examples of Western blots (top) and normalized protein expression (bottom) are shown for SERCA2a (A) and PLN (B). NF (n=6) indicates the group of non-failing hearts; Isch-F (n=6), the group of failing hearts with ischemic cardiomyopathy; and Nonischemic-F (n=6), the group of failing hearts with nonischemic/idiopathic cardiomyopathy.³¹ ENDO indicates endocardium; EPI, epicardium.