

Chapter 9 Discrete Mathematics Notes

Miagao

Contents

9 Counting and Probability	2
9.1 Introduction to Probability	2
Counting the Elements of a List	2
9.2 Possibility Trees and the Multiplication Rule	3
The Multiplication Rule	3
Permutations	4
Permutations of Selected Elements	5
9.3 Counting Elements of Disjoint Sets: The Addition Rule	6
The Difference Rule	6
The Inclusion/Exclusion Rule	7
9.4 The Pigeonhole Principle	8
Generalized Pigeonhole Principle	8
9.5 Counting Subsets of a Set: Combinations	9
Double Counting	11
9.6 r -Combinations with Repetition Allowed	13
9.7 Pascal's Formula and the Binomial Theorem	14

9 Counting and Probability

9.1 Introduction to Probability

- A process is **random** when it leads to an outcome from some set of outcomes, and it is impossible to predict with certainty what it may be.
 - In a coin flip, the set of outcomes is heads and tails, but it cannot be predicted with certainty if it is heads or tails, even considering a calculated probability of 50%.

Definition:

A **sample space** is the set of all possible outcomes of a random process or experiment.

An **event** is a subset of a sample space.

- Formally, given S , a finite sample space where all outcomes are equally like, and E , an event in S , then **probability of E** , denoted $P(E)$ is:

$$P(E) = \frac{\text{the number of outcomes in } E}{\text{the total number of outcomes in } S}$$

- Or, given $N(A)$, the number of elements in some set A :

$$P(E) = \frac{N(E)}{N(S)}$$

Counting the Elements of a List

- Sometimes, counting list elements can be easy, such as listing the number of elements from one integer through another.
- However, there is an exact formula for this, which is helpful for determining the number of iterations with variables.

Theorem 9.1.1:

If m and n are integers such that $m \leq n$, then there are $n - m + 1$ integers from m to n , inclusive.

Example: Counting the Elements of a Sublist

How many three-digit integers exist are divisible by 5?

- For each three-digit integer divisible by 5, there is a corresponding integer from 20 to 199, inclusive.
- Hence, by **Theorem 9.1.1.**, there are $199 - 20 + 1 = 180$ three-digit integers divisible by 5.

What is the probability that a randomly chosen three-digit integer is divisible by 5?

- By **Theorem 9.1.1.**, there are $999 - 100 + 1 = 900$ total three-digit integers.
- Using the previous answer, we can now calculate the probability:

$$\frac{180}{900} = \frac{1}{5} \text{ by algebra}$$

9.2 Possibility Trees and the Multiplication Rule

- Tree structures are useful for keeping track of all the possibilities in situations where events happen in a particular order.

Example: Tournament Play Possibilities

Suppose there are two teams, *A* and *B*, who continuously play against each other until one team wins twice in a row or wins three times total. A series of wins may be denoted by the team names, like *A, B, B*.

How many ways can the tournament be played?

- Using a tree, we can outline all the possibilities up to 5 levels:




Diagram 9.2.1: The possibility tree for the tournament.

- As outlined in the possibility tree, there are ten different ways the tournament can be played, with *A* and *B* each having five different events where they win.

If each outcome is assumed to be equally likely, what is the likelihood that a team wins after five games?

- There are 4 leaf nodes on the tree, each representing a win on the fifth game.
- Thus, the probability is as follows:

$$\frac{4}{10} = \frac{1}{5} \text{ by algebra}$$

$$= 40\% \text{ by converting to percent}$$

The Multiplication Rule

- Imagine a computer installation with four IO units *A*, *B*, *C*, and *D*, and three CPUs *X*, *Y*, and *Z*.
- There should be twelve ways to pair the IO units with the CPUs.
- Note that there are CPU options available for each IO unit, and there are four IO units.
- Thus, there are $4 \cdot 3 = 12$ pairs.

Theorem 9.2.1:

If an operation consists of k steps, and if every i th step for each integer $1 \leq i \leq k$ can be completed in n_i ways, then the entire operation may be completed in $n_1 \cdot n_2 \cdots n_k$ ways.

Example: Multiplication Rule with Nested Loops

How many times does the following nested loop run?

```
for i in range(4):  
    for j in range(3):  
        #Loop body
```

- The outer loop has four iterations.
- For each iteration in the outer loop, the inner loop has three iterations.
- By the multiplication rule, the nested loop has $4 \cdot 3 = 12$ iterations.

Permutations

- A **permutation** of a set of objects is a particular ordering of the objects in a row.
- For instance, the set $\{a, b, c\}$ has 6 permutations:
- Note that for each successive step, there is one less way to perform it.
- Thus, by the multiplication rule, there are $n \cdot (n - 1) \cdot (n - 2) \cdots 2 \cdot 1$ ways to perform the entire operation.
 - Note that this matches the recursive definition for factorial.

Theorem 9.2.6:

For any integer $n \geq 1$, a set with n elements has $n!$ permutations.

Example: Possible License Plates

Suppose there is a license plate format starting with three uppercase Latin alphabet letters and ending with four digits.

How many license plates are possible?

- This problem can be split into two parts.
 - There are 26 letters in the Latin alphabet, thus there are $26^3 = 17576$ possible letter combinations.
 - There are 10 Arabic numerals, so there are $10^4 = 10000$.
- Hence, there are $17576 \cdot 10000 = 175,760,000$ possible license plates.

How many possible license plates start with an A and end with a 0?

- Because there is only 1 possible option for the first and last character of the license plate, there are only $1 \cdot 26^2 \cdot 10^3 \cdot 1 = 676,000$ possible combinations.

Continued on next page

Example: Possible License Plates continued

How many possible license plates have distinct letters and digits?

- Again, this problem can be split into two parts.
 - For each successive letter, there is one less option because they are distinct. Thus, there are $26 \cdot 25 \cdot 24 = 15600$ possible letter combinations.
 - Likewise, there is one less option for each successive digit. Thus, there are $10 \cdot 9 \cdot 8 \cdot 7 = 5040$ possible digit combinations.
- Hence, there are $15600 \cdot 5040 = 78,624,000$ possible combinations.

Permutations of Selected Elements

- Again, consider the set $\{a, b, c\}$.
- There are six different ways to select two different letters from the set and write them in a particular order:

$$ab, ac, ba, bc, ca, cb$$

- Altogether, these orderings are the 2-permutations of $\{a, b, c\}$.

Definition:

An **r -permutation** for a set of n elements is an ordered selection of r elements from the set. The number of r -permutations in a set of n elements is denoted $P(n, r)$.

Theorem 9.2.3:

If n and r are integers such that $1 \leq r \leq n$, the number of r -permutations of a set of n elements is given by the formula:

$$P(n, r) = n(n-1)(n-2)\cdots(n-r+1)$$

or

$$P(n, r) = \frac{n!}{(n-r)!}$$

- Logically, given a set of length n , the number of n -permutations in the set is equal to the number of permutations, or just $n!$.

9.3 Counting Elements of Disjoint Sets: The Addition Rule

- We may count the number elements in unions, intersections, and differences between sets.
- The number of elements in a union of mutually disjoint finite sets equals the sum of the number of elements in each of the operands.

Theorem 9.3.1:

Suppose a finite set A equals the union of k distinct mutually disjoint sets A_1, A_2, \dots, A_k . Thus,

$$N(A) = N(A_1) + N(A_2) + \dots + N(A_k)$$

- Logically, this only applies to mutually disjoint finite sets because overlapping elements are not counted separately.

Example: Possible Passwords of Different Lengths

Suppose a company has the following password restrictions:

1. Password lengths must be from 8 to 10 characters in length.
2. The 26 letters of the Latin alphabet, lowercase or uppercase.
3. The 10 Arabic numerals.
4. The 14 symbols, !, @, #, \$, %, ^, &, *, (,), -, +, {, and }.

How many passwords are possible?

- There are $26 \cdot 2 + 10 + 14 = 76$ valid symbols and three valid lengths, 3, 4, and 5.
- Thus, there are $76^3 + 76^4 + 76^5 = 6,514,592,610,973,974,528$ possible passwords.

The Difference Rule

- As a consequence of the addition rule, if the number of elements in set A and its subset B are known, then the number of elements in $A - B$ may be computed.

Theorem 9.3.2:

If A is a finite set and $B \subseteq A$, then:

$$N(A - B) = N(A) - N(B)$$

- Additionally, there is a related formula for the probability of an event's complement:

If S is a finite sample space, and A is an event in S , then

$$P(A^c) = 1 - P(A)$$

Example: Possible Passwords with Repeated Characters

Following the same restrictions as the previous example, how many passwords have at least one repeated symbol?

- First, we must calculate the number of passwords without repeated characters.
- Because the amount of possible characters decreases for each character, there are $76 \cdot 75 \cdot 74 \cdots 69 + 76 \cdot 75 \cdot 74 \cdots 68 + 76 \cdot 75 \cdot 74 \cdots 67 = 3,516,245,169,066,000,000$ total combinations.
- Because this event is a subset of the total number of passwords, we can subtract it to find the possible passwords with at least one repeated symbol:

$$\begin{aligned} &6,514,592,610,973,974,528 - 3,516,245,169,066,000,000 \\ &= 2,998,347,441,907,974,528 \text{ passwords with repeated characters} \end{aligned}$$

The Inclusion/Exclusion Rule

- Recall that the addition rule is for unions of sets that are *mutually disjoint*.
- Thus, to count the number of elements in the union of sets with overlapping elements, we must subtract the number of overlapping elements after applying the addition rule.
 - This is because overlapping elements will be counted more than once.

Theorem 9.3.3:

If A , b , and C are any finite sets, then

$$N(A \cup B) = N(A) + N(B) - N(A \cap B)$$

and

$$N(A \cup B \cup C) = N(A) + N(B) + N(C) - N(A \cap B) - N(A \cap C) - N(B \cap C) + N(A \cap B \cap C)$$

9.4 The Pigeonhole Principle

- According to the **pigeonhole principle**, given $m < n$, if n pigeons fly into m holes, then at least one hole will contain more than one pigeon.
- In terms of functions, a function from one finite set to a smaller finite set cannot be one-to-one. That is, there must be at least two elements in the domain that have the same image in the co-domain.

Example: Applications of the Pigeonhole Principle

In a group of six people, must there be at least two people who were born in the same month?

- No, there are twelve possible months, but only six people.

What about in a group of thirteen people?

- Yes; equating the people to pigeons and the birth months to pigeonholes, it can be seen that a birth month must be shared between at least two people because there are more “pigeons” than “pigeonholes.”

- Another important consequence of the pigeonhole principle is that the decimal expansion of any rational number either terminates or repeats.

Generalized Pigeonhole Principle

- A generalization of the pigeonhole principle states that if n pigeons fly into m pigeonholes, and for some positive integer k , $km < n$, then at least one pigeonhole contains at least $k + 1$ pigeons.
- Formally, for any function f from a finite set X with n elements to a finite set Y with m elements, and for any positive integer k , if $km < n$, there is some $y \in Y$ such that y is the image of at least $k + 1$ distinct elements of X .
- Or, in the contrapositive form, for any function f from a finite set X with n elements to a finite set Y with m elements, and for any positive integer k , if for each $y \in Y$, $f^{-1}(y)$ has at most k elements, then X has at most km elements; essentially, $n \leq km$.




Diagram 9.4.3: Given 5 pigeons and 2 pigeon holes, at least one pigeonhole should have 3 pigeons by the generalization of the pigeonhole principle.

9.5 Counting Subsets of a Set: Combinations

Definition:

Let n and r be nonnegative integers such that $r \leq n$. An r -**combination** of a set of n elements is a subset of r of the n elements.

- Additionally, recall the notation n choose r , the number of subsets of size r or r -combinations that can be formed from a set with n elements, denoted as

$$\binom{n}{r} = C(n, r) = \frac{n!}{r!(n-r)!}$$

- Consequently, there are two distinct methods to select r objects from a set of n elements.
 - In an **ordered selection**, both the elements chosen and their order matters. Thus, two ordered selections are the same if they contain the same elements in the same order.
 - In an **unordered selection**, only the identity of the elements chosen matter. Thus, two unordered selections are the same if they consist of the same elements.
- Notice how ordered selections are akin to r -**permutations** while the latter is akin to r -**combinations**.

Example: Relation between n Choose r and r -Permutations

n choose r and r -permutations have very similar formulas and only differ by their consideration of order. To illustrate the methodology behind their formulas, we may start by finding the number of 2-permutations of $\{6, 7, 8, 9\}$.

- Using the r -permutations formula, there are:

$$\frac{4!}{(4-2)!} = \frac{4!}{2!} = \frac{24}{2} = 12 \text{ 2-permutations}$$

- Thinking about how n choose r is an unordered selection and how r -permutations is an ordered selection, we can split the process of finding the r -Permutations into two steps:
 1. Find the r -combinations.
 2. Find the permutations of those r -combinations.
- Logically, it follows that there are $r!$ permutations for each r -combination.
- Thus, by the multiplication rule, there are $r!$ times as many r -permutations as there are r -combinations.
- In this case, there are $\frac{12}{2!} = \frac{12}{2} = 6$ r -combinations.
- Using a possibility tree, this is even easier to outline.

Continued on next page

Example: *Relation between n Choose r and r -Permutations continued*




Diagram 9.5.5: *The possibility tree for the r -combinations and r -permutations of $\{6, 7, 8, 9\}$. Notice how the r -permutations branch off of the r -combinations as the order varies.*

Theorem 9.5.1:

The number of subsets of size r or r -combinations that can be chosen from a set of n elements, $\binom{n}{r}$, is given by

$$\binom{n}{r} = \frac{P(n, r)}{r!}$$

or

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}$$

Example: Calculating the Number of Teams

Suppose there is a need to create a five person team from a pool of twelve people. How many distinct five person teams can be created?

- In terms of sets, we are looking for the number of distinct subsets of length 5 of a set of length 12.
- Thus, the number of distinct teams can be calculated using 12 choose 5:

$$\begin{aligned}\binom{12}{5} &= \frac{12!}{5!(12-5)!} \\ &= \frac{12!}{5!(7!)} \\ &= \frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8 \cdot 7!}{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1 \cdot 7!} \text{ by recursive definition of !} \\ &= \frac{12 \cdot 11 \cdot 10 \cdot 9 \cdot 8}{5 \cdot 4 \cdot 3 \cdot 2} \text{ by canceling } 7! \\ &= \frac{12 \cdot 11 \cdot 10 \cdot 9}{5 \cdot 3} \text{ by canceling } 4 \cdot 2 \text{ and } 8 \\ &= 12 \cdot 11 \cdot 2 \cdot 3 \text{ by simplifying } \frac{10}{5} \text{ and } \frac{9}{3} \\ &= 792 \text{ distinct teams}\end{aligned}$$

- However, n choose r may be involved in finding the r -permutations if *some* elements in the ordered selection do not consider order.

Theorem 9.5.2:

Suppose a collection consists of n objects with n_k objects for each k th type, and that each type is distinct, but also indistinguishable from other objects of the same type. Additionally, suppose that $n_1 + n_2 + \cdots + n_k = n$.

Then, the number of distinguishable permutations of the n objects is given by

$$\begin{aligned}\binom{n}{n_1} \binom{n-n_1}{n_2} \binom{n-n_1-n_2}{n_3} \cdots \binom{n-n_1-n_2-\cdots-n_{k-1}}{n_k} \\ = \frac{n!}{n_1! n_2! \cdots n_k!}\end{aligned}$$

Double Counting

- As displayed in the inclusion/exclusion rule for addition, there has to be consideration for overlapping elements.
- Suppose we are attempting to create a team of five containing at least one man from a pool of five men and seven women.
 - You might be tempted to choose a subset of one man from the group of five men, and then choose a subset of four people from the remaining eleven people. Then, by the multiplication rule:

$$\binom{5}{1} \cdot \binom{11}{4} = 1650 \text{ five person teams}$$

- The problem with this logic is that the separation of two unordered selection means that teams of the same people may be counted multiple times for each person pulled from the 5 choose 1 instead of the 11 choose 4.

9.6 r -Combinations with Repetition Allowed

- Previously, the formula we used did not account for combinations where elements from the set of length n were chosen multiple times.
- To understand the logic behind the formula,

Theorem 9.6.1:

The number of r -combinations with repetition allowed that can be selected from a set of n elements is

$$\binom{r+n-1}{r}$$

9.7 Pascal's Formula and the Binomial Theorem