

# $\alpha$ -Hölder smoothness

Chanwoo Lee, August 12, 2021

For one variate function, many papers follow definition of  $\alpha$ -Hölder class in [1].

**Definition 1** (Univariate case [1]). A function  $f: \mathcal{D} \rightarrow \mathbb{R}$  is in Hölder class if

1.  $\sup_{x \in \mathcal{D}} |f^{(j)}(x)| \leq \infty$ , for all  $j = 0, 1, \dots, \lfloor \alpha \rfloor$ .
2.  $\sup_{x \neq x' \in \mathcal{D}} \frac{|f^{(\lfloor \alpha \rfloor)}(x) - f^{(\lfloor \alpha \rfloor)}(x')|}{|x - x'|^{\alpha - \lfloor \alpha \rfloor}} \leq \infty$ .

Define  $\alpha$ -Hölder norm

$$\|f\|_{\mathcal{H}_\alpha} = \max_{j \leq \lfloor \alpha \rfloor} \sup_{x \in \mathcal{D}} |f^{(j)}(x)| + \sup_{x \neq x' \in \mathcal{D}} \frac{|f^{(\lfloor \alpha \rfloor)}(x) - f^{(\lfloor \alpha \rfloor)}(x')|}{|x - x'|^{\alpha - \lfloor \alpha \rfloor}}.$$

Notice we have equivalent definition of Hölder class with Definition 1 using  $\alpha$ -Hölder norm.

**Definition 1\*** (Variation for univariate case). A function  $f: \mathcal{D} \rightarrow \mathbb{R}$  is in Hölder class if

$$\|f\|_{\mathcal{H}_\alpha} \leq M,$$

for some  $M > 0$ .

Now consider multivariate function  $f: \mathcal{X} \rightarrow \mathbb{R}$ , where  $\mathcal{X} = \mathcal{D}^d$ . For multi-index  $\kappa$ , we denote partial derivatives,

$$\nabla_\kappa f(x) = \frac{\partial^{|\kappa|} f(x)}{(\partial x)^\kappa}.$$

Then we define multivariate version of  $\alpha$ -Hölder class as in [2],

**Definition 2** (Multivariate case [2]). Define  $\alpha$ -Hölder norm with respect to norm  $\|\cdot\|$ ,

$$\|f\|_{\mathcal{H}_\alpha} = \max_{j: |j| \leq \lfloor \alpha \rfloor} \sup_{\mathbf{x} \in \mathcal{X}} |\nabla_j f(\mathbf{x})| + \max_{\kappa = \lfloor \alpha \rfloor} \sup_{\mathbf{x} \neq \mathbf{x}' \in \mathcal{D}} \frac{|\nabla_\kappa f(\mathbf{x}) - \nabla_\kappa f(\mathbf{x}')|}{\|\mathbf{x} - \mathbf{x}'\|^{\alpha - \lfloor \alpha \rfloor}}.$$

Then  $f$  is in  $\alpha$ -Hölder class if  $\|f\|_{\mathcal{H}_\alpha} \leq M$ , for some  $M > 0$ .

**Remark 1.** In [2], they choose to use  $\|\cdot\|$  as zero norm when they define  $\alpha$ -Hölder norm.

**Remark 2.** In a relation to  $\alpha$ -Hölder class, we define  $\alpha$ -Hölder smooth function. We will show that  $\alpha$ -Hölder class is included in a collection of  $\alpha$ -Hölder smooth functions in Lemma 1.

**Definition 3** ( $\alpha$ -Hölder smooth). A function  $f: \mathcal{X} \rightarrow \mathbb{R}$  is  $\alpha$ -Hölder smooth with respect to  $\|\cdot\|$  if there exists a polynomial  $P_k(\cdot - \mathbf{x}_0)$  of degree  $k = \lfloor \alpha \rfloor$ , such that

$$|f(\mathbf{x}) - P_k(\mathbf{x} - \mathbf{x}_0)| \leq c \|\mathbf{x} - \mathbf{x}_0\|^\alpha, \text{ for all } \mathbf{x}, \mathbf{x}_0 \in \mathcal{X}.$$

**Lemma 1.** If a function  $f$  is in  $\alpha$ -Hölder class, then  $f$  is a  $\alpha$ -Hölder smooth.

*Proof.* Let  $P_k(\cdot - \mathbf{x}_0)$  be the Taylor polynomial of degree  $\lfloor \alpha \rfloor$ ,

$$P_k(\cdot - \mathbf{x}_0) = \sum_{\kappa: |\kappa| \leq \lfloor \alpha \rfloor} \frac{\nabla_{\kappa} f(\mathbf{x}_0)}{\kappa!} (\mathbf{x} - \mathbf{x}_0)^{\kappa}.$$

Then,  $P_k(\cdot - \mathbf{x}_0)$  satisfies,

$$\begin{aligned} |f(\mathbf{x}) - P_k(\mathbf{x} - \mathbf{x}_0)| &= \sum_{\kappa: |\kappa| = \lfloor \alpha \rfloor} \frac{|\nabla_{\kappa} f(\mathbf{z}) - \nabla_{\kappa} f(\mathbf{x}_0)|}{\kappa!} (\mathbf{x} - \mathbf{x}_0)^{\kappa}, \text{ where } \mathbf{z} = \mathbf{x}_0 + c(\mathbf{x} - \mathbf{x}_0), c \in (0, 1), \\ &\lesssim \sum_{\kappa: |\kappa| = \lfloor \alpha \rfloor} \frac{|\nabla_{\kappa} f(\mathbf{z}) - \nabla_{\kappa} f(\mathbf{x}_0)|}{\kappa!} \|\mathbf{x} - \mathbf{x}_0\|^{\lfloor \alpha \rfloor} \\ &\leq \sum_{\kappa: |\kappa| = \lfloor \alpha \rfloor} \frac{\|\mathbf{x} - \mathbf{x}_0\|^{\alpha - \lfloor \alpha \rfloor}}{\kappa!} \|\mathbf{x} - \mathbf{x}_0\|^{\lfloor \alpha \rfloor} \\ &\leq M_{\alpha} \|\mathbf{x} - \mathbf{x}_0\|^{\alpha}. \end{aligned}$$

□

## References

- [1] Lawrence C Evans. Partial differential equations and monge-kantorovich mass transfer. *Current developments in mathematics*, 1997(1):65–126, 1997.
- [2] Chao Gao, Yu Lu, and Harrison H Zhou. Rate-optimal graphon estimation. *The Annals of Statistics*, 43(6):2624–2652, 2015.