

# Compatibility between kernel SMM and SVM

Miaoyan Wang, May 18, 2020

**Theorem 0.1** (Compatibility with classical SVM). *Let  $X \in \mathbb{R}^d$  denote a column vector and  $X^* = \begin{bmatrix} 0 & X^T \\ X & 0 \end{bmatrix} \in \mathbb{R}^{(d+1) \times (d+1)}$  be the symmetrized matrix. Then, under our proposed column-wise kernels, the SMM classifier trained on  $X^*$  equals the SVM classifier trained on  $X$ .*

**Remark 1.** The decision boundaries generated by two classifiers are the same, but the optimal objective value may not be the same.

*Proof.* We prove the following stronger result.

**(Set-up.)** Let  $X \in \mathbb{R}^{m \times n}$  be the original matrix feature,

$$X = \begin{bmatrix} | & | & | & | \\ x_1 & x_2 & \vdots & x_n \\ | & | & | & | \end{bmatrix} = \begin{bmatrix} - & y_1^T & - \\ - & y_2^T & - \\ - & \dots & - \\ - & y_m^T & - \end{bmatrix},$$

where  $x_i$  and  $y_j^T$  denotes the  $i$ -th column and  $j$ -th row of the matrix  $X$ , respectively. Let  $X^* \in \mathbb{R}^{(m+n) \times (m+n)}$  be the symmetrized matrix feature. Based on the assumption (of sufficiently valid kernels), the feature mapping  $h$  is applied to matrix  $X^*$  in a column-wise fashion; that is,

$$X^* = \begin{bmatrix} 0 & X^T \\ X & 0 \end{bmatrix} \Rightarrow h(X^*) = \begin{bmatrix} & & & & | & | & | \\ & & & & h(y_1^T) & \dots & h(y_m^T) \\ & & & & | & | & | \\ \dots & & & & \dots & \dots & \dots \\ h(x_1) & \dots & h(x_n) & & & & \\ | & | & | & & & & 0 \end{bmatrix} =: \begin{bmatrix} 0 & A \\ C & 0 \end{bmatrix}, \quad (1)$$

where, without a little abuse of notation, we have used  $h(x_i)$  and  $h(y_i^T)$  to denote the non-zero coordinates in the image vector  $h((0, \dots, 0, x_i^T))$  and  $h((y_i^T, 0, \dots, 0))$ , respectively.

**(Conclusion.)** Assume the set of elements in  $C$  contains all elements in  $A$ . Then, the classifier trained on  $X$  equals to the classifier trained on  $X^*$ .

**(Proof of the above conclusion.)** Because the elements in  $C$  contains all elements in  $A$ , we re-arrange the elements in  $C$  into a possibly larger matrix:

$$C = \begin{bmatrix} 0 & A^T \\ D & 0 \end{bmatrix}. \quad (2)$$

Such reshaping is possible by, for example, setting  $D = \text{diag}(\{C\}/\{A\})$ . (The choice of  $D$  is non-important.) To prove the equivalence between  $X$ - and  $X^*$ -trained classifiers, it suffices to prove the equivalence between

$$h(X^*) = \begin{bmatrix} 0 & 0 & A \\ 0 & A^T & 0 \\ D & 0 & 0 \end{bmatrix}, \quad \text{and} \quad h(X) = \begin{bmatrix} 0 & A^T \\ D & 0 \end{bmatrix} \quad (\text{c.f. (1) and (2)}).$$

Now we apply the same proof techniques as in 051820\_SMMK\_modification.pdf. Specifically, consider the primal problems with  $h(X^*)$  and  $h(X)$ . The relevant linear predictors are

$$\langle B^*, h(X^*) \rangle = \left\langle \begin{bmatrix} 0 & 0 & B_3^* \\ 0 & B_2^* & 0 \\ B_1^* & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & A \\ 0 & A^T & 0 \\ D & 0 & 0 \end{bmatrix} \right\rangle, \quad \text{and} \quad \langle B, h(X) \rangle = \left\langle \begin{bmatrix} 0 & B_2 \\ B_1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & A^T \\ D & 0 \end{bmatrix} \right\rangle.$$

The optimizations are

$$(P1) \quad \min_{B_1^*, B_2^*, B_3^*} \|B_1^*\|_F^2 + \|B_2^*\|_F^2 + \|B_3^*\|_F^2 + C \sum_i \xi_i, \\ \text{s.t. .... where } \langle B^*, h(X^*) \rangle = \langle B_1^*, D \rangle + \langle B_2^*, A^T \rangle + \langle B_3^{*T}, A \rangle \\ = \langle B_1^*, D \rangle + \langle B_2^{*T} + B_3^*, A \rangle,$$

and

$$(P2) \quad \min_{B_1, B_2} \|B_1\|_F^2 + \|B_2\|_F^2 + C \sum_i \xi_i, \\ \text{s.t. ... where } \langle B, h(X) \rangle = \langle B_1, D \rangle + \langle B_2, A \rangle.$$

It is easy to see that the optimal solution to (P1) is achieved at  $B_2^* = B_3^{*T}$  because

$$\|B_2^*\|_F^2 + \|B_3^*\|_F^2 \geq \frac{1}{2} \|B_2^* + B_3^*\|_F^2 = \frac{1}{2} \|B_2^*\|_F^2,$$

where the equality holds when  $B_2^{*T} = B_3^*$ . Moreover, under this choice, the optimal objective values

$$(P1) \quad \min_{B_1^*, B_2^*} \|B_1\|_F + \frac{1}{2} \|B_2\|_F^2 + C \sum_i \xi_i, \quad (P2) \quad \min_{B_1, B_2} \|B_1\|_F + \|B_2\|_F^2 + C \sum_i \xi_i,$$

are not necessarily equal unless  $B_1 = 0$ . □