

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity**, **equivalence**, **entailment** can be reduced to satisfiability checking.

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity**, **equivalence**, **entailment** can be reduced to satisfiability checking.

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity**, **equivalence**, **entailment** can be reduced to satisfiability checking.

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity**, **equivalence**, **entailment** can be reduced to satisfiability checking.

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity**, **equivalence**, **entailment** can be reduced to satisfiability checking.

# Propositional Satisfiability Problem

Given a propositional formula  $A$ , check whether it is **satisfiable** or **unsatisfiable**.

If  $A$  is satisfiable, we also want to find a **satisfying assignment** for  $A$ , that is, a **model** of  $A$ .

It is one of the **most famous** combinatorial problems in computer science.

It is a **very hard** problem with a surprisingly **large number of practical applications**.

It is also the first ever problem to be proved **NP-complete**.

Checking **validity, equivalence, entailment** can be reduced to satisfiability checking.

# Russian spy puzzle



There are three persons: Stirlitz, Müller, and Eismann. It is known that **exactly one** of them is Russian, while the **other two** are Germans. Moreover, **every Russian must be a spy**.

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, **you are as German as I am Russian**”. It is known that Stirlitz always tells the truth when he is joking.

**We have to show that Eismann is not a Russian spy.**

How can we solve problems of this kind?

# Russian spy puzzle



There are three persons: Stirlitz, Müller, and Eismann. It is known that **exactly one** of them is Russian, while the **other two** are Germans. Moreover, **every Russian must be a spy**.

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, **you are as German as I am Russian**”. It is known that Stirlitz always tells the truth when he is joking.



We have to show that Eismann is not a Russian spy.

How can we solve problems of this kind?



# Russian spy puzzle



There are three persons: Stirlitz, Müller, and Eismann. It is known that **exactly one** of them is Russian, while the **other two** are Germans. Moreover, **every Russian must be a spy**.

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, **you are as German as I am Russian**”. It is known that Stirlitz always tells the truth when he is joking.



**We have to show that Eismann is not a Russian spy.**

How can we solve problems of this kind?

# Russian spy puzzle



There are three persons: Stirlitz, Müller, and Eismann. It is known that **exactly one** of them is Russian, while the **other two** are Germans. Moreover, **every Russian must be a spy**.

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, **you are as German as I am Russian**”. It is known that Stirlitz always tells the truth when he is joking.



**We have to show that Eismann is not a Russian spy.**

**How can we solve problems of this kind?**

# Formalisation in propositional logic

Introduce nine propositional variables as in the following table:

	Stirlitz	Müller	Eismann
Russian	RS	RM	RE
German	GS	GM	GE
Spy	SS	SM	SE

For example,

*SE* : Eismann is a Spy

*RS* : Stirlitz is Russian

# Formalisation in propositional logic

Introduce nine propositional variables as in the following table:

	Stirlitz	Müller	Eismann
Russian	RS	RM	RE
German	GS	GM	GE
Spy	SS	SM	SE

For example,

*SE* : Eismann is a Spy

*RS* : Stirlitz is Russian

# Formalisation in propositional logic

There are three persons: **Stirlitz**, **Müller**, and **Eismann**. It is known that exactly one of them is **Russian**, while the other two are **Germans**.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every **Russian** must be a **spy**.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When **Stirlitz** meets **Müller** in a corridor, he makes the following joke: “you know, **Müller**, you are as **German** as I am **Russian**”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that **Eismann** is not a **Russian spy**.

# Formalisation in propositional logic

There are three persons: Stirlitz, Müller, and Eismann. It is known that exactly one of them is Russian, while the other two are Germans.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every Russian must be a spy.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, you are as German as I am Russian”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that Eismann is not a Russian spy.

# Formalisation in propositional logic

There are three persons: Stirlitz, Müller, and Eismann. It is known that exactly one of them is Russian, while the other two are Germans.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every Russian must be a spy.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, you are as German as I am Russian”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that Eismann is not a Russian spy.

# Formalisation in propositional logic

There are three persons: Stirlitz, Müller, and Eismann. It is known that exactly one of them is Russian, while the other two are Germans.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every Russian must be a spy.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, you are as German as I am Russian”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that Eismann is not a Russian spy.



# Formalisation in propositional logic

There are three persons: Stirlitz, Müller, and Eismann. It is known that exactly one of them is Russian, while the other two are Germans.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every Russian must be a spy.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, you are as German as I am Russian”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that Eismann is not a Russian spy.

# Formalisation in propositional logic

There are three persons: Stirlitz, Müller, and Eismann. It is known that exactly one of them is Russian, while the other two are Germans.

$$(RS \wedge GM \wedge GE) \vee (GS \wedge RM \wedge GE) \vee (GS \wedge GM \wedge RE).$$

Moreover, every Russian must be a spy.

$$(RS \rightarrow SS) \wedge (RM \rightarrow SM) \wedge (RE \rightarrow SE).$$

When Stirlitz meets Müller in a corridor, he makes the following joke: “you know, Müller, you are as German as I am Russian”.

$$RS \leftrightarrow GM.$$

Hidden: Russians are not Germans.

$$(RS \leftrightarrow \neg GS) \wedge (RM \leftrightarrow \neg GM) \wedge (RE \leftrightarrow \neg GE).$$

We have to show that Eismann is not a Russian spy.


To this end, we add the following formula

$$RE \wedge SE.$$

and check whether the resulting set of formulas is satisfiable. If it is unsatisfiable, then Eismann cannot be a Russian spy.

# Circuit Equivalence


Given two circuits, check if they are equivalent. For example:



We know that equivalence-checking for propositional formulas can be reduced to unsatisfiability-checking.

# Circuit Equivalence


Given two circuits, check if they are equivalent. For example:



We know that equivalence-checking for propositional formulas can be reduced to unsatisfiability-checking.

# Circuit Equivalence

Given two circuits, check if they are equivalent. For example:



We know that equivalence-checking for **propositional formulas** can be **reduced to unsatisfiability-checking**.

# Satisfiability: use formula evaluation methods

Consider  $\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$ .

We can evaluate it in any interpretation, for example,

$\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ :

	subformula				$I_0$
1	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$				0
2	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$				1
3	$p \rightarrow r$				1
4	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$				1
5	$p \wedge q \rightarrow r$				1
6	$p \rightarrow q$				1
7	$p \wedge q$				0
8	$p$	$p$	$p$		0
9	$q$	$q$			0
10			$r$	$r$	0

# Satisfiability: Truth tables

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)).$$

Likewise, we can evaluate it in **all** interpretations:

	subformula				$l_1$	$l_2$	$l_3$	$l_4$	$l_5$	$l_6$	$l_7$	$l_8$
1	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$				0	0	0	0	0	0	0	0
2	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$				1	1	1	1	1	1	1	1
3	$p \rightarrow r$				1	1	1	1	0	1	0	1
4	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$				1	1	1	1	0	0	0	1
5	$p \wedge q \rightarrow r$				1	1	1	1	1	1	0	1
6	$p \rightarrow q$				1	1	1	1	0	0	1	1
7	$p \wedge q$				0	0	0	0	0	0	1	1
8	$p$	$p$		$p$	0	0	0	0	1	1	1	1
9		$q$	$q$		0	0	1	1	0	0	1	1
10				$r$	0	1	0	1	0	1	0	1

The formula is **unsatisfiable** since it is false in every interpretation.

**Problem:** a formula with  $n$  propositional variables has  $2^n$  different interpretations.

# Satisfiability: Truth tables

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)).$$

Likewise, we can evaluate it in **all** interpretations:

	subformula				$l_1$	$l_2$	$l_3$	$l_4$	$l_5$	$l_6$	$l_7$	$l_8$
1	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$				0	0	0	0	0	0	0	0
2	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$				1	1	1	1	1	1	1	1
3	$p \rightarrow r$				1	1	1	1	0	1	0	1
4	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$				1	1	1	1	0	0	0	1
5	$p \wedge q \rightarrow r$				1	1	1	1	1	1	0	1
6	$p \rightarrow q$				1	1	1	1	0	0	1	1
7	$p \wedge q$				0	0	0	0	0	0	1	1
8	$p$	$p$		$p$	0	0	0	0	1	1	1	1
9		$q$	$q$		0	0	1	1	0	0	1	1
10				$r$	0	1	0	1	0	1	0	1

The formula is **unsatisfiable** since it is false in every interpretation.

**Problem:** a formula with  $n$  propositional variables has  $2^n$  different interpretations.



# Satisfiability: Truth tables

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)).$$

Likewise, we can evaluate it in **all** interpretations:

	subformula				$l_1$	$l_2$	$l_3$	$l_4$	$l_5$	$l_6$	$l_7$	$l_8$
1	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$				0	0	0	0	0	0	0	0
2	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$				1	1	1	1	1	1	1	1
3	$p \rightarrow r$				1	1	1	1	0	1	0	1
4	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$				1	1	1	1	0	0	0	1
5	$p \wedge q \rightarrow r$				1	1	1	1	1	1	0	1
6	$p \rightarrow q$				1	1	1	1	0	0	1	1
7	$p \wedge q$				0	0	0	0	0	0	1	1
8	$p$	$p$	$p$	$p$	0	0	0	0	1	1	1	1
9	$q$	$q$	$q$	$q$	0	0	1	1	0	0	1	1
10	$r$	$r$	$r$	$r$	0	1	0	1	0	1	0	1

The formula is **unsatisfiable** since it is false in every interpretation.

**Problem:** a formula with  $n$  propositional variables has  $2^n$  different interpretations.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

## Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$			0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

## Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

## Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$			0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.



# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$			0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$			0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

## Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is **unsatisfiable**.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

# Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$	$p$	$p$	$p$		0	1	1	
	$q$	$q$				0	1	
			$r$	$r$	0	0	0	1

The formula is unsatisfiable.

**Note:** the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.

## Compact truth table

Idea: we can sometimes evaluate a formula based on values of only a **subset of all variables**.

subformula					$l_2$	$l_3$	$l_4$	$l_1$
$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$					0	0	0	0
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$					1	1	1	1
$p \rightarrow r$					1	0	0	1
$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$						0	0	
$p \wedge q \rightarrow r$						1	0	1
$p \rightarrow q$						0	1	
$p \wedge q$						0	1	
$p$		$p$		$p$	0	1	1	
	$q$		$q$			0	1	
				$r$	0	0	0	1

The formula is unsatisfiable.

Note: the size of the compact table (but not the result) depends on the order of atoms!

The ideas of **guessing variable values** (or **case analysis**) and **propagation** are the key ideas in nearly all propositional satisfiability algorithms.



## The splitting algorithm for propositional satisfiability

# Splitting: idea

$A_p^\perp$  and  $A_p^\top$ : the formulas obtained by replacing in  $A$  all occurrences of  $p$  by  $\perp$  and  $\top$ , respectively.

## Lemma

Let  $p$  be an atom,  $A$  be a formula, and  $I$  be an interpretation.

1. If  $I \models p$ , then  $I \models p \leftrightarrow \top$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\top$  in  $I$ .*

2. If  $I \not\models p$ , then  $I \models p \leftrightarrow \perp$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\perp$  in  $I$ .*

- Pick a variable  $p$  and perform case analysis on this variable:
  - If  $p$  is **true**, replace  $p$  by  $\top$ .
  - If  $p$  is **false**, replace  $p$  by  $\perp$ ;
- When a formula contains occurrences of  $\top$  or  $\perp$ , **simplify it**.

# Splitting: idea

$A_p^\perp$  and  $A_p^\top$ : the formulas obtained by replacing in  $A$  all occurrences of  $p$  by  $\perp$  and  $\top$ , respectively.

## Lemma

Let  $p$  be an atom,  $A$  be a formula, and  $I$  be an interpretation.

1. If  $I \models p$ , then  $I \models p \leftrightarrow \top$ .

By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\top$  in  $I$ .

2. If  $I \not\models p$ , then  $I \models p \leftrightarrow \perp$ .

By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\perp$  in  $I$ .

- ▶ Pick a variable  $p$  and perform case analysis on this variable:
  - ▶ If  $p$  is true, replace  $p$  by  $\top$ .
  - ▶ If  $p$  is false, replace  $p$  by  $\perp$ ;
- ▶ When a formula contains occurrences of  $\top$  or  $\perp$ , simplify it.

# Splitting: idea

$A_p^\perp$  and  $A_p^\top$ : the formulas obtained by replacing in  $A$  all occurrences of  $p$  by  $\perp$  and  $\top$ , respectively.

## Lemma

Let  $p$  be an atom,  $A$  be a formula, and  $I$  be an interpretation.

1. If  $I \models p$ , then  $I \models p \leftrightarrow \top$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\top$  in  $I$ .*

2. If  $I \not\models p$ , then  $I \models p \leftrightarrow \perp$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\perp$  in  $I$ .*

- Pick a variable  $p$  and perform case analysis on this variable:
  - If  $p$  is **true**, replace  $p$  by  $\top$ .
  - If  $p$  is **false**, replace  $p$  by  $\perp$ ;
- When a formula contains occurrences of  $\top$  or  $\perp$ , **simplify it**.

# Splitting: idea

$A_p^\perp$  and  $A_p^\top$ : the formulas obtained by replacing in  $A$  all occurrences of  $p$  by  $\perp$  and  $\top$ , respectively.

## Lemma

Let  $p$  be an atom,  $A$  be a formula, and  $I$  be an interpretation.

1. If  $I \models p$ , then  $I \models p \leftrightarrow \top$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\top$  in  $I$ .*

2. If  $I \not\models p$ , then  $I \models p \leftrightarrow \perp$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\perp$  in  $I$ .*

- Pick a variable  $p$  and perform case analysis on this variable:
  - If  $p$  is **true**, replace  $p$  by  $\top$ .
  - If  $p$  is **false**, replace  $p$  by  $\perp$ ;
- When a formula contains occurrences of  $\top$  or  $\perp$ , **simplify it**.

# Splitting: idea

$A_p^\perp$  and  $A_p^\top$ : the formulas obtained by replacing in  $A$  all occurrences of  $p$  by  $\perp$  and  $\top$ , respectively.

## Lemma

Let  $p$  be an atom,  $A$  be a formula, and  $I$  be an interpretation.

1. If  $I \models p$ , then  $I \models p \leftrightarrow \top$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\top$  in  $I$ .*

2. If  $I \not\models p$ , then  $I \models p \leftrightarrow \perp$ .

*By equivalent replacement lemma,  $A$  is equivalent to  $A_p^\perp$  in  $I$ .*

- Pick a variable  $p$  and perform case analysis on this variable:
  - If  $p$  is **true**, replace  $p$  by  $\top$ .
  - If  $p$  is **false**, replace  $p$  by  $\perp$ ;
- When a formula contains occurrences of  $\top$  or  $\perp$ , **simplify it**.

# Simplification rules for $\top$ and $\perp$

## Simplification rules for $\top$ :

$$\begin{aligned}\neg \top &\Rightarrow \perp \\ \top \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow A_1 \wedge \dots \wedge A_n \\ \top \vee A_1 \vee \dots \vee A_n &\Rightarrow \top \\ A \rightarrow \top &\Rightarrow \top & \top \rightarrow A &\Rightarrow A \\ A \leftrightarrow \top &\Rightarrow A & \top \leftrightarrow A &\Rightarrow A\end{aligned}$$

## Simplification rules for $\perp$ :

$$\begin{aligned}\neg \perp &\Rightarrow \top \\ \perp \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow \perp \\ \perp \vee A_1 \vee \dots \vee A_n &\Rightarrow A_1 \vee \dots \vee A_n \\ A \rightarrow \perp &\Rightarrow \neg A & \perp \rightarrow A &\Rightarrow \top \\ A \leftrightarrow \perp &\Rightarrow \neg A & \perp \leftrightarrow A &\Rightarrow \neg A\end{aligned}$$

Note that they cover all cases when  $\perp$  or  $\top$  occurs in the formula apart from the trivial ones.

Thus, if we apply these rules until they are no more applicable we obtain either  $\perp$ , or  $\top$ , or a formula containing neither  $\perp$  nor  $\top$ .

# Simplification rules for $\top$ and $\perp$

Simplification rules for  $\top$ :

$$\begin{aligned}\neg \top &\Rightarrow \perp \\ \top \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow A_1 \wedge \dots \wedge A_n \\ \top \vee A_1 \vee \dots \vee A_n &\Rightarrow \top \\ A \rightarrow \top &\Rightarrow \top & \top \rightarrow A &\Rightarrow A \\ A \leftrightarrow \top &\Rightarrow A & \top \leftrightarrow A &\Rightarrow A\end{aligned}$$

Simplification rules for  $\perp$ :

$$\begin{aligned}\neg \perp &\Rightarrow \top \\ \perp \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow \perp \\ \perp \vee A_1 \vee \dots \vee A_n &\Rightarrow A_1 \vee \dots \vee A_n \\ A \rightarrow \perp &\Rightarrow \neg A & \perp \rightarrow A &\Rightarrow \top \\ A \leftrightarrow \perp &\Rightarrow \neg A & \perp \leftrightarrow A &\Rightarrow \neg A\end{aligned}$$

Note that they cover **all cases** when  $\perp$  or  $\top$  occurs in the formula apart from the trivial ones.

Thus, if we apply these rules until they are no more applicable we obtain either  $\perp$ , or  $\top$ , or a formula containing neither  $\perp$  nor  $\top$ .



# Simplification rules for $\top$ and $\perp$

Simplification rules for  $\top$ :

$$\begin{aligned}\neg \top &\Rightarrow \perp \\ \top \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow A_1 \wedge \dots \wedge A_n \\ \top \vee A_1 \vee \dots \vee A_n &\Rightarrow \top \\ A \rightarrow \top &\Rightarrow \top & \top \rightarrow A &\Rightarrow A \\ A \leftrightarrow \top &\Rightarrow A & \top \leftrightarrow A &\Rightarrow A\end{aligned}$$

Simplification rules for  $\perp$ :

$$\begin{aligned}\neg \perp &\Rightarrow \top \\ \perp \wedge A_1 \wedge \dots \wedge A_n &\Rightarrow \perp \\ \perp \vee A_1 \vee \dots \vee A_n &\Rightarrow A_1 \vee \dots \vee A_n \\ A \rightarrow \perp &\Rightarrow \neg A & \perp \rightarrow A &\Rightarrow \top \\ A \leftrightarrow \perp &\Rightarrow \neg A & \perp \leftrightarrow A &\Rightarrow \neg A\end{aligned}$$


Note that they cover all cases when  $\perp$  or  $\top$  occurs in the formula apart from the trivial ones.

Thus, if we apply these rules until they are no more applicable we obtain either  $\perp$ , or  $\top$ , or a formula containing neither  $\perp$  nor  $\top$ .


# Splitting algorithm

```
procedure split( $G$ )  
parameters: function select  
input: formula  $G$   
output: “satisfiable” or “unsatisfiable”  
begin  
   $G := \text{simplify}(G)$   
  if  $G = \top$  then return “satisfiable”  
  if  $G = \perp$  then return “unsatisfiable”  
   $(p, b) := \text{select}(G)$   
  case  $b$  of  
    1  $\Rightarrow$   
      if split( $G_p^\top$ ) = “satisfiable”  
        then return “satisfiable”  
        else return split( $G_p^\perp$ )  
    0  $\Rightarrow$   
      if split( $G_p^\perp$ ) = “satisfiable”  
        then return “satisfiable”  
        else return split( $G_p^\top$ )  
end
```


## Splitting algorithm, example




## Splitting algorithm, example




## Splitting algorithm, example




## Splitting algorithm, example



# Splitting algorithm, example




## Splitting algorithm, example







## Splitting algorithm, example




# Splitting algorithm, example




# Splitting algorithm, example




# Splitting algorithm, example




## Splitting algorithm, example




# Splitting algorithm, example



# Splitting algorithm, example




# Splitting algorithm, example







# Splitting algorithm, example




# Splitting algorithm, example



## Splitting algorithm, example




## Splitting algorithm, example



The formula is **unsatisfiable**.

# Splitting algorithm, example



The formula is **unsatisfiable**.

What is going on here is very similar to using compact truth tables, but on the syntactic level.

## Splitting algorithm, example 2

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r))$$



The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r))$$

$p=0$

---

$$\neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$r=0$


---

The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2




The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .



## Splitting algorithm, example 2




The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\perp$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2




The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2




The formula is **satisfiable**.

To find a model of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2




The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Splitting algorithm, example 2



The formula is **satisfiable**.

To **find a model** of this formula, we should simply collect choices made on the branch terminating at  $\top$ .

Any interpretation  $I$  such that  $I(p) = I(r) = 0$  satisfies the formula, for example the interpretation  $\{p \mapsto 0, q \mapsto 0, r \mapsto 0\}$ .

## Next:

- ▶ **monotonicity**
- ▶ **position** of a subformula occurrence,
- ▶ **polarity** of a subformula occurrence,
- ▶ **monotonic replacement** based on polarity,
- ▶ **optimizations based on monotonic replacement:** pure atom rule.

Monotonicity, position, polarity

# Monotonicity

- ▶ Introduce an **order**  $<$  on truth values by defining  $0 < 1$  and
- ▶ A function  $f(x_1, \dots, x_n)$  is called **monotonic** on its  $k$ -th argument (w.r.t. an order  $<$ ) if  $a_k \leq a'_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ A function  $f(x_1, \dots, x_n)$  is called **anti-monotonic** on its  $k$ -th argument if  $a'_k \leq a_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ consider the behaviour of the logical connectives w.r.t. monotonicity.



# Monotonicity

- ▶ Introduce an **order**  $<$  on truth values by defining  $0 < 1$  and
- ▶ A function  $f(x_1, \dots, x_n)$  is called **monotonic** on its  $k$ -th argument (w.r.t. an order  $<$ ) if  $a_k \leq a'_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ A function  $f(x_1, \dots, x_n)$  is called **anti-monotonic** on its  $k$ -th argument if  $a'_k \leq a_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ consider the behaviour of the logical connectives w.r.t. monotonicity.

# Monotonicity

- ▶ Introduce an **order**  $<$  on truth values by defining  $0 < 1$  and
- ▶ A function  $f(x_1, \dots, x_n)$  is called **monotonic** on its  $k$ -th argument (w.r.t. an order  $<$ ) if  $a_k \leq a'_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ A function  $f(x_1, \dots, x_n)$  is called **anti-monotonic** on its  $k$ -th argument if  $a'_k \leq a_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ consider the behaviour of the logical connectives w.r.t. monotonicity.

# Monotonicity

- ▶ Introduce an **order**  $<$  on truth values by defining  $0 < 1$  and
- ▶ A function  $f(x_1, \dots, x_n)$  is called **monotonic** on its  $k$ -th argument (w.r.t. an order  $<$ ) if  $a_k \leq a'_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ A function  $f(x_1, \dots, x_n)$  is called **anti-monotonic** on its  $k$ -th argument if  $a'_k \leq a_k$  implies
$$f(a_1, \dots, a_k, \dots, a_n) \leq f(a_1, \dots, a'_k, \dots, a_n).$$
- ▶ consider the behaviour of the logical connectives w.r.t. monotonicity.

# Monotonicity and logical connectives

- ▶ The connectives  $\wedge$  and  $\vee$  are **monotonic** on **all** of their arguments.
- ▶ The negation  $\neg$  is **anti-monotonic**.
- ▶ The implication  $\rightarrow$  is **monotonic on its second argument**, but **anti-monotonic on its first argument**.
- ▶ The equivalence  $\leftrightarrow$  is **neither monotonic nor anti-monotonic** on either of its arguments.

# Monotonicity and logical connectives

- ▶ The connectives  $\wedge$  and  $\vee$  are **monotonic** on **all** of their arguments.
- ▶ The negation  $\neg$  is **anti-monotonic**.
- ▶ The implication  $\rightarrow$  is **monotonic on its second argument**, but **anti-monotonic on its first argument**.
- ▶ The equivalence  $\leftrightarrow$  is **neither monotonic nor anti-monotonic** on either of its arguments.

# Monotonicity and logical connectives


- ▶ The connectives  $\wedge$  and  $\vee$  are **monotonic** on **all** of their arguments.
- ▶ The negation  $\neg$  is **anti-monotonic**.
- ▶ The implication  $\rightarrow$  is **monotonic on its second argument**, but **anti-monotonic on its first argument**.
- ▶ The equivalence  $\leftrightarrow$  is **neither monotonic nor anti-monotonic** on either of its arguments.

# Monotonicity and logical connectives

- ▶ The connectives  $\wedge$  and  $\vee$  are **monotonic** on **all** of their arguments.
- ▶ The negation  $\neg$  is **anti-monotonic**.
- ▶ The implication  $\rightarrow$  is **monotonic on its second argument**, but **anti-monotonic on its first argument**.
- ▶ The equivalence  $\leftrightarrow$  is **neither monotonic nor anti-monotonic** on either of its arguments.

# Parse tree

$$A \stackrel{\text{def}}{=} \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$




- ▶ Position in the formula: 1.1.2.1;
- ▶ Subformula at this position:  $p \wedge q$ ; denoted  $A|_{1.1.2.1} = p \wedge q$ .
- ▶ Position of  $A$  is  $\epsilon$ .



# Parse tree


$$A \stackrel{\text{def}}{=} \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$



- ▶ Position in the formula: 1.1.2.1;
- ▶ Subformula at this position:  $p \wedge q$ ; denoted  $A|_{1.1.2.1} = p \wedge q$ .
- ▶ Position of  $A$  is  $\epsilon$ .

# Parse tree

$$A \stackrel{\text{def}}{=} \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$



- ▶ Position in the formula: 1.1.2.1;
- ▶ Subformula at this position:  $p \wedge q$ ; denoted  $A|_{1.1.2.1} = p \wedge q$ .
- ▶ Position of  $A$  is  $\epsilon$ .

# Positions and Subformulas

- **Position** is any sequence of positive integers  $a_1, \dots, a_n$ , where  $n \geq 0$ , written as  $a_1.a_2.\dots.a_n$ .
- **Empty position**, denoted by  $\epsilon$ : when  $n = 0$ .
- **Position  $\pi$  in a formula  $A$ , subformula at a position**, denoted  $A|_\pi$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$  and  $A|_\epsilon \stackrel{\text{def}}{=} A$ .

2. Let  $A|_\pi = B$ .

2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ .

2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ .

2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ;

2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ .

If  $A|_\pi = B$ , we also say that  $B$  occurs in  $A$  at the position  $\pi$ .

# Positions and Subformulas

- ▶ **Position** is any sequence of positive integers  $a_1, \dots, a_n$ , where  $n \geq 0$ , written as  $a_1.a_2.\dots.a_n$ .
- ▶ **Empty position**, denoted by  $\epsilon$ : when  $n = 0$ .
- ▶ **Position  $\pi$  in a formula  $A$ , subformula at a position**, denoted  $A|_\pi$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$  and  $A|_\epsilon \stackrel{\text{def}}{=} A$ .

2. Let  $A|_\pi = B$ .

2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all

$i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ .

2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ .

2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ;

2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ .

If  $A|_\pi = B$ , we also say that  $B$  occurs in  $A$  at the position  $\pi$ .

# Polarity

Polarity of subformula at a position. Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.



# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity

**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive** respectively.

# Polarity


**Polarity of subformula at a position.** Notation:  $pol(A, \pi)$ .

1. For every formula  $A$ ,  $\epsilon$  is a position in  $A$ ,  $A|_{\epsilon} \stackrel{\text{def}}{=} A$  and  $pol(A, \epsilon) \stackrel{\text{def}}{=} 1$ .
  2. Let  $A|_{\pi} = B$ .
    - 2.1 If  $B$  has the form  $B_1 \wedge \dots \wedge B_n$  or  $B_1 \vee \dots \vee B_n$ , then for all  $i \in \{1, \dots, n\}$  the position  $\pi.i$  is a position in  $A$ ,  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$ , and  $pol(A, \pi.i) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.2 If  $B$  has the form  $\neg B_1$ , then  $\pi.1$  is a position in  $A$ ,  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$  and  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ .
    - 2.3 If  $B$  has the form  $B_1 \rightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and we have  $A|_{\pi.1} \stackrel{\text{def}}{=} B_1$ ,  $A|_{\pi.2} \stackrel{\text{def}}{=} B_2$ ,  $pol(A, \pi.1) \stackrel{\text{def}}{=} -pol(A, \pi)$ ,  $pol(A, \pi.2) \stackrel{\text{def}}{=} pol(A, \pi)$ .
    - 2.4 If  $B$  has the form  $B_1 \leftrightarrow B_2$ , then  $\pi.1$  and  $\pi.2$  are positions in  $A$  and  $A|_{\pi.i} \stackrel{\text{def}}{=} B_i$  and  $pol(A, \pi.i) \stackrel{\text{def}}{=} 0$  for  $i = 1, 2$ .
- If  $pol(A, \pi) = 1; -1; 0$  and  $A|_{\pi} = B$ , then we call the occurrence of  $B$  at the position  $\pi$  in  $A$  **positive**; **negative**; **neutral** respectively.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is 0.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is 0.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is **0**.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.



# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is 0.
- Otherwise, its polarity is  $-1$  if it has an **odd number of red arcs** above it and  $1$  if **even**.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is **0**.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.




- If a position has **at least one blue arc** above it, its polarity is 0.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.

# The coloring algorithm for determining polarity

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \leftrightarrow (r \rightarrow q))).$$

- Color in **blue** all arcs below an equivalence.
- Color in **red** all uncolored arcs going down from a negation or left-hand side of an implication.



- If a position has **at least one blue arc** above it, its polarity is **0**.
- Otherwise, its polarity is **-1** if it has an **odd number of red arcs** above it and **1** if **even**.

## Position and polarity, again

position	subformula	polarity
$\epsilon$	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$ $(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	1

# Position and polarity, again

position	subformula	polarity
$\in$	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	$1$
$1$	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	$-1$
	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	

## Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
	$p \rightarrow q$	

# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
	$p$	



# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
1.1.1.1	$p$	-1
	$q$	

# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
1.1.1.1	$p$	-1
1.1.1.2	$q$	1
	$p \wedge q \rightarrow r$	

# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
1.1.1.1	$p$	-1
1.1.1.2	$q$	1
1.1.2	$p \wedge q \rightarrow r$	1
	$p \wedge q$	

# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
1.1.1.1	$p$	-1
1.1.1.2	$q$	1
1.1.2	$p \wedge q \rightarrow r$	1
1.1.2.1	$p \wedge q$	-1
	$p$	

# Position and polarity, again

position	subformula	polarity
€	$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r))$	1
1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (p \rightarrow r)$	-1
1.1	$(p \rightarrow q) \wedge (p \wedge q \rightarrow r)$	1
1.1.1	$p \rightarrow q$	1
1.1.1.1	$p$	-1
1.1.1.2	$q$	1
1.1.2	$p \wedge q \rightarrow r$	1
1.1.2.1	$p \wedge q$	-1
1.1.2.1.1	$p$	-1
1.1.2.1.2	$q$	-1
1.1.2.2	$r$	1
1.2	$p \rightarrow r$	-1
1.2.1	$p$	1
1.2.2	$r$	-1

Monotonic replacement  
pure atom rule

# Connection between $\rightarrow$ and $\leq$

Notation:

- ▶  $A[B]_{\pi}$  denotes a formula  $A$  with the subformula  $B$  at the position  $\pi$ ;
- ▶  $A[B']_{\pi}$  denotes  $A$  with the subformula at the position  $\pi$  replaced by  $B'$ .

Remind: For any interpretation  $I$ :

$$I(A) = I(B) \text{ if and only if } I \models A \leftrightarrow B$$

Lemma. For any interpretation  $I$ :

$$I(A) \leq I(B) \text{ if and only if } I \models A \rightarrow B.$$

# Connection between $\rightarrow$ and $\leq$

Notation:

- ▶  $A[B]_{\pi}$  denotes a formula  $A$  with the subformula  $B$  at the position  $\pi$ ;
- ▶  $A[B']_{\pi}$  denotes  $A$  with the subformula at the position  $\pi$  replaced by  $B'$ .

**Remind:** For any interpretation  $I$ :

$$I(A) = I(B) \text{ if and only if } I \models A \leftrightarrow B$$

**Lemma.** For any interpretation  $I$ :

$$I(A) \leq I(B) \text{ if and only if } I \models A \rightarrow B.$$



# Monotonic replacement lemma

Let  $I$  be an interpretation.

Remind:

Lemma (Equivalent Replacement)

If  $I \models B \leftrightarrow B'$ , then

$$I \models A[B] \leftrightarrow A[B'].$$

$I(B) = I(B')$  then

$$I(A[B]_{\pi}) = I(A[B']_{\pi})$$

# Monotonic replacement lemma

Let  $I$  be an interpretation.

Remind:

## Lemma (Equivalent Replacement)

If  $I \models B \leftrightarrow B'$ , then

$$I \models A[B] \leftrightarrow A[B'].$$

$$I(B) = I(B') \text{ then}$$

$$I(A[B]_{\pi}) = I(A[B']_{\pi})$$

## Lemma (Monotonic Replacement)

► Let  $pol(A, \pi) = 1$  and  $I \models B \rightarrow B'$ , then

$$I(B) \leq I(B') \text{ then}$$

$$I \models A[B]_{\pi} \rightarrow A[B']_{\pi}$$

$$I(A[B]_{\pi}) \leq I(A[B']_{\pi})$$

# Monotonic replacement lemma

Let  $I$  be an interpretation.

Remind:

Lemma (Equivalent Replacement)

If  $I \models B \leftrightarrow B'$ , then

$$I \models A[B] \leftrightarrow A[B'].$$

$I(B) = I(B')$  then

$$I(A[B]_{\pi}) = I(A[B']_{\pi})$$

Lemma (Monotonic Replacement)

► Let  $pol(A, \pi) = 1$  and  $I \models B \rightarrow B'$ , then

$$I \models A[B]_{\pi} \rightarrow A[B']_{\pi}$$

$I(B) \leq I(B')$  then

$$I(A[B]_{\pi}) \leq I(A[B']_{\pi})$$

► Let  $pol(A, \pi) = -1$  and  $I \models B' \rightarrow B$ , then

$$I \models A[B]_{\pi} \rightarrow A[B']_{\pi}$$

$I(B') \leq I(B)$  then

$$I(A[B]_{\pi}) \leq I(A[B']_{\pi})$$

# Monotonic replacement theorem

Note:  $A \leftrightarrow B$  is **valid** if and only if  $A \equiv B$ .

Remind:

## Theorem (Equivalent Replacement)

If  $B \leftrightarrow B'$  is valid then

$A[B] \leftrightarrow A[B']$  is valid.

## Theorem (Monotonic Replacement)

- ▶ Let  $pol(A, \pi) = 1$  and  $B \rightarrow B'$  is valid, then

$A[B]_{\pi} \rightarrow A[B']_{\pi}$  is valid.

- ▶ Let  $pol(A, \pi) = -1$  and  $B' \rightarrow B$  is valid, then

$A[B]_{\pi} \rightarrow A[B']_{\pi}$  is valid.

*Positive/negative polarity is sufficient condition for monotonicity/anti-monotonicity (but not necessary).*

# Pure Atom

Atom  $p$  is **pure in a formula  $A$** , if either all occurrences of  $p$  in  $A$  are positive or all occurrences of  $p$  in  $A$  are negative.

$$p \wedge r \rightarrow (\neg q \rightarrow (r \wedge \neg p))$$




- Both occurrences of  $p$  are negative, so  $p$  is pure.
- The only occurrence of  $q$  is positive, so  $q$  is pure.
- $r$  is not pure, since it has both negative and positive occurrences.

# Pure Atom

Atom  $p$  is **pure in a formula**  $A$ , if either all occurrences of  $p$  in  $A$  are positive or all occurrences of  $p$  in  $A$  are negative.

$$p \wedge r \rightarrow (\neg q \rightarrow (r \wedge \neg p))$$




- Both occurrences of  $p$  are negative, so  $p$  is pure.
- The only occurrence of  $q$  is positive, so  $q$  is pure.
- $r$  is not pure, since it has both negative and positive occurrences.

# Pure Atom

Atom  $p$  is **pure in a formula**  $A$ , if either all occurrences of  $p$  in  $A$  are positive or all occurrences of  $p$  in  $A$  are negative.

$$p \wedge r \rightarrow (\neg q \rightarrow (r \wedge \neg p))$$




- ▶ Both occurrences of  $p$  are negative, so  $p$  is pure.
- ▶ The only occurrence of  $q$  is positive, so  $q$  is pure.
- ▶  $r$  is not pure, since it has both negative and positive occurrences.

# Pure Atom

Atom  $p$  is **pure in a formula**  $A$ , if either all occurrences of  $p$  in  $A$  are positive or all occurrences of  $p$  in  $A$  are negative.

$$p \wedge r \rightarrow (\neg q \rightarrow (r \wedge \neg p))$$




- ▶ Both occurrences of  $p$  are negative, so  $p$  is pure.
- ▶ The only occurrence of  $q$  is positive, so  $q$  is pure.
- ▶  $r$  is not pure, since it has both negative and positive occurrences.



# Pure Atom

Atom  $p$  is **pure in a formula**  $A$ , if either all occurrences of  $p$  in  $A$  are positive or all occurrences of  $p$  in  $A$  are negative.

$$p \wedge r \rightarrow (\neg q \rightarrow (r \wedge \neg p))$$



- ▶ Both occurrences of  $p$  are negative, so  $p$  is pure.
- ▶ The only occurrence of  $q$  is positive, so  $q$  is pure.
- ▶  $r$  is not pure, since it has both negative and positive occurrences.

# Properties of Pure Atoms

## Theorem (Pure Atom)

Let an atom  $p$  has only *positive* (respectively, only *negative*) occurrences in  $A$ . Then  $A$  is *satisfiable* if and only if so is  $A_p^\top$  (respectively,  $A_p^\perp$ ).

We can prove *Pure Atom Theorem* by applying *Monotonic Replacement Theorem*:

- $p \rightarrow \top$  is valid. If  $p$  has only positive occurrences in  $A$  then  $A \rightarrow A_p^\perp$  is valid. In other words,  $I(A) \leq I(A_p^\perp)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\perp)$ .
- $\perp \rightarrow p$  is valid. If  $p$  has only negative occurrences in  $A$  then  $A \rightarrow A_p^\top$  is valid. In other words,  $I(A) \leq I(A_p^\top)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\top)$ .

# Properties of Pure Atoms

## Theorem (Pure Atom)

Let an atom  $p$  has only *positive* (respectively, only *negative*) occurrences in  $A$ . Then  $A$  is *satisfiable* if and only if so is  $A_p^\top$  (respectively,  $A_p^\perp$ ).

We can prove **Pure Atom Theorem** by applying **Monotonic Replacement Theorem**:

- $p \rightarrow \top$  is valid. If  $p$  has only positive occurrences in  $A$  then  $A \rightarrow A_p^\top$  is valid. In other words,  $I(A) \leq I(A_p^\top)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $(A_p^\top)$ .
- $\perp \rightarrow p$  is valid. If  $p$  has only negative occurrences in  $A$  then  $A \rightarrow A_p^\perp$  is valid. In other words,  $I(A) \leq I(A_p^\perp)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $(A_p^\perp)$ .

# Properties of Pure Atoms

## Theorem (Pure Atom)

Let an atom  $p$  has only *positive* (respectively, only *negative*) occurrences in  $A$ . Then  $A$  is *satisfiable* if and only if so is  $A_p^\top$  (respectively,  $A_p^\perp$ ).

We can prove **Pure Atom Theorem** by applying **Monotonic Replacement Theorem**:

- ▶  $p \rightarrow \top$  is valid. If  $p$  has only *positive* occurrences in  $A$  then  $A \rightarrow A_p^\top$  is valid. In other words,  $I(A) \leq I(A_p^\top)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\top)$ .
- ▶  $\perp \rightarrow p$  is valid. If  $p$  has only *negative* occurrences in  $A$  then  $A \rightarrow A_p^\perp$  is valid. In other words,  $I(A) \leq I(A_p^\perp)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\perp)$ .

# Properties of Pure Atoms

## Theorem (Pure Atom)


Let an atom  $p$  has only *positive* (respectively, only *negative*) occurrences in  $A$ . Then  $A$  is *satisfiable* if and only if so is  $A_p^\top$  (respectively,  $A_p^\perp$ ).

We can prove *Pure Atom Theorem* by applying *Monotonic Replacement Theorem*:

- ▶  $p \rightarrow \top$  is valid. If  $p$  has only *positive* occurrences in  $A$  then  $A \rightarrow A_p^\top$  is valid. In other words,  $I(A) \leq I(A_p^\top)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\top)$ .
- ▶  $\perp \rightarrow p$  is valid. If  $p$  has only *negative* occurrences in  $A$  then  $A \rightarrow A_p^\perp$  is valid. In other words,  $I(A) \leq I(A_p^\perp)$  for any interpretation  $I$ . In particular, if  $I$  satisfies  $A$  then  $I$  also satisfies  $I(A_p^\perp)$ .

# Pure atom rule, example


Consider  $\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r))$ .



All occurrences of  $p$  are negative, so, for the purpose of checking satisfiability we can replace  $p$  by  $\perp$ .

# Pure atom rule, example


Consider  $\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r))$ .



All occurrences of  $p$  are negative, so, for the purpose of checking satisfiability we can replace  $p$  by  $\perp$ .

## Pure atom rule, example

Consider  $\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r))$ .



All occurrences of  $p$  are negative, so, for the purpose of checking satisfiability we can replace  $p$  by  $\perp$ .



## Example, continued

$$\begin{aligned}& \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) \\& \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\& \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\& \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\& \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\& \neg(\top \rightarrow (\neg \perp \rightarrow r)) \\& \neg(\neg \perp \rightarrow r) \\& \neg(\top \rightarrow r) \\& \neg r \\& \neg \perp \\& \top\end{aligned}$$

All occurrences of  $p$  are negative, so, for the purpose of checking satisfiability we can replace  $p$  by  $\perp$ .

## Example, continued

$$\begin{aligned} & \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) \quad \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) \\ & \neg(\neg \perp \rightarrow r) \\ & \neg(\top \rightarrow r) \\ & \neg r \\ & \neg \perp \\ & \top \end{aligned}$$

All occurrences of  $p$  are negative, so, for the purpose of checking satisfiability we can **replace  $p$  by  $\perp$** .

## Example, continued

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) \Rightarrow$$

$$\neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\top \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\neg \perp \rightarrow r)$$

$$\neg(\top \rightarrow r)$$

$$\neg r$$

$$\neg \perp$$

$$\top$$

## Example, continued

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) \Rightarrow$$

$$\neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\top \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\neg \perp \rightarrow r)$$

$$\neg(\top \rightarrow r)$$

$$\neg r$$

$$\neg \perp$$

$$\top$$

## Example, continued

$$\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) \Rightarrow$$

$$\neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) \Rightarrow$$

$$\neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\top \rightarrow (\neg \perp \rightarrow r))$$

$$\neg(\neg \perp \rightarrow r)$$

$$\neg(\top \rightarrow r)$$

$$\neg r$$

$$\neg \perp$$

$$\top$$

## Example, continued

$$\begin{aligned} & \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) \\ & \quad \neg(\neg \perp \rightarrow r) \\ & \quad \neg(\top \rightarrow r) \\ & \quad \neg r \\ & \quad \neg \perp \\ & \quad \top \end{aligned}$$

## Example, continued

$$\begin{aligned} & \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\neg \perp \rightarrow r) \\ & \neg(\top \rightarrow r) \\ & \neg r \\ & \neg \perp \\ & \top \end{aligned}$$

## Example, continued

$$\begin{aligned} & \neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\neg \perp \rightarrow r) && \Rightarrow \\ & \neg(\top \rightarrow r) \\ & \neg r \\ & \neg \perp \\ & \top \end{aligned}$$



## Example, continued

$$\begin{aligned} & \neg((\textcolor{red}{p} \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg \textcolor{red}{p} \rightarrow r)) & \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) & \Rightarrow \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) & \Rightarrow \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) & \Rightarrow \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) & \Rightarrow \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) & \Rightarrow \\ & \neg(\neg \perp \rightarrow r) & \Rightarrow \\ & \neg(\top \rightarrow r) & \Rightarrow \\ & \neg \textcolor{red}{r} \\ & \neg \perp \\ & \top \end{aligned}$$

All occurrences of  $\textcolor{blue}{r}$  are negative, so, for the purpose of checking satisfiability we can replace  $\textcolor{red}{r}$  by  $\perp$ .

## Example, continued

$$\begin{aligned} & \neg((\textcolor{red}{p} \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg \textcolor{red}{p} \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\top \rightarrow (\neg \perp \rightarrow r)) && \Rightarrow \\ & \neg(\neg \perp \rightarrow r) && \Rightarrow \\ & \neg(\top \rightarrow r) && \Rightarrow \\ & \neg \textcolor{red}{r} && \Rightarrow \\ & \neg \perp && \\ & \top && \end{aligned}$$

All occurrences of  $\textcolor{blue}{r}$  are negative, so, for the purpose of checking satisfiability we can **replace  $r$  by  $\perp$** .

## Example, continued

$$\begin{aligned}\neg((p \rightarrow q) \wedge (p \wedge q \rightarrow r) \rightarrow (\neg p \rightarrow r)) &\Rightarrow \\ \neg((\perp \rightarrow q) \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) &\Rightarrow \\ \neg(\top \wedge (\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) &\Rightarrow \\ \neg((\perp \wedge q \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) &\Rightarrow \\ \neg((\perp \rightarrow r) \rightarrow (\neg \perp \rightarrow r)) &\Rightarrow \\ \neg(\top \rightarrow (\neg \perp \rightarrow r)) &\Rightarrow \\ \neg(\neg \perp \rightarrow r) &\Rightarrow \\ \neg(\top \rightarrow r) &\Rightarrow \\ \neg r &\Rightarrow \\ \neg \perp &\Rightarrow \\ \top &\end{aligned}$$

We have shown satisfiability of this formula deterministically, using only the **pure atom rule**.

# Summary

We have studied:

- ▶ how to **formalise** problems in propositional logic,
- ▶ **splitting algorithm** for checking satisfiability,
- ▶ **position/polarity** of a subformula occurrence,
- ▶ **monotonic replacement**,
- ▶ **pure atom rule**.