NEUROSCIENCES JOURNAL 2026;31:108–126

Scan the QR to view
the full-text article on
the journal website

REVIEW ARTICLE

The Role of the Ventral Tegmental Area in Reward and Addiction: A Comprehensive Review

Emad Ali Albadawi, MSc, PhD

Department of Basic Medical Sciences, College of Medicine, Taibah University, Madinah, Kingdom of Saudi Arabia

ABSTRACT

The ventral tegmental area (VTA) is a central hub in reward and motivation, integrating heterogeneous neuronal populations and projection-defined circuits to regulate learning, decision-making, and addiction. This review synthesizes anatomical organization and cellular diversity (dopaminergic, GABAergic, and glutamatergic neurons, including multi-transmitter co-release), and maps major efferent pathways constituting the mesolimbic and mesocortical systems. We highlight computational roles of VTA activity, including reward prediction errors, incentive salience, and context memory integration, and summarize drug class-specific mechanisms that perturb VTA output. We then outline time-dependent neuroadaptations across mesocorticolimbic networks and discuss therapeutic implications spanning pharmacological, behavioral, and emerging circuit-targeted interventions. Together, these perspectives frame the VTA as a projection- and state-dependent controller whose dysfunction underlies core features of substance use disorders and other reward-related psychopathologies, pointing toward precision, circuit-informed treatments.

Keywords: Ventral tegmental area (VTA), Mesolimbic pathway, Mesocortical pathway, Neuromodulation

Introduction

The ventral tegmental area (VTA) is one of the most intensively studied brain regions in neuroscience, serving as a critical component of the brain's reward system and playing a central role in the pathophysiology of addiction [1]. Located in the midbrain, the VTA has been recognized for over five decades as a key structure mediating the rewarding effects of natural stimuli and drugs of abuse. The significance of this region was first highlighted by the pioneering work of James Olds and Peter Milner in the 1950s, who demonstrated that electrical stimulation of the VTA and related areas could serve as a powerful reinforcer of instrumental behavior, driving animals to repeatedly perform actions (such as lever pressing) to obtain the stimulation in laboratory animals [2]. The VTA's importance extends beyond addiction research, as it plays fundamental roles in motivation, learning, decision-making, and various cognitive processes. Its dysfunction has been implicated in numerous psychiatric and neurological conditions, including depression, schizophrenia, attention deficit hyperactivity disorder, and Parkinson's disease. The VTA's complex anatomy and diverse cellular composition enable it to integrate multiple types of information and modulate behavior in sophisticated ways [3–5]. This review comprehensively examines the VTA's anatomical organization, neural connections, and functional roles in reward processing and addiction. We explore the cellular and molecular mechanisms that underlie VTA function, discuss how these processes become dysregulated in addiction, and consider the therapeutic implications of current research findings.

Terminology note: Throughout this review, we use the term "addiction" to refer to the human clinical syndrome, typically operationalized as substance use disorders in diagnostic classifications. In contrast, when we discuss preclinical studies in rodents

Received 27 August 2025; accepted 13 December 2025.
Available online 30 April 2026

E-mail address: ebadawi@taibahu.edu.sa, dr-albadawi@hotmail.com (E. A. Albadawi).

https://doi.org/10.17712/1658-3183.2787
1658-3183/© 2026 Neurosciences Journal. All rights reserved. This is an open access article under the CC BY-NC 4.0 Licence (https://creativecommons.org/licenses/by-nc/4.0/).




Fig. 1. Schematic representation of the ventral tegmental area (VTA) and its major dopaminergic projection pathways. The VTA (green dot) gives rise to the mesolimbic pathway (green) to the nucleus accumbens (NAc) and the mesocortical pathway (magenta) to cortical targets. For comparison, the figure also depicts the nigrostriatal pathway (blue), which originates primarily from the adjacent substantia nigra pars compacta (SNc) and innervates the dorsal striatum. The relative positions of the cortex, striatum, nucleus accumbens (NAc), and midbrain structures are shown for anatomical context.

and other animals, we refer to "drug dependence" or "dependence-like"/"addiction-like" behaviors, because these paradigms model specific components of the disorder (such as escalation of intake, withdrawal, and relapse-like responding) rather than the whole phenomenology of human addiction.

Anatomical organization of the ventral tegmental area

Location and boundaries

The VTA is situated in the ventral midbrain, medial to the substantia nigra and dorsal to the interpeduncular nucleus. In humans, it extends approximately from the level of the posterior commissure to the mammillary bodies, spanning roughly 5–6 millimeters in the rostrocaudal dimension as presented in Fig. 1 [6–9]. The VTA's boundaries have been refined over decades of research, with modern anatomical studies defining it based on cytoarchitecture, immunohistochemistry, and connectivity patterns. The VTA can be subdivided into several distinct subnuclei, each with unique anatomical and functional characteristics [10]. The para-nigral nucleus (PN) forms the medial portion and contains the highest concentration of dopaminergic neurons. The parabrachial pigmented nucleus (PBP) occupies the central region and shows intermediate dopamine neuron density [11, 12]. The rostral VTA (VTAR) and caudal VTA also represent distinct subdivisions with different projection patterns and functional roles (Fig. 1) [13].

Cellular composition

The VTA contains a heterogeneous population of neurons broadly classified into three major types based on their neurotransmitter phenotype: dopaminergic neurons, GABAergic neurons, and glutamatergic neurons. This cellular diversity underlies the VTA's

complex functional properties and its ability to modulate behavior in a nuanced manner [8, 11]. Critically, these subtypes form a local microcircuit in which dopaminergic neurons drive reward-related signaling, GABAergic neurons provide inhibitory control essential for aversive processing and for regulating dopamine output, and glutamatergic neurons contribute excitatory drive that influences behavioral valence. The interplay of these distinct populations allows the VTA to compute motivational value and guide adaptive behavior [13].

Neuronal subtypes in the ventral tegmental area

Dopaminergic neurons (DA)

DA represent the predominant neuronal population in the VTA, comprising approximately 65–70% of its cells [14]. These neurons are critically involved in reward processing, motivational salience, learning, and goal-directed behavior [8]. They are identified by their tyrosine hydroxylase (TH) expression, the rate-limiting enzyme in dopamine synthesis. They exhibit distinct electrophysiological characteristics, including slow, pacemaker-like firing and broad action potentials with prominent after-hyperpolarizations [15]. Anatomically, VTA DA can be categorized based on their projection targets and molecular signatures. Those projecting to the Nucleus accumbens (NAc), particularly the core and shell regions, form the mesolimbic dopamine pathway, essential for reinforcing behavior and encoding reward prediction errors. Another major subpopulation targets the prefrontal cortex (PFC), creating the mesocortical pathway implicated in executive function, working memory, and value-based decision-making. Additional projections extend to the amygdala, hippocampus, and lateral septum, supporting emotional regulation, stress responsivity, and contextual information integration [16, 17]. Importantly, DA are not homogeneous. Recent studies employing single-cell transcriptomics and electrophysiological profiling have revealed distinct subtypes with specialized roles. Moreover, some VTA DA corelease glutamate or GABA, expanding their functional repertoire beyond classical DA signaling. These co-transmission modalities allow bidirectional modulation of target regions, particularly under stress, learning, or substance exposure conditions [18, 19].

GABAergic neurons

GABAergic neurons constitute approximately 20–30% of the VTA and serve local inhibitory functions and long-range modulatory roles [20]. As the primary inhibitory cells of the VTA, they play a crucial role in maintaining the excitatory-inhibitory balance and in regulating dopamine output [20]. Canonical VTA GABAergic neurons are defined by their robust expression of the glutamate decarboxylase isoforms GAD1 (GAD67) and GAD2 (GAD65), whereas subsets of VTA dopamine neurons can also express GAD65 and thereby engage GABA co-transmission [21]. Electrophysiologically, GABAergic neurons are distinct from DA cells, with typically faster firing rates and shorter-duration action potentials [22, 23]. Within the VTA, GABAergic neurons act as local interneurons, regulating the excitability of nearby DA through inhibitory synapses [24]. This inhibition is crucial in controlling dopamine neuron burst firing and dopamine release in target regions. Additionally, GABAergic projection neurons from the VTA extend to structures such as the nucleus accumbens, prefrontal cortex, and central amygdala (CeA), influencing motivation, emotional reactivity, and behavioral flexibility [24]. The functional relevance of VTA GABA neurons has been increasingly recognized in both reward and aversion paradigms [20, 25, 26]. For example, optogenetic activation of these neurons can suppress reward-seeking behavior [25] and modulate responses to aversive stimuli [26]. Importantly, these neurons are also involved in the feedforward inhibition circuit from the lateral habenula (LHb) via the rostromedial tegmental nucleus (RMTg), which plays a pivotal role in aversive learning and stress-induced modulation of reward processing [27, 28].

Glutamatergic neurons

Although comprising only 5–10% of the total VTA population, glutamatergic neurons play indispensable roles in local excitation and long-range signaling [29]. These neurons provide a critical excitatory counterpart to dopamine and GABA signaling, directly exciting target regions and influencing behavioral valence [1]. These neurons express vesicular glutamate transporter 2 (VGLUT2) and can activate local VTA neurons and distant targets through excitatory transmission. Glutamatergic VTA neurons project to the prefrontal cortex, lateral habenula, and nucleus accumbens, contributing to the encoding of rewarding and aversive stimuli. Recent studies have demonstrated that selective activation of VTA glutamate neurons can reinforce behavior or induce aversive responses depending on the projection target and firing pattern. This dual functionality emphasizes their importance in valence processing and behavioral flexibility [1, 30]. Moreover, glutamatergic VTA neurons often participate in co-release mechanisms,

Table 1. Characteristics of VTA Neuronal Populations.

Neuron Type	Neurotransmitter(s)	Molecular Markers	Projection Targets	Functional Roles	Ref.
DA	Dopamine	TH, GCH1, DAT	Nucleus accumbens (NAc), prefrontal cortex (PFC), amygdala	Reward processing, motivation, and learning	[32, 33]
GABAergic	GABA	GAD1, GAD2	Local interneurons, PFC, NAc	Inhibition of DA neurons, modulation of reward circuits	[34]
Glutamatergic	Glutamate	VGLUT2	PFC, lateral habenula (LHb)	Excitation of target neurons, involvement in aversion and reward	[1]
DA-Glutamate Co-releasing	Dopamine & Glutamate	TH, VGLUT2	NAc, PFC	Modulation of reward and aversive behaviors	[35]
DA-GABA Co-releasing	Dopamine & GABA	TH, GAD2	NAc, PFC	Fine-tuning of reward circuits, behavioral flexibility	[35, 36]
GABA-Glutamate Co-releasing	GABA & Glutamate	GAD2, VGLUT2	Local circuits	Balance of excitation and inhibition within the VTA	[37]
Triple Co-releasing	Dopamine, GABA & Glutamate	TH, GAD2, VGLUT2	Various targets	Complex modulation of neural circuits	[35]

especially in DA glutamate co-releasing neurons expressing both TH and VGLUT2 [31]. These hybrid neurons allow for rapid excitatory transmission alongside modulatory DA signaling, enhancing the dynamic range of VTA output during motivationally relevant events. Table 1 provides a detailed classification of VTA neuronal subtypes, including DA, GABAergic, and glutamatergic populations, as well as co-releasing phenotypes, based on their neurotransmitter profiles, molecular markers, projection targets, and roles in modulating reward-related behaviors.

Afferent connections of the ventral tegmental area

VTA is a central integrative node within the brain's reward and motivation circuits. Its ability to orchestrate various behaviors, including reward processing, executive function, affect regulation, and learning, depends on a complex matrix of reciprocal and feedforward connections [38]. This intricate input-output architecture enables the VTA to dynamically regulate activity across limbic, cortical, and subcortical regions. Major afferent and efferent connections include projections to the nucleus accumbens (NAc), prefrontal cortex (PFC), and the central amygdala (CeA), while receiving modulatory input from structures such as (LHb) and the rostromedial tegmental nucleus (RMTg) [39]. The following subsections outline these key circuits' anatomical and functional relevance.

VTA–NAc circuitry

The mesolimbic dopamine pathway, extending from the VTA to the NAc, forms the neurobiological core of reinforcement learning and motivational behavior. While the VTA DA project to the NAc, this circuit is also modulated by GABAergic feedback arising primarily from D1 receptor-expressing medium spiny neurons (MSNs) within the NAc [8]. These reciprocal connections differ across NAc subregions: lateral shell (NAcLat) MSNs preferentially target VTA GABAergic neurons, resulting in disinhibition of dopamine neurons, whereas medial shell (NAcMed) MSNs exert more selective inhibition. Specifically, they suppress dopamine neurons projecting to NAcMed via $GABA_A$ receptors and those projecting to NAcLat via $GABA_B$ receptors. This dual inhibitory mechanism finely calibrates motivational

drive and action selection, as demonstrated by optogenetic manipulation studies [40, 41]. In addition to its role in reinforcement, the VTA-NAc pathway is integral to feeding behavior. Exposure to palatable food elevates dopamine release in the NAc, while optogenetic activation of VTA dopamine neurons has been shown to suppress feeding, even under caloric deprivation. This anorexigenic effect is mediated through D1 receptor activation in the NAc and can be reversed by blocking NAc–VTA projections. Furthermore, hypothalamic neuropeptides such as orexin and neurotensin modulate this circuitry by acting directly on VTA dopamine neurons, integrating energy homeostasis with reward-driven behavior [42, 43].

VTA-PFC circuitry

The mesocortical pathway, linking the VTA to the PFC, is essential for cognitive flexibility, emotional regulation, and behavioral control domains often disrupted in psychiatric disorders such as depression and addiction. VTA dopamine neurons modulate prefrontal function through dopamine release and co-release of glutamate and GABA, adding a layer of temporal and synaptic specificity [44]. Short bursts of VTA activity enhance excitability in both pyramidal neurons and parvalbumin-positive (PV+) interneurons in the PFC through glutamatergic transmission. However, sustained stimulation recruits additional neuromodulators: dopamine promotes transient excitation of pyramidal neurons, while GABAergic co-release prolongs interneuron activation. This bidirectional modulation contributes to the dynamic excitatory/inhibitory (E/I) balance required for complex cognitive operations. Notably, the PFC exerts top-down control over VTA function [45]. Anatomical and tracing studies indicate that medial PFC pyramidal neurons send both ipsilateral and decussating projections to the VTA, including synapses onto dopamine neurons, although the density of these direct inputs is modest relative to other major afferents [46, 47]. In parallel, indirect glutamatergic input via intermediary circuits can significantly elevate dopamine release in the NAc. Functional studies reveal heterogeneous responses in VTA neurons to PFC stimulation, including enhanced burst firing and inhibition, suggesting the involvement of relay inhibitory networks. These mechanisms underscore the nuanced role of PFC–VTA interactions in behavioral adaptation and stress resilience [8, 48, 49].

VTA–CeA circuitry

Projections from the VTA to the (CeA) form a critical interface between emotional valence and motivated behavior. VTA output to the CeA includes dopaminergic, glutamatergic, and GABAergic projections, highlighting the multimodal nature of this connection [50]. This circuit has been implicated in various alcohol-related behaviors. For example, in alcohol-preferring rodent models, increased activity of CeA-projecting VTA dopamine neurons during withdrawal reinforces alcohol-seeking. Orexinergic signaling in the VTA further amplifies flavor preference through CeA D1 receptor activation [51]. Beyond reward, this pathway also contributes to pain modulation and defensive behavior. Dopamine release in the CeA is associated with analgesia, which is reversed by D2 receptor activation. Moreover, VTA GABAergic neurons mediate aversive responses, while glutamatergic projections have been linked to arousal and wakefulness. These findings position the VTA-CeA axis as vital for integrating sensory, affective, and motivational cues [52, 53].

VTA–LHb and RMTg circuitry

The (LHb) is a major aversive input to the VTA, largely mediated through the rostromedial tegmental nucleus (RMTg), a GABAergic relay structure. Activation of LHb neurons stimulates RMTg outputs, which in turn inhibit VTA dopamine neurons. This trisynaptic inhibitory circuit plays a central role in encoding reward prediction errors, regulating aversive learning, and modulating stress responsiveness [54]. Chronic stress enhances glutamatergic drive from the LHb to the RMTg, potentiating inhibition of VTA dopamine neurons and contributing to depressive-like phenotypes [55]. In contrast, acute stress may transiently suppress LHb D-neuron activity via trace amine signaling, disinhibiting dopamine neurons and facilitating adaptive arousal. Importantly, LHb inputs exhibit projection-specific modulation, preferentially targeting VTA neurons projecting to the medial PFC [56]. This topography underlies aversive decision-making and is particularly relevant in models of cocaine relapse during extinction phases. Given its dual role in reward suppression and affective regulation, the LHb–VTA axis represents a promising therapeutic target for mood and substance use disorders [57–59].

Efferent connections of the ventral tegmental area

The efferent projections of the ventral tegmental area (VTA) form the anatomical and functional substrate through which it exerts widespread influence on behavior, cognition, and affective states. These output pathways primarily originate from DA, GABAergic, and glutamatergic neurons, often

Table 2. Major efferent projections of the VTA.

Target region (examples)	Principal VTA neuron type(s)	Transmitter(s)	Representative functional roles	Ref.
Nucleus accumbens (core/shell)	DA; DA–Glu co-releasing; GABAergic	Dopamine, glutamate, GABA	Reward processing, motivation, reinforcement learning; VTA→NAc glutamate can suppress psychostimulant seeking	[8, 69]
Prefrontal cortex (mPFC, OFC, ACC)	DA (mesocortical)	Dopamine	Cognitive control, decision-making, emotion regulation; VTA→mPFC DA required for rapid threat-avoidance learning	[70]
Amygdala (BLA, CeA)	DA; (some) glutamatergic	Dopamine, glutamate	Emotion/fear and reward valence; VTA→BLA DA controls anxiety-related behavior	[71, 72]
Hippocampus (ventral/dorsal)	DA; (reports of VTA Glu inputs)	Dopamine (± glutamate)	Contextual memory, novelty signals; VTA→Hip DA triggers LTP and supports contextual learning	[65]
Lateral septum	DA; Glutamatergic	Dopamine, glutamate	Social behavior, aggression, reward modulation	[73]
Hypothalamus (LHA, ARC, PVN)	DA	Dopamine	Arousal, energy balance, feeding regulation; VTA→LHA DA release during cue-driven feeding	[74]

displaying complex co-release profiles that fine-tune neural signaling (Table 2). Rather than reiterating functions detailed in earlier circuit sections, this part emphasizes how projection-defined VTA outputs differentially engage downstream targets to coordinate reward learning, executive control, and emotional regulation across multiple brain regions [60, 61].

Mesolimbic pathway

The mesolimbic dopamine pathway is the most extensively studied VTA output system and constitutes the brain's reward circuitry core. It projects predominantly to the nucleus accumbens (NAc), including core and shell subregions and adjacent ventral striatal structures. VTA neurons projecting to the NAc are primarily DA, although subpopulations exhibit co-release of glutamate or GABA, adding complexity to the modulation of postsynaptic targets. In addition to these dopaminergic and co-releasing projections, VTA GABA projection neurons also innervate nucleus accumbens cholinergic interneurons, providing an additional route by which VTA output can shape accumbal microcircuit excitability and plasticity [24]. Together, these DA-only, co-releasing, and purely GABAergic efferents allow the VTA to regulate NAc microcircuits through multiple neurochemical channels, complementing the functional roles of this pathway in reward prediction, motivational salience, and reinforcement learning described earlier. Dysregulation of the mesolimbic projection, particularly under chronic drug exposure, underlies key features of addiction such as craving, compulsive seeking, and sensitization [62].

Mesocortical pathway

The mesocortical pathway comprises DA projections from the VTA to various prefrontal cortical regions, including the medial prefrontal cortex (mPFC), orbitofrontal cortex (OFC), and anterior cingulate cortex (ACC). This pathway is critically involved in executive function, working memory, decision-making, and emotional regulation [63]. In contrast to the more phasic, event-locked dynamics of mesolimbic dopamine, mesocortical dopamine transmission is generally more tonic and projection-specific, providing sustained modulatory input that shapes prefrontal network activity over longer timescales [63]. Importantly, these projections exhibit projection-specific modulation of local circuits: dopamine enhances the excitability of pyramidal neurons and modulates the recruitment of interneurons, thereby shaping the balance of cortical excitation and inhibition. Alterations in mesocortical signaling have been implicated in various psychiatric disorders, including schizophrenia, depression, and attention-deficit/hyperactivity disorder (ADHD) [64].

Extended limbic and hypothalamic projections

Beyond the striatum and cortex, the VTA sends projections to several limbic and hypothalamic regions, supporting its role in integrating motivational and homeostatic signals. Key targets include the: Amygdala: VTA outputs to the central and basolateral amygdala contribute to emotional learning, fear conditioning, and affective modulation. Both DA and non-DA projections are involved, with complex effects depending on neuronal subtype and receptor expression. Hippocampus: Although less dense than cortical or striatal projections, VTA innervation of the ventral hippocampus enhances contextual memory formation, particularly under emotionally salient or rewarding conditions [65]. Lateral septum: DA projections to the septal area are involved

in social behavior, reward modulation, and stress reactivity [66]. Hypothalamic nuclei: VTA projections to the lateral hypothalamus, arcuate nucleus, and paraventricular nucleus mediate interactions between energy balance, arousal, and reward valuation. These connections integrate metabolic and hedonic drives, particularly in feeding and drug-seeking contexts [41].

Neurochemical diversity and Co-transmission

A hallmark of VTA efferent organization is the presence of co-transmission, where neurons can release combinations of dopamine, GABA, and glutamate onto distinct or overlapping targets [13]. This arrangement allows a single VTA projection to convey excitatory, inhibitory, and neuromodulatory signals in a temporally and spatially coordinated manner. For instance, VTA glutamate neurons that co-release GABA can promote positive reinforcement and exert rapid inhibitory–excitatory control over downstream targets [67], while dopamine–glutamate co-releasing neurons targeting mesolimbic regions such as the NAc can rapidly excite postsynaptic neurons and concurrently modulate plasticity via dopamine [18]. Triple-transmitting neurons identified in recent single-cell and single-nucleus RNA-sequencing studies suggest even greater computational complexity within VTA output channels [68]. This neurochemical heterogeneity underlies the functional diversity of VTA outputs, allowing the region to participate flexibly in reward, aversion, stress adaptation, and learning.

Neural circuitry and functional organization

Reward prediction and associative learning

Through DA prediction error signaling, VTA neurons encode the discrepancy between expected and actual outcomes an essential process for reinforcement learning and adaptive behavior. Seminal work by Wolfram Schultz and others has revealed that VTA dopamine neurons exhibit phasic changes in firing that reflect the value and timing of rewards. These changes in activity can be categorized into three canonical phases:

Phasic response to unexpected reward

When a reward occurs unexpectedly, VTA dopamine neurons fire a brief burst of action potentials, signaling the occurrence and magnitude of a positive outcome. This reinforces recently performed actions that led to the reward [75–77].

1. Reward prediction and cue learning

With experience, the phasic dopamine response shifts from the reward itself to the cue that reliably predicts the reward. This anticipatory activity reflects the formation of a learned association between the cue and the expected outcome.

2. Negative prediction error signaling

If an expected reward is omitted, dopamine neurons exhibit a pause or dip below baseline activity, signaling a negative prediction error. This teaches the brain that a prior expectation was incorrect, reducing the value assigned to that cue or behavior.

These bidirectional reward prediction error signals positive when outcomes are better than expected and negative when they are worse serve as teaching signals that drive plasticity in downstream structures such as the nucleus accumbens (NAc) and prefrontal cortex (PFC). Through this mechanism, the VTA contributes to adaptive decision-making, learning, and habit formation. The dynamics of DA prediction error signaling are illustrated in Fig. 2, which shows the shift in VTA firing from unpredicted reward to predictive cue and the suppression of activity following omitted rewards [75–77].

Motivation and incentive salience

VTA is a central hub in the neural circuitry of motivation, orchestrating the attribution of incentive salience to environmental cues that predict biologically relevant outcomes [78]. This process enables organisms to selectively direct attention and behavior toward goal-relevant stimuli. Crucially, the VTA mediates this transformation through phasic dopamine release to downstream regions, notably the nucleus accumbens (NAc), prefrontal cortex (PFC), and amygdala [68, 79].

Incentive salience: From neutral cue to motivational target

Incentive salience refers to the motivational "wanting" of a stimulus, distinct from its hedonic "liking." Phasic activation of VTA dopamine neurons imbues previously neutral cues with motivational significance, thereby driving cue-triggered approach behaviors [80]. This mechanism is adaptive in contexts such as food seeking, social bonding, and reward learning, but can become maladaptive under conditions of pathological salience attribution, such as in substance use disorders [81]. In addiction, for example, drug-paired cues acquire abnormally high incentive




Fig. 2. DA signaling and neural circuitry underlying reward prediction error.

salience, triggering compulsive craving and drug-seeking even in the absence of the substance. This aberrant learning is mediated by hyper-responsive VTA dopamine signaling to conditioned stimuli and drug-evoked synaptic plasticity in the NAc and PFC, including long-term potentiation and recruitment of calcium-permeable AMPA receptors at corticostriatal synapses [82, 83].

Circuit-specific DA modulation

Motivational processing by the VTA is not monolithic. Rather, functionally heterogeneous populations of dopamine neurons exhibit projection-specific roles:

- **VTA→NAc medial shell**: Encodes cue-triggered motivation and goal-directed action.
- **VTA→PFC**: Modulates executive oversight over motivational impulses, allowing for behavioral flexibility.
- **VTA→Amygdala**: Contributes to emotionally valenced salience, particularly in threat-related contexts.

This specificity is further enhanced by neurochemical co-release. DA neurons capable of co-releasing glutamate or GABA provide dual modulatory control exciting or inhibiting target circuits based on behavioral demands [15, 84]. Additionally, afferent inputs from the lateral hypothalamus, laterodorsal tegmentum (LDT), and paraventricular thalamus (PVT) shape VTA activity in response to internal physiological states (e.g., hunger, arousal) and environmental contingencies, ensuring a state-dependent encoding of motivational relevance.

Contextual processing and memory integration

VTA functions as a hub that binds spatiotemporal context with goals and rules to shape behavior

in a state-dependent manner by integrating convergent hippocampal and prefrontal inputs [85]. Hippocampal inputs convey contextual (space–time) information, while prefrontal inputs provide task rules, goals, and abstract associations; together they enable the VTA to modulate its output to downstream targets according to current contingencies and internal state. Notably, both hippocampus and prefrontal cortex receive DA innervation from the VTA, forming reciprocal loops through which prediction, context, and action selection are coordinated [86]. Within this loop, dopamine signaling in the ventral hippocampus–medial prefrontal (vHip–mPFC) circuit provides a mechanistic bridge from context to memory [87]. VTA projections to ventral hippocampus can enhance contextual memory formation under emotionally salient or rewarding conditions, positioning dopamine as a facilitator of consolidation for behaviorally significant events [85, 88]. Consistent with this, opiate paradigms demonstrate phase- and receptor-specific gating of contextual memory within the vHip–mPFC pathway: acquisition of drug–context associations engages a D1 receptor (D1R)–ERK–CREB pathway in both vHip and mPFC, whereas retrieval or reinstatement depends on D2 receptor (D2R)–Akt–GSK3 signaling in the same circuit [87]. This phase specificity is mirrored in circuit physiology. During morphine conditioned place preference (CPP), plasticity in the vHip→mPFC pathway is altered, high-frequency stimulation fails to induce LTP in morphine-paired animals, and manipulating D1R or D2R signaling bidirectionally rescues or impairs memory depending on exposure state [87]. In morphine-naïve animals, D1R-driven ERK–CREB activity supports contextual memory formation; after chronic exposure and withdrawal, D2R-linked Akt–GSK3 signaling becomes the dominant substrate for retrieval. Overactivation of D1R (during acquisition) or D2R (during reinstatement) is associated with deficits in hippocampus-dependent spatial memory, which are alleviated by blocking the corresponding receptor within the vHip–mPFC pathway. Together, these findings position the VTA as a dynamic integrator that (i) reads contextual states from the hippocampus, (ii) incorporates goal and rule signals from the prefrontal cortex, and (iii) writes back dopamine-dependent teaching signals to vHip and mPFC to shape consolidation and retrieval according to behavioral relevance [85, 87]. This closed-loop architecture helps explain how pathological salience in addiction can co-opt normal contextual processing via receptor- and phase-specific modulation within the vHip–mPFC–VTA network [89].

Role in addiction

Acute drug effects

All major drugs of abuse directly or indirectly increase dopamine levels in the VTA and its target regions, particularly the nucleus accumbens. Different classes of drugs achieve this effect through distinct mechanisms in animal models of drug dependence:

Stimulants

Psychostimulants such as cocaine, methylphenidate, and amphetamine acutely elevate mesolimbic dopamine by acting as monoamine transporters, with two separable components: a canonical dopamine-transporter (DAT) mechanism and a DAT-independent circuit mechanism. At the synapse, cocaine and methylphenidate bind DAT and inhibit reuptake, whereas amphetamine both interferes with reuptake and drives reverse transport of dopamine through DAT, boosting extracellular levels. Crucially, multiple lines of evidence show that the acute behavioral and neurochemical effects of psychostimulants cannot be explained by DAT alone. In DAT knockout mice, nucleus accumbens (NAc) dopamine still rises, and cocaine or amphetamine remains rewarding in rodent dependence-like paradigms, while a selective DAT blocker (GBR12909) is ineffective, implicating non-DAT routes. These non-DAT contributions likely arise from psychostimulant actions at norepinephrine (NET) and serotonin transporters, which can influence dopamine signaling even when DAT is absent [90, 91]. The mPFC initiates a key DAT-independent pathway. Systemic d-amphetamine elevates NAc dopamine and drives locomotion via NET-dependent norepinephrine release in mPFC and subsequent activation of α1-adrenergic receptors; blocking α1 receptors with prazosin systemically or directly in mPFC largely prevents both the dopamine rise and locomotor activation. These findings support two forms of NAc dopamine release: a “functional” form, generated distally via mPFC α1 signaling (DAT-independent) that correlates with behavior, and a “non-functional” local form within NAc driven by direct DAT interaction that lacks behavioral impact. Consistently, very high intra-NAc psychostimulant concentrations can massively raise extracellular dopamine yet fail to recapitulate the robust locomotion seen with modest systemic dosing, underscoring that the magnitude of dopamine alone is insufficient without the appropriate circuit context [91, 92]. At the level of VTA output dynamics,

psychostimulants also reshape the temporal patterning of dopamine neuron activity. They increase a slow oscillation (~0.5–1.5 Hz) in VTA dopamine neurons that is tightly coupled to prefrontal activity; this oscillatory drive can selectively gate glutamatergic inputs in NAc depending on timing, potentially explaining why mPFC-dependent, DAT-independent signaling has outsized behavioral consequences [91]. Behaviorally, mPFC norepinephrine is therefore pivotal for the acute stimulant phenotype. Depleting mPFC norepinephrine or antagonizing adrenergic receptors attenuates stimulant-evoked NAc dopamine increases, locomotor activation, and reward (e.g., conditioned place preference). Pharmacologically, α1-adrenergic antagonists (prazosin; longer-acting doxazosin) blunt sensitized locomotion and drug seeking across cocaine, amphetamine, methamphetamine, and MDMA in preclinical rodent models of drug dependence and addiction-like behavior, with early clinical signals that doxazosin can reduce cocaine's positive subjective effects, highlighting this pathway as a translational target [91, 93]. In sum, acute stimulant effects emerge from an interplay between local DAT actions and a NET/α1-driven prefrontal control loop that shapes VTA firing patterns and the temporal structure of dopamine release in NAc. This framework reconciles why systemic psychostimulants produce stronger, behaviorally relevant outcomes than isolated NAc DAT manipulation and motivates α1-adrenergic strategies to modulate acute stimulant responses [91].

Opioids disinhibition of VTA dopamine via μ-Opioid receptors

Opioids acutely increase mesolimbic dopamine by engaging μ-opioid receptors (MORs) located on GABAergic interneurons within the ventral tegmental area (VTA). MOR activation hyperpolarizes these interneurons and suppresses GABA release onto dopamine (DA) cells, thereby disinhibiting VTA DA neurons and elevating DA output to nucleus accumbens and prefrontal targets [94, 95]. More recent work has refined this view by showing that a major MOR-sensitive inhibitory input arises from GABAergic neurons in the rostromedial tegmental nucleus (RMTg), which densely innervate VTA DA neurons and encode aversive signals; opioids potently inhibit RMTg GABA cells, thereby relieving tonic inhibition of midbrain DA neurons [96, 97]. Several converging observations support this circuit mechanism. Intra-VTA pharmacology shows that a MOR agonist (e.g., DAMGO) enhances stimulant self-administration in rodent models of drug dependence, whereas VTA opioid antagonism (e.g., naltrexone) retards acquisition consistent with an opiate-driven disinhibition of VTA DA neurons. Genetic loss of MORs reduces cocaine self-administration and is accompanied by higher frequencies of spontaneous inhibitory postsynaptic currents (sIPSCs) onto identified VTA DA neurons, indicating increased GABAergic drive when MOR-mediated inhibition of GABA neurons is removed. By contrast, spontaneous glutamatergic input to DA cells is largely unchanged in MOR knockouts, highlighting the primacy of the GABA pathway in the acute opioid modulation of DA output [95, 97]. Taken together, these data support an updated "GABA-gate" model in which opioids act at MORs on both RMTg and local VTA GABA neurons to relieve inhibition of DA neurons, producing the rapid DA surge that underlies the acute reinforcing effects of opioids and potentiates the impact of other drugs (e.g., psychostimulants) [95].

Cannabis CB1 modulation of VTA microcircuits and dopamine output

Cannabinoids elevate mesolimbic dopamine predominantly by activating CB1 receptors on GABAergic, and to a lesser extent glutamatergic, terminals impinging on ventral tegmental area (VTA) dopamine (DA) neurons, thereby biasing local microcircuits toward disinhibition rather than directly acting on DA somata [98]. In vivo single-unit recordings show that Δ^9-THC dose-dependently increases VTA DA-cell firing and bursting; this excitation is prevented by the CB1-selective antagonist SR141716A but is unaffected by naloxone, demonstrating a CB1-dependent and opioid-independent mechanism. Consistent with an impulse-dependent process, cannabinoids also enhance nucleus accumbens dopamine measured by microdialysis and fast-scan voltammetry in rodent models of drug dependence [99]. Together, these findings support a circuit model in which CB1 receptor activation primarily relieves GABAergic inhibition and modulates excitatory drive within VTA microcircuits, mechanistically similar to opioid-induced disinhibition, thereby facilitating dopamine output to limbic targets. Table 3 summarizes the effects of major drug classes on the ventral tegmental area (VTA) dopamine system, detailing primary targets, circuit loci, neuronal responses, and behavioral outcomes in rodent models of drug dependence, as well as, where available, human addiction studies.

Alcohol and multiple actions in the VTA

Ethanol engages the VTA's circuit-level and cell-intrinsic mechanisms to elevate mesolimbic dopamine. Systemically, it increases dopamine

Table 3. Drug effects on VTA dopamine system.

Drug/class	Primary receptor/target(s)	Dominant VTA/circuit site	Acute effect on VTA DA neurons	Acute effect on NAc DA/behavior	Ref.
Psychostimulants (cocaine, amphetamine, methylphenidate)	DAT blockade (cocaine, MPH); DAT reverse transport (amphetamine); plus NET/α1 circuit mechanism	Local NAc DAT action; distal mPFC→VTA loop via NET/α1	↑ Firing and altered temporal patterning; systemic effects exceed isolated intra-NAc DAT action	↑ NAc DA; locomotion/reward depend on mPFC α1 signaling; α1 antagonists blunt DA rise and locomotion	[91]
Opioids (e.g., morphine)	μ-opioid receptor (MOR)	**MORs on RMTg and VTA GABA neurons → disinhibit DA neurons**	Disinhibition increases DA neuron firing	↑ Mesolimbic DA; intra-VTA MOR agonism facilitates, and VTA opioid blockade reduces, reinforcement	[95]
Alcohol (ethanol)	Direct ion-channel effects (↑ h-current; ↓ M-current; ↑ barium-sensitive K^+); modulation of GABA, glutamate, ACh, glycine inputs	Direct action on DA neuron membrane; modulatory effects via afferents	Net excitation of identified VTA DA neurons even with broad synaptic receptor blockade	↑ DA release in NAc and PFC; intra-VTA ethanol supports self-administration	[100]
Nicotine	High-affinity nAChRs on DA neurons (β2-containing; α4/α6 with α5 accessories); desensitization on GABA cells; α7 on glutamate terminals	VTA DA somata/dendrites; local GABA and glutamate afferents	↑ Firing and phasic bursting; synchronized DA output	↑ NAc DA; genetic loss of α4/β2/α5 blunts DA bursting and reinforcement; α6 contributes to reinforcement	[101]
Cannabis (Δ^9-THC; CB1 agonists)	CB1 cannabinoid receptors on GABA (and some glutamate) afferents	CB1-dependent modulation within VTA; disinhibition of DA neurons and modulation of excitatory input	↑ DA-cell firing and bursting via reduced GABAergic inhibition; effects blocked by CB1 antagonist (SR141716A)	↑ NAc DA (impulse-dependent) and reinforcement-related activity in dependence paradigms	[98, 99]

release in the nucleus accumbens and prefrontal cortex and supports intra-VTA self-administration in rodent dependence-like paradigms, establishing a direct link to reinforcement [102]. Ethanol also acts directly on VTA dopamine neurons by enhancing the hyperpolarization-activated current (h-current) and suppressing the M-current, changes that increase excitability; conversely, it can boost a barium-sensitive K^+ current, an opposing influence that helps explain dose- and state-dependent effects [102]. Repeated exposure augmented the net excitatory response to acute ethanol, consistent with VTA-level neuroadaptation. Beyond intrinsic conductances, ethanol reshapes afferent control of VTA neurons: glutamatergic [103] and cholinergic [104] drives tend to be increased, while GABAergic [105] and glycinergic inputs [106] are reduced, collectively biasing the network toward dopamine-neuron activation. Importantly, many of these synaptic effects emerge at low, behaviorally relevant ethanol concentrations, where ethanol preferentially suppresses VTA GABA neuron excitability and GABAergic transmission onto dopamine cells [105], highlighting that GABA microcircuits make a critical contribution to ethanol's initial impact on VTA output. At the receptor level, these synaptic influences are modulatory rather than obligatory ethanol's excitation of identified VTA dopamine neurons persists after dissociation from local circuitry and during broad blockade of GABA_A/B, NMDA/AMPA/mGlu, and muscarinic/nicotinic receptors [107], underscoring a bona fide direct action on dopamine-neuron membranes. Finally, ethanol perturbs dopamine autoregulation by impairing D_2-receptor desensitization in VTA neurons and alters synaptic plasticity (LTP/LTD) in both dopamine and GABA cells, changes thought to shift the reward/aversion balance during the transition from use to dependence in preclinical models [100].

Nicotine and direct activation of nAChRs on VTA dopamine neurons

Nicotine engages high-affinity nicotinic acetylcholine receptors (nAChRs) expressed on VTA dopamine (DA) neurons predominantly β2-containing assemblies (e.g., $\alpha4\beta2$, $\alpha6\beta2$ with $\alpha5/\beta3$ accessories) to increase firing rate, promote phasic bursting, and enhance synchrony across DA

populations. In parallel, nicotine modulates local afferents: rapid desensitization of $\alpha4\beta2$ nAChRs on VTA GABA neurons favors disinhibition, while $\alpha7$ nAChRs facilitate glutamatergic drive, together yielding a net rise in DA neuron excitability and dopamine release in nucleus accumbens. Genetic and circuit evidence underscore this mechanism. Mice lacking the $\beta2$ subunit failed to show nicotine-evoked DA activation or self-administration in dependence-like paradigms, and these deficits are rescued by $\beta2$ re-expression in the VTA; $\alpha4$ deletion blunts nicotine-induced bursting and limits self-administration, whereas loss of $\alpha5$ shifts dose–response to require higher nicotine for comparable DA activation [108, 109]. Regionally, posterior VTA (rich in $\alpha4/\alpha6/\beta3$ expression) exhibits stronger nicotine-evoked increases in DA firing and bursting than anterior VTA, aligning with the posterior VTA's greater reinforcing sensitivity to intra-VTA nicotine, as illustrated in Fig. 2. Collectively, these data establish that nicotine's acute reinforcing actions arise from direct nAChR-mediated excitation of DA neurons, complemented by nAChR-dependent control of local inhibitory and excitatory inputs, as illustrated in Fig. 3 [101].

Neuroadaptations in addiction

Addictive drugs leave enduring "traces" across mesocorticolimbic circuits that progressively shift behavior from goal-directed use to compulsive seeking and taking. These adaptations unfold over time from rapid synaptic changes after initial exposure to circuit- and gene-regulatory remodeling during abstinence, ultimately altering motivation, stress responsivity, and executive control. A complementary clinical framework is the allostatic model, in which negative affect and hyperkatifeia drive drug seeking via negative reinforcement as dependence develops [110].

Synaptic and circuit plasticity in VTA–NAc–striatal networks

Early exposures to drugs of abuse rapidly remodel glutamatergic synapses onto VTA dopamine neurons, providing a "teaching signal" that seeds longer-term circuit changes. Within hours of a first cocaine exposure in rodent models, VTA DA neurons show an increased AMPA/NMDA ratio and insertion of calcium-permeable, GluA2-lacking AMPARs; mGluR1 signaling can reverse this state, highlighting a labile early memory trace of drug exposure [111]. With repeated use and withdrawal, plasticity consolidates downstream in the nucleus accumbens (NAc): PFC→NAc and other inputs acquire silent synapses that later "mature," and CP-AMPARs accumulate during protracted abstinence, cellular changes causally linked to incubating cue-evoked drug seeking in dependence-like paradigms. Ensemble activity in the NAc shell increases across withdrawal and biases relapse propensity [112, 113]. Over longer timescales, addiction and chronic drug dependence reshape corticostriatal loops: direct (D1) and indirect (D2) medium spiny neuron pathways adopt cell-type–specific gene programs and synaptic rules that favor drug seeking and blunt aversion/suppression, and behavioral control shifts from ventral (NAc) to dorsal striatal circuits that encode habits and compulsion in rodent models and human neuroimaging studies [114]. In parallel with these dopamine-centered changes, drugs of abuse also induce distinct but complementary plasticity within VTA GABA microcircuits. Distinct GABAergic inputs onto VTA dopamine neurons undergo drug- and stress-induced alterations in synaptic strength and firing that reshape their inhibitory control over dopamine output and learning-related behavior in rodent models [97, 105]. Together with the better-characterized adaptations in VTA dopamine neurons, these inhibitory-circuit changes position VTA GABA plasticity as an integral component of the mesocorticolimbic "addiction engram," rather than a secondary consequence of dopamine-centered signaling [20, 24].

Glial, neuroimmune, and epigenetic "Scar" mechanisms

Neuroadaptations are not restricted to neurons. During withdrawal in rodent models of cocaine dependence, microglia are required for CP-AMPAR buildup at NAc synapses, and astrocytes show abstinence- and sex-dependent alterations in glutamate handling and synaptic coverage, pointing to glia as active drivers, not mere bystanders, of persistent plasticity [115, 116]. In parallel, drug exposure engages epigenetic programs that stabilize drug memories and vulnerability over weeks to months. Chromatin remodeling, DNA methylation, histone modifications, and noncoding RNAs reconfigure transcriptional landscapes in the NAc and related regions. Time-dependent DNA methylation increases in the NAc have been linked to incubated cocaine seeking in rodent dependence-like paradigms, and broader work catalogs histone/DNA modifications (e.g., acetylation, methylation, dopaminylation) across various substances [112, 117]. These glial and epigenetic




Fig. 3. The distribution of nicotinic acetylcholine receptor (nAChR) subunits on VTA DA, GABAergic, and glutamatergic neurons shows key afferent inputs and projection targets. DA neurons (green) receive glutamatergic input (yellow) from cortex/LDTg and GABAergic input (blue) from VP/NAc, while cholinergic afferents (purple) from PPTg/LDTg activate distinct nAChR subtypes. Outputs to NAc, PFC, and striatum mediate reward, motivation, and reinforcement.

mechanisms together act as "molecular and cellular scars" that maintain high relapse risk long after drug clearance, providing a substrate for persistent sensitivity to cues and stress.

Stress, Anti-reward systems, and translational targets

A further layer of adaptation involves recruitment of stress and anti-reward systems. Chronic exposure to drugs recruits corticotropin-releasing factor and dynorphin/κ-opioid signaling in extended amygdala circuits, producing hyperkatifeia and dysphoria; drug seeking is then maintained partly to relieve this negative state, reinforcing the cycle of dependence [118]. These insights have motivated translational strategies that target plasticity mechanisms rather than only acute drug effects, for example, restoring mGluR1/5 tone, normalizing CP-AMPARs, or modulating gene-regulatory programs to reduce incubated craving and relapse [119]. By integrating synaptic/circuit, glial–neuroimmune, and epigenetic adaptations with stress-system engagement, contemporary models view addiction as a distributed network disorder with multiple interlocking intervention points.

Therapeutic implications: Current treatment approaches

Pharmacological interventions

Medications that modulate mesolimbic dopamine often act on receptors that gate VTA output and remain first-line when approved for substance use disorders (SUDs). For alcohol use disorder (AUD), opioid antagonism with naltrexone reduces alcohol's ability to recruit β-endorphin–mediated disinhibition of VTA dopamine neurons and lowers accumbal dopamine, contributing to reduced drinking; across head-to-head reviews, naltrexone shows consistent, moderate efficacy [120]. Glutamate and GABA-directed agents also leverage VTA-linked circuitry: acamprosate mitigates withdrawal-related glutamatergic hyperexcitability (with debated molecular targets) and improves

abstinence outcomes. In contrast, topiramate (mixed GABAergic/glutamatergic actions) shows moderate evidence for reducing drinking days and heavy drinking [121, 122]. A GAB_A_B-receptor approach (baclofen) likely increases abstinent days and reduces relapse, particularly in detoxified patients though effects on heavy drinking are less consistent [123]. For nicotine, the $\alpha4\beta2$-partial agonist varenicline dampens nicotine-evoked VTA firing and striatal dopamine, supporting its clinical efficacy for cessation [124]. For stimulant use disorders, no medications are yet approved; circuit-informed adrenergic strategies (e.g., $\alpha1$-adrenergic antagonists such as doxazosin) have shown human laboratory signals but mixed/limited clinical data to date [93, 125].

Behavioral interventions

Behavioral therapies capitalize on reward learning to strengthen non-drug reinforcement and executive control. Contingency management (CM) produces reliable, objective improvements in abstinence (urine-verified) with benefits that can persist months after treatment; recent syntheses and federal guidance now endorse CM as an evidence-based standard [126, 127]. Cognitive-behavioral therapy (CBT) remains a core modality with small-to-moderate effects across alcohol and other drugs, and it pairs well with CM and pharmacotherapy [128].

Neuromodulation (emerging)

Noninvasive brain stimulation aimed at top-down control of mesolimbic circuits shows promise: meta-analyses report medium-to-large reductions in craving and substance use with repetitive TMS, particularly when targeting left dorsolateral prefrontal cortex over multiple sessions [129]. Deep brain stimulation (DBS) of VTA-connected nodes (most commonly nucleus accumbens/ventral capsule) is under active investigation for treatment-refractory cases; an exploratory randomized trial in AUD and recent reviews suggest feasibility and potential benefit, but evidence remains preliminary and not yet practice-changing [130, 131].

Future therapeutic directions

Precision medicine

Advances in imaging and genomics are enabling individualized treatment selection for substance use disorders (SUDs). Systematic frameworks now integrate behavioral, environmental, and biological markers to stratify patients and forecast outcomes. In particular, resting-state and task-based connectivity within mesocorticolimbic networks (including VTA–striatal–prefrontal links) are being used as predictive features and mechanistic readouts, while early trials relate medication effects (e.g., naltrexone or topiramate) to changes in reward-network connectivity. Genetic variation in receptors and channels that shape VTA function (e.g., CHRNA5 affecting nAChR physiology on VTA DA neurons) provides plausible pharmacogenomic levers for dosing, drug choice, and relapse risk models [132].

Circuit-specific interventions

Projection-defined studies in preclinical rodent models continue to map the most effective control points within VTA microcircuits. Optogenetic and chemogenetic manipulations reveal that inputs from ventral pallidum, nucleus accumbens, and rostromedial tegmental nucleus bidirectionally regulate reward seeking and cocaine-evoked adaptations, nominating these nodes and their synapses onto identified VTA cell types as translational targets for neuromodulation or targeted pharmacology [133, 134].

Epigenetic therapies

Drug exposure drives durable gene-regulatory changes across VTA- striatal circuits (histone acetylation/methylation, DNA methylation, noncoding RNAs). Recent reviews argue for “epidrugs” (e.g., histone deacetylase and DNA-methyltransferase inhibitors) as adjuncts to reverse drug dependence–linked transcriptional states observed in preclinical models and associated with substance use disorders in humans; preclinical data show normalization of synaptic plasticity and reduced drug seeking after such interventions [135].

Neurofeedback and brain-computer interfaces

Real-time fMRI neurofeedback (rt-fMRI-NF) is being tested to down- or up-regulate activity within reward circuits, including studies that explicitly train VTA regulation and report reductions in craving-related measures. Emerging reviews synthesize growing evidence for NF as a circuit-informed behavioral tool, and early trials explore closed-loop paradigms that detect craving-related patterns and deliver contingent feedback or stimulation [136].

Combination and algorithm-guided care

Converging work suggests the greatest benefit will come from rational combinations integrating

pharmacotherapies with behavioral treatments and neuromodulation selected by individual biomarker profiles (connectivity fingerprints, cue-reactivity signatures, genotype). Imaging-guided TMS programs illustrate how biomarkers can steer target selection and scheduling; similar algorithms are developed for integrating medications and behavioral plans around a patient's VTA-centric dysfunctions [137].

Conclusion

The VTA exerts wide-ranging control over motivation and learning through heterogeneous cell types and projection-specific signaling to limbic and cortical targets. Multi-transmitter co-release (dopamine, GABA, and glutamate) endows VTA outputs with fast synaptic and slower modulatory control, enabling flexible encoding of reward, aversion, and context. In parallel, closed-loop interactions with the hippocampus and prefrontal cortex bind spatiotemporal context to task rules, allowing state-dependent gating of behavior. These architectural and computational features explain how VTA activity confers incentive salience and generates prediction-error signals that train downstream circuits. Addictive drugs and drugs producing dependence-like states in preclinical models perturb these same motifs through class-specific actions on VTA neurons and afferents, seeding rapid synaptic changes that consolidate into enduring mesocorticolimbic adaptations. Recognizing addiction as a circuit-level disorder manifesting clinically as substance use disorders (SUDs) and informed by mechanistic insights from rodent models of drug dependence motivates multifaceted treatment strategies. Alongside established pharmacological and behavioral approaches, emerging interventions such as biomarker-guided neuromodulation, projection-informed circuit targets, epigenetic therapies, and neurofeedback aim to normalize dysregulated network dynamics. Future work should prioritize projection- and cell-type resolved biomarkers, longitudinal tracking of circuit function, and combinatorial, algorithm-guided care to match interventions to individual circuit phenotypes. Precision frameworks promise more durable remission by restoring balanced VTA-centered computation across reward, control, and context networks.

Declarations

Ethics approval and consent to participate

Not applicable as this study did not involve human participants, human data, or human tissue.

Consent for publication

Not applicable.

Availability of data and materials

Not applicable to this article.

Authors' contributions

The author conceptualized the study, performed the analysis, wrote the original draft, reviewed and edited the manuscript, and is the corresponding author. The author read and approved the final manuscript.

Acknowledgment

The authors gratefully acknowledge Ayman A. Fayek for providing the medical illustrations for this manuscript. Also the author would like to acknowledge Scribendi Company for editing the manuscript.

Disclosure

The author has no conflicts of interest, and the work was not supported or funded by any company or organization.

References

1. Cai J, Tong Q. Anatomy and function of ventral tegmental area glutamate neurons. Front Neural Circuits. 2022; 16:867053.
2. Zaharia C. Strain electricity. J Appl Phys. 1973;44(11): 5198–5199.
3. Saleh Y, Jarratt-Barnham I, Fernandez-Egea E, Husain M. Mechanisms underlying motivational dysfunction in schizophrenia. Front Behav Neurosci. 2021;15:709753.
4. Döbrössy MD, Ramanathan C, Ashouri Vajari D, Tong Y, Schlaepfer T, Coenen VA. Neuromodulation in psychiatric disorders: experimental and clinical evidence for reward and motivation network deep brain stimulation: focus on the medial forebrain bundle. Eur J Neurosci. 2021;53(1):89–113.
5. Valenti O, Rekawek KA, Wieser S, Bulut H, Scholze P, Boehm S. Plasticity of ventral tegmental area disturbance during abstinence after repeated amphetamine exposure: restoration by selective activation of group II metabotropic glutamate receptors. Front Pharmacol. 2025;16:1–17.
6. Huang D, Grady FS, Peltekian L, Geerling JC. Efferent projections of Vglut2, Foxp2, and Pdyn parabrachial neurons in mice. J Comp Neurol 2021;529(4):657–693.
7. Huang D, Grady FS, Peltekian L, Laing JJ, Geerling JC. Efferent projections of CGRP/Calca-expressing parabrachial neurons in mice. J Comp Neurol 2021;529(11):2911–2957.
8. Hou G, Hao M, Duan J, Han MH. The formation and function of the VTA dopamine system. Int J Mol Sci. 2024;25(7): 3875.

9. Trutti AC, Fontanesi L, Mulder MJ, Bazin PL, Hommel B, Forstmann BU. A probabilistic atlas of the human ventral tegmental area (VTA) based on 7 Tesla MRI data. Brain Struct Funct. 2021;226(4):1155–1167.
10. Prasad AA, Wallén-Mackenzie Å. Architecture of the subthalamic nucleus. Commun Biol. 2024;7(1):78.
11. Lu X, Xue J, Lai Y, Tang X. Heterogeneity of mesencephalic dopaminergic neurons: From molecular classifications, electrophysiological properties to functional connectivity. FASEB J. 2024;38(3):1–14.
12. Salmani bY, lahti l, gillberg l, Jacobsen jk, Mantas I, Svenningsson P, *et al.* Transcriptomic atlas of midbrain dopamine neurons uncovers differential vulnerability in a Parkinsonism lesion model. Elife. 2024;12:1–32.
13. Morales M, Margolis EB. Ventral tegmental area: cellular heterogeneity, connectivity and behaviour. Nat Rev Neurosci. 2017;18(2):73–85.
14. Holly EN, Miczek KA. Ventral tegmental area dopamine revisited: effects of acute and repeated stress. Psychopharmacology (Berl). 2016;233(2):163–186.
15. Soares-Cunha C, Heinsbroek JA. Ventral pallidal regulation of motivated behaviors and reinforcement. Front Neural Circuits. 2023;17:1086053.
16. Han Y, Ai L, Song S, Zhou Y, Chen D, Sha S, *et al.* Midbrain glutamatergic circuit mechanism of resilience to socially transferred allodynia in male mice. Nat Commun. 2024;15(1):4947.
17. Basso V, Döbrössy MD, Thompson LH, Kirik D, Fuller HR, Gates MA. State of the art in sub-phenotyping midbrain dopamine neurons. Biology (Basel). 2024;13(9):690.
18. Eskenazi D, Malave L, Mingote S, Yetnikoff L, Ztaou S, Velicu V, *et al.* Dopamine neurons that cotransmit glutamate, from synapses to circuits to behavior. Front Neural Circuits. 2021;15:665386.
19. Michel L, Molina P, Mameli M. The behavioral relevance of a modular organization in the lateral habenula. Neuron. 2024;112(16):2669–2685.
20. Bouarab C, Thompson B, Polter AM. VTA GABA neurons at the interface of stress and reward. Front Neural Circuits. 2019;13:78.
21. Merrill CB, Friend LN, Newton ST, Hopkins ZH, Edwards JG. Ventral tegmental area dopamine and GABA neurons: Physiological properties and expression of mRNA for endocannabinoid biosynthetic elements. Sci Rep. 2015;5(1): 16176.
22. Urrutia-Piñones J, Morales-Moraga C, Sanguinetti-González N, Escobar AP, Chiu CQ. Long-range GABAergic projections of cortical origin in brain function. Front Syst Neurosci. 2022;16:841869.
23. Wallace ML, Sabatini BL. Synaptic and circuit functions of multitransmitter neurons in the mammalian brain. Neuron. 2023;111(19):2969–2983.
24. Creed MC, Ntamati NR, Tan KR. VTA GABA neurons modulate specific learning behaviors through the control of dopamine and cholinergic systems. Front Behav Neurosci. 2014;8:8.
25. van Zessen R, Phillips JL, Budygin EA, Stuber GD. Activation of VTA GABA neurons disrupts reward consumption. Neuron. 2012;73(6):1184–1194.
26. Tan KR, Yvon C, Turiault M, Mirzabekov JJ, Doehner J, Labouèbe G, *et al.* GABA neurons of the VTA drive conditioned place aversion. Neuron. 2012;73(6):1173–1183.
27. Yu YH, Tsai AC, Ou CY, Cheng CN, Chang FC, Shyu BC, *et al.* Optogenetic stimulation in the medial prefrontal cortex modulates stimulus valence from rewarding and aversive to neutral states. Front Psychiatry. 2023;14:1119803.
28. Zhou K, Xu H, Lu S, Jiang S, Hou G, Deng X, *et al.* Reward and aversion processing by input-defined parallel nucleus accumbens circuits in mice. Nat Commun. 2022;13(1):6244.
29. Taylor SR, Badurek S, Dileone RJ, Nashmi R, Minichiello L, Picciotto MR. GABAergic and glutamatergic efferents of the mouse ventral tegmental area. J Comp Neurol. 2014;522(14):3308–3334.
30. McGovern DJ, Polter AM, Prévost ED, Ly A, McNulty CJ, Rubinstein B, *et al.* Ventral tegmental area glutamate neurons establish a mu-opioid receptor gated circuit to mesolimbic dopamine neurons and regulate opioid–seeking behavior. Neuropsychopharmacology. 2023;48(13):1889–1900.
31. Hnasko TS, Hjelmstad GO, Fields HL, Edwards RH. Ventral tegmental area glutamate neurons: electrophysiological properties and projections. J Neurosci. 2012;32(43):15076–15085.
32. Fitzgerald ND, Day JJ. Neuronal heterogeneity in the ventral tegmental area: distinct contributions to reward circuitry and motivated behavior. Addict Neurosci. 2025;14:100191.
33. Jiang Q, Bakhurin KI, Hughes RN, Lu B, Ruan S, Yin HH. GABAergic neurons in the ventral tegmental area represent and regulate force vectors. Cell Rep. 2025;44(2):115313.
34. Oriol L, Chao M, Kollman GJ, Dowlat DS, Singhal SM, Steinkellner T, *et al.* Ventral tegmental area interneurons revisited: GABA and glutamate projection neurons make local synapses. Elife. 2025;13:1–24.
35. Srinivasan S, Limani F, Hanzlova M, La Batide-Alanore S, Klotz S, Hnasko TS, *et al.* Evidence for low affinity of GABA at the vesicular monoamine transporter VMAT2 – Implications for transmitter co-release from dopamine neurons. Neuropharmacology. 2025;270:110367.
36. Patel JC, Sherpa AD, Melani R, Witkovsky P, Wiseman MR, O'Neill B, *et al.* GABA co-released from striatal dopamine axons dampens phasic dopamine release through autoregulatory GABAA receptors. Cell Rep. 2024;43(3):113834.
37. Miranda-Barrientos J, Chambers I, Mongia S, Liu B, Wang HL, Mateo-Semidey GE, *et al.* Ventral tegmental area GABA, glutamate, and glutamate-GABA neurons are heterogeneous in their electrophysiological and pharmacological properties. Eur J Neurosci. 2021;54(1):4061–4084.
38. Oliva I, Wanat MJ. Ventral tegmental area afferents and drug-dependent behaviors. Front Psychiatry. 2016;7:30.
39. Elum JE, Szelenyi ER, Juarez B, Murry AD, Loginov G, Zamorano CA, *et al.* Distinct dynamics and intrinsic properties in ventral tegmental area populations mediate reward association and motivation. Cell Rep. 2024;43(9):114668.
40. Xu Y, Lin Y, Yu M, Zhou K. The nucleus accumbens in reward and aversion processing: insights and implications. Front Behav Neurosci. 2024;18:1420028.
41. Marinescu AM, Labouesse MA. The nucleus accumbens shell: a neural hub at the interface of homeostatic and hedonic feeding. Front Neurosci. 2024;18:1437210.
42. King N, Dafny N. The ventral tegmental area (VTA), the nucleus accumbens (NAc), the caudate nucleus (CN) and the prefrontal cortex (PFC) role in the response to acute and chronic methylphenidate. J Exp Neurol. 2023;4(1):21–36.
43. Ebrahimi MN, Banazadeh M, Alitaneh Z, Suha AJ, Esmaeili A, Hasannejad-Asl B, *et al.* The distribution of neurotransmitters in the brain circuitry: Mesolimbic pathway and addiction. Physiol Behav. 2024;284:114639.
44. Zhong P, Qin L, Yan Z. Dopamine differentially regulates response dynamics of prefrontal cortical principal neurons and interneurons to optogenetic stimulation of inputs from ventral tegmental area. Cereb Cortex. 2020;30(8):4402–4409.
45. Tritsch NX, Sabatini BL. Dopaminergic modulation of synaptic transmission in cortex and striatum. Neuron. 2012;76(1): 33–50.

46. Carr DB, Sesack SR. Projections from the rat prefrontal cortex to the ventral tegmental area: target specificity in the synaptic associations with mesoaccumbens and mesocortical neurons. J Neurosci. 2000;20(10):3864–3873.
47. Geisler S, Zahm DS. Afferents of the ventral tegmental area in the rat-anatomical substratum for integrative functions. J Comp Neurol. 2005;490(3):270–294.
48. Li M, Yang G. A mesocortical glutamatergic pathway modulates neuropathic pain independent of dopamine co-release. Nat Commun. 2024;15(1):643.
49. Li Y, Zhang B, Pan X, Wang Y, Xu X, Wang R, *et al.* Dopamine-mediated major depressive disorder in the neural circuit of ventral tegmental area-nucleus accumbens-medial prefrontal cortex: From biological evidence to computational models. Front Cell Neurosci. 2022;16:923039.
50. Kong MS, Zweifel LS. Central amygdala circuits in valence and salience processing. Behav Brain Res. 2021;410:113355.
51. Avegno EM, Gilpin NW. Reciprocal midbrain-extended amygdala circuit activity in preclinical models of alcohol use and misuse. Neuropharmacology. 2022;202:108856.
52. Sladky R, Kargl D, Haubensak W, Lamm C. An active inference perspective for the amygdala complex. Trends Cogn Sci. 2024;28(3):223–236.
53. Liu A, Cheng Y, Huang J. Neurons innervating both the central amygdala and the ventral tegmental area encode different emotional valences. Front Neurosci. 2023;17:1178693.
54. Xin J, Shan W, Li J, Yu H, Zuo B. Activation of the lateral habenula-ventral tegmental area neural circuit contributes to postoperative cognitive dysfunction in mice. Adv Sci. 2022;9(22):2202228.
55. Cerniauskas I, Winterer J, W de Jong J, Lukacsovich D, Yang H, Fawwad Khan F, *et al.* Chronic stress induces activity, synaptic, and transcriptional remodeling of the lateral habenula associated with deficits in motivated behaviors. Neuron. 2019;104(5):899–915.
56. Hones VI, Mizumori SJY. Response flexibility: the role of the lateral habenula. Front Behav Neurosci. 2022;16:852235.
57. Sun Y, Cao J, Xu C, Sun J, Liu X, Shi Z, *et al.* Lateral habenula-rostromedial tegmental nucleus circuit mediates inflammatory pain in mice. J Headache Pain. 2025;26(1):105.
58. Godfrey N, Qiao M, Borgland SL. Activation of LH GABAergic inputs counteracts fasting-induced changes in tVTA/RMTG neurons. J Physiol. 2022;600(9):2203–2224.
59. Liu Y, Wu K, Dong Y, Jia R, Chen X, Ge A, *et al.* Lateral habenula induces cognitive and affective dysfunctions in mice with neuropathic pain via an indirect pathway to the ventral tegmental area. Neuropsychopharmacology. 2025;50(7):1039–1050.
60. Vincent KF, Zhang ER, cho AJ, Kato-Miyabe R, Mallari OG, Moody OA, *et al.* Electrical stimulation of the ventral tegmental area restores consciousness from sevoflurane-, dexmedetomidine-, and fentanyl-induced unconsciousness in rats. Brain Stimul. 2024;17(3):687–697.
61. Bracht T, Mertse N, Walther S, Lüdi K, Breit S, Federspiel A, *et al.* Link between structural connectivity of the medial forebrain bundle, functional connectivity of the ventral tegmental area, and anhedonia in unipolar depression. NeuroImage Clin. 2022;34:102961.
62. Blum K, Bowirrat A, Sunder K, Thanos PK, Hanna C, Gold MS, *et al.* Dopamine dysregulation in reward and autism spectrum disorder. Brain Sci. 2024 Jul;14(7):733.
63. Vander Weele CM, Siciliano CA, Tye KM. Dopamine tunes prefrontal outputs to orchestrate aversive processing. Brain Res. 2019;1713:16–31.
64. Hui M, Beier KT. Defining the interconnectivity of the medial prefrontal cortex and ventral midbrain. Front Mol Neurosci. 2022;15:971349.
65. Sayegh FJ, Mouledous L, Macri C, Macedo JP, Lejards C, Rampon C. Ventral tegmental area dopamine projections to the hippocampus trigger long-term potentiation and contextual learning. Nat Commun. 2024;15(1):4100.
66. Rizzi-Wise CA, Wang DV. Putting together pieces of the lateral septum: multifaceted functions and its neural pathways. eNeuro. 2021;8(6). ENEURO.0315-21.2021.
67. Yoo JH, Zell V, Gutierrez-Reed N, Wu J, Ressler R, Shenasa MA, *et al.* Ventral tegmental area glutamate neurons co-release GABA and promote positive reinforcement. Nat Commun. 2016;7(1):13697.
68. Phillips RA, Tuscher JJ, Black SL, Andraka E, Fitzgerald ND, Ianov L, *et al.* An atlas of transcriptionally defined cell populations in the rat ventral tegmental area. Cell Rep. 2022;39(1):110616.
69. Barbano MF, Qi J, Chen E, Mohammad U, Espinoza O, Candido M, *et al.* VTA glutamatergic projections to the nucleus accumbens suppress psychostimulant-seeking behavior. Neuropsychopharmacology. 2024;49(12):1905–1915.
70. Zeidler Z, Gomez MF, Gupta TA, Shari M, Wilke SA, DeNardo LA. Dopaminergic projections to the prefrontal cortex are critical for rapid threat avoidance learning. Curr Biol. 2025;35(17):4259–4269.
71. Smith DM, Torregrossa MM. The ventral tegmental area dopamine to basolateral amygdala projection supports acquisition of cocaine self-administration. Neuropharmacology. 2024;261:110160.
72. Morel C, Montgomery SE, Li L, Cuttoli RD, Teichman EM, Juarez B, *et al.* Midbrain projection to the basolateral amygdala encodes anxiety-like but not depression-like behaviors. Nat Commun. 2022;13(1):1532.
73. Mahadevia D, Saha R, Manganaro A, Chuhma N, Annette Ziolkowski-Blake AZ, Morgan AA, *et al.* Dopamine promotes aggression in mice via ventral tegmental area to lateral septum projections. Nat Commun. 2021;12(1):6796.
74. Grove JCR, Gray LA, La Santa Medina N, Sivakumar N, Ahn JS, Corpuz TV, *et al.* Dopamine subsystems that track internal states. Nature. 2022;608(7922):374–380.
75. Deng Y, Song D, Ni J, Qing H, Quan Z. Reward prediction error in learning-related behaviors. Front Neurosci. 2023;17:1171612.
76. Solié C, Girard B, Righetti M, Tapparel M, Bellone C. VTA dopamine neuron activity encodes social interaction and promotes reinforcement learning through social prediction error. Nat Neurosci. 2022;25(1):86–97.
77. Kaushik P, Naudé J, Raju SB, Alexandre F. A VTA GABAergic computational model of dissociated reward prediction error computation in classical conditioning. Neurobiol Learn Mem. 2022;193:107653.
78. Bromberg-Martin ES, Matsumoto M, Hikosaka O. Dopamine in motivational control: rewarding, aversive, and alerting. Neuron. 2010;68(5):815–834.
79. Salamone JD, Correa M. The mysterious motivational functions of mesolimbic dopamine. Neuron. 2012;76(3):470–485.
80. Berridge KC. Motivation concepts in behavioral neuroscience. Physiol Behav. 2004;81(2):179–209.
81. Robinson T. The neural basis of drug craving: an incentive-sensitization theory of addiction. Brain Res Rev. 1993;18(3):247–291.
82. Wolf ME. Synaptic mechanisms underlying persistent cocaine craving. Nat Rev Neurosci. 2016;17:351–365.

83. Lüscher C, Malenka RC. Drug-evoked synaptic plasticity in addiction: from molecular changes to circuit remodeling. Neuron. 2011;69(4):650–663.
84. Ramirez-Mejia G, Gil-Lievana E, Urrego-Morales O, Galvez-Marquez D, Hernández-Ortiz E, Carrillo-Lorenzo JA, *et al.* Salience to remember: VTA-IC dopaminergic pathway activity is necessary for object recognition memory formation. Neuropharmacology. 2023;228:109464.
85. Lisman JE, Grace AA. The hippocampal-VTA loop: controlling the entry of information into long-term memory. Neuron. 2005;46(5):703–713.
86. Flores-García M, Rizzo A, Garçon-Poca MZ, Fernández-Dueñas V, Bonaventura J. Converging circuits between pain and depression: the ventral tegmental area as a therapeutic hub. Front Pharmacol. 2023;14:1278023.
87. Wang Y, Zhang H, Cui J, Zhang J, Yin F, Guo H, *et al.* Opiate-associated contextual memory formation and retrieval are differentially modulated by dopamine D1 and D2 signaling in hippocampal–prefrontal connectivity. Neuropsychopharmacology. 2019;44(2):334–343.
88. Yu TH, Wu YJ, Chien ME, Hsu KS. Transcranial direct current stimulation induces hippocampal metaplasticity mediated by brain-derived neurotrophic factor. Neuropharmacology. 2019;144(1):358–367.
89. Pereira AR, Alemi M, Cerqueira-Nunes M, Monteiro C, Galhardo V, Cardoso-Cruz H. Dynamics of lateral habenula–ventral tegmental area microcircuit on pain-related cognitive dysfunctions. Neurol Int. 2023;15(4):1303–1319.
90. Howell LL, Kimmel HL. Monoamine transporters and psychostimulant addiction. Biochem Pharmacol. 2008;75(1):196–217.
91. dela Peña I, Gevorkiana R, Shi WX. Psychostimulants affect dopamine transmission through both dopamine transporter-dependent and independent mechanisms. Eur J Pharmacol. 2015;764:562–570.
92. Ventura R, Cabib S, Alcaro A, Orsini C, Puglisi-Allegra S. Norepinephrine in the prefrontal cortex is critical for amphetamine-induced reward and mesoaccumbens dopamine release. J Neurosci. 2003;23(5):1879–1885.
93. Newton TF, De La Garza R, Brown G, Kosten TR, Mahoney JJ, Haile CN. Noradrenergic α1 receptor antagonist treatment attenuates positive subjective effects of cocaine in humans: a randomized trial. PLoS One. 2012;7(2):e30854.
94. Johnson S, North R. Opioids excite dopamine neurons by hyperpolarization of local interneurons. J Neurosci. 1992;12(2):483–488.
95. Mathon DS, Lesscher HMB, Gerrits MAF, Kamal A, Pintar JE, Schuller AG, *et al.* Increased GABAergic input to ventral tegmental area dopaminergic neurons associated with decreased cocaine reinforcement in μ-opioid receptor knockout mice. Neuroscience. 2005;130(2):359–367.
96. Jhou TC, Fields HL, Baxter MG, Saper CB, Holland PC. The rostromedial tegmental nucleus (RMTg), a GABAergic afferent to midbrain dopamine neurons, encodes aversive stimuli and inhibits motor responses. Neuron. 2009;61(5):786–800.
97. Matsui A, Jarvie BC, Robinson BG, Hentges ST, Williams JT. Separate GABA afferents to dopamine neurons mediate acute action of opioids, development of tolerance, and expression of withdrawal. Neuron. 2014;(82):1346–1356.
98. Maldonado R, Valverde O, Berrendero F. Involvement of the endocannabinoid system in drug addiction. Trends Neurosci. 2006;29(4):225–232.
99. Cheer JF, Wassum KM, Heien MLAV, Phillips PEM, Wightman RM. Cannabinoids enhance subsecond dopamine release in the nucleus accumbens of awake rats. J Neurosci. 2004;24(18):4393–4400.
100. You C, Vandegrift B, Brodie MS. Ethanol actions on the ventral tegmental area: novel potential targets on reward pathway neurons. Psychopharmacology (Berl). 2018;235(6):1711–1726.
101. Faure P, Tolu S, Valverde S, Naudé J. Role of nicotinic acetylcholine receptors in regulating dopamine neuron activity. Neuroscience. 2014;282:86–100.
102. Di Chiara G, Imperato A. Drugs abused by humans preferentially increase synaptic dopamine concentrations in the mesolimbic system of freely moving rats. Proc Natl Acad Sci U S A. 1988;85(14):5274–5278.
103. Deng C, Li KY, Zhou C, Ye JH. Ethanol enhances glutamate transmission by retrograde dopamine signaling in a postsynaptic neuron/synaptic bouton preparation from the ventral tegmental area. Neuropsychopharmacology. 2009;34(5):1233–1244.
104. Larsson A, Edström L, Svensson L, Söderpalm B, Engel JA. Voluntary ethanol intake increases extracellular acetylcholine levels in the ventral tegmental area in the rat. Alcohol Alcohol. 2005;40(5):349–358.
105. Stobbs SH, Ohran AJ, Lassen MB, Allison DW, Brown JE, Steffensen SC. Ethanol suppression of ventral tegmental area GABA neuron electrical transmission involves N-methyl-D-aspartate receptors. J Pharmacol Exp Ther. 2004;311(1):282–289.
106. Ye JH, Tao L, Zhu L, Krnjević K, McArdle JJ. Ethanol inhibition of glycine-activated responses in neurons of ventral tegmental area of neonatal rats. J Neurophysiol. 2001;86(5):2426–2434.
107. Brodie MS, Shefner SA, Dunwiddie TV. Ethanol increases the firing rate of dopamine neurons of the rat ventral tegmental area in vitro. Brain Res. 1990;508(1):65–69.
108. Mathon DS, Lesscher HMB, Gerrits MAF, Kamal A, Pintar JE, Schuller AG, *et al.* Nicotine reinforcement and cognition restored by targeted expression of nicotinic receptors. Nature. 2005;436(7047):103–107.
109. Picciotto MR, Zoli M, Rimondini R, Léna C, Marubio LM, Pich EM, *et al.* Acetylcholine receptors containing the β2 subunit are involved in the reinforcing properties of nicotine. Nature. 1998;391(6663):173–177.
110. Poisson CL, Engel L, Saunders BT. Dopamine circuit mechanisms of addiction-like behaviors. Front Neural Circuits. 2021;15:752420.
111. Luscher C. Cocaine-evoked synaptic plasticity of excitatory transmission in the ventral tegmental area. Cold Spring Harb Perspect Med. 2013;3(5):a012013.
112. He G, Huai Z, Jiang C, Huang B, Tian Z, Le Q, *et al.* Persistent increase of accumbens cocaine ensemble excitability induced by IRK downregulation after withdrawal mediates the incubation of cocaine craving. Mol Psychiatry. 2023;28(1):448–462.
113. Zinsmaier AK, Dong Y, Huang YH. Cocaine-induced projection-specific and cell type-specific adaptations in the nucleus accumbens. Mol Psychiatry. 2022;27(1):669–686.
114. Lipton DM, Gonzales BJ, Citri A. Dorsal striatal circuits for habits, compulsions and addictions. Front Syst Neurosci. 2019;13:28.
115. Kim R, Testen A, Harder EV, Brown NE, Witt EA, Bellinger TJ, *et al.* Abstinence-dependent effects of long-access cocaine self-administration on nucleus accumbens astrocytes are observed in male, but not female, rats. eNeuro. 2022;9(5). ENEURO.0310-22.2022.
116. Reverte I, Marchetti C, Pezza S, Zenoni SF, Scaringi G, Ferrucci L, *et al.* Microglia-mediated calcium-permeable AMPAR accumulation in the nucleus accumbens drives hyperlocomotion during cocaine withdrawal. Brain Behav Immun. 2024;115:535–542.

117. Liu M, Si Z. An update: epigenetic mechanisms underlying methamphetamine addiction. Front Cell Dev Biol. 2024;12:1494557.
118. Koob GF. Drug addiction: hyperkatifeia/negative reinforcement as a framework for medications development. Pharmacol Rev. 2021;73(1):163–201.
119. Wolf ME. Targeting neuroplasticity in substance use disorders: implications for therapeutics. Annu Rev Pharmacol Toxicol. 2025;65(1):259–280.
120. Hood LE, Leyrer-Jackson JM, Olive MF. Pharmacotherapeutic management of co-morbid alcohol and opioid use. Expert Opin Pharmacother. 2020;21(7):823–839.
121. Spanagel R, Vengeliene V, Jandeleit B, Fischer W, Grindstaff K, Zhang X, *et al.* Acamprosate produces its antirelapse effects via calcium. Neuropsychopharmacology. 2014;39(4):783–791.
122. Kalk NJ, Lingford-Hughes AR. The clinical pharmacology of acamprosate. Br J Clin Pharmacol. 2014;77(2):315–323.
123. Agabio R, Saulle R, Rösner S, Minozzi S. Baclofen for alcohol use disorder. Cochrane Database Syst Rev. 2023;1: CD012557.
124. Goldstein N, Carty JRE, Betley JN. Specificity of varenicline in blocking mesolimbic circuit activation to natural and drug rewards. Neuroscience. 2022;483:40–51.
125. Shorter D, Lindsay JA, Kosten TR. The alpha-1 adrenergic antagonist doxazosin for treatment of cocaine dependence: a pilot study. Drug Alcohol Depend. 2013;131(1–2): 66–70.
126. Beaumont J. Clinical neuropsychology in rehabilitation. In: Stokes M, editor. Physical management in neurological rehabilitation. 2nd ed. Oxford: Elsevier;2004. p. 461–468.
127. Petry NM. Contingency management treatments: controversies and challenges. Addiction. 2010;105(9):1507–1509.
128. McHugh RK, Hearon BA, Otto MW. Cognitive behavioral therapy for substance use disorders. Psychiatr Clin North Am. 2010;33(3):511–525.
129. Mehta DD, Praecht A, Ward HB, Sanches M, Sorkhou M, Tang VM, *et al.* A systematic review and meta-analysis of neuromodulation therapies for substance use disorders. Neuropsychopharmacology. 2024;49(4):649–680.
130. Zammit Dimech D, Zammit Dimech AA, Hughes M, Zrinzo L. A systematic review of deep brain stimulation for substance use disorders. Transl Psychiatry. 2024;14(1):361.
131. Bach P, Luderer M, Müller UJ, Jakobs M, Baldermann JC, Voges J, *et al.* Deep brain stimulation of the nucleus accumbens in treatment-resistant alcohol use disorder: a double-blind randomized controlled multi-center trial. Transl Psychiatry. 2023;13(1):49.
132. Logge WB, Haber PS, Hurzeler T, Gallagher H, Kranzler H, Morley KC. Neural cue reactivity and intrinsic functional connectivity in individuals with alcohol use disorder following treatment with topiramate or naltrexone. Psychopharmacology (Berl). 2025;242(7):1641–1652.
133. Palmer D, Cayton CA, Scott A, Lin I, Newell B, Paulson A, *et al.* Ventral pallidum neurons projecting to the ventral tegmental area reinforce but do not invigorate reward-seeking behavior. Cell Rep. 2024;43(1):113669.
134. Khayat A, Yaka R. Activation of RMTg projections to the VTA reverse cocaine-induced molecular adaptation in the reward system. Transl Psychiatry. 2024;14(1):40.
135. Anderson EM, Taniguchi M. Epigenetic effects of addictive drugs in the nucleus accumbens. Front Mol Neurosci. 2022;15:828055.
136. Gu A, Chan CL, Xu X, Dexter JP, Becker B, Zhao Z. Real-time fMRI neurofeedback modulation of dopaminergic midbrain activity in young adults with elevated internet gaming disorder risk: randomized controlled trial. J Med Internet Res. 2025;27:e64687.
137. Sahay S, Reddy MVSRR, Lennox C, Wolinsky E, McCullumsmith RE, Singh T. Harnessing neuroimaging-guided transcranial magnetic stimulation for precision therapy in substance use disorders. Mol Psychiatry. 2025;30(8): 3804–3816.