

Weak Convergence of a Metropolis Algorithm for Bimodal Target Distributions

Michaël Lalancette¹ Mylène Bédard²

¹Department of Statistical Sciences, University of Toronto,
lalancette@utstat.toronto.edu

²Département de Mathématiques et de Statistique, Université de Montréal,
bedard@dms.umontreal.ca

June 4, 2018

- 1 Random Walk Metropolis algorithm
- 2 Weak convergence of our algorithm
- 3 Summary and Future work

- 1 Random Walk Metropolis algorithm
 - The Random Walk Metropolis algorithm
 - Optimal scaling
 - The problem with bimodal distributions
 - Solution: A new instrumental distribution
- 2 Weak convergence of our algorithm
- 3 Summary and Future work

MCMC in a nutshell

- Probability measure Π

MCMC in a nutshell

- Probability measure Π
- Want to estimate (finite) expectation $\int h d\Pi$

MCMC in a nutshell

- Probability measure Π
- Want to estimate (finite) expectation $\int h d\Pi$
- Idea: create a Markov chain X_1, X_2, \dots that has target distribution Π

MCMC in a nutshell

- Probability measure Π
- Want to estimate (finite) expectation $\int h d\Pi$
- Idea: create a Markov chain X_1, X_2, \dots that has target distribution Π
- If the chain is well behaved (irreducible, aperiodic, Harris recurrent),

$$\frac{1}{n} \sum_{i=1}^n h(X_i) \longrightarrow \int h d\Pi \quad \text{a.s.}$$

MCMC in a nutshell

- Probability measure Π
- Want to estimate (finite) expectation $\int h d\Pi$
- Idea: create a Markov chain X_1, X_2, \dots that has target distribution Π
- If the chain is well behaved (irreducible, aperiodic, Harris recurrent),

$$\frac{1}{n} \sum_{i=1}^n h(X_i) \longrightarrow \int h d\Pi \quad \text{a.s.}$$

- In general, less efficient than iid sampling, but iid sampling is sometimes unrealizable

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d
- Initialize \mathbf{X}_0

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d
- Initialize \mathbf{X}_0
- Given \mathbf{X}_t , independently generate \mathbf{Y} *symmetric around 0*

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d
- Initialize \mathbf{X}_0
- Given \mathbf{X}_t , independently generate \mathbf{Y} *symmetric around 0*
- Set

$$\mathbf{X}_{t+1} = \begin{cases} \mathbf{X}_t + \mathbf{Y}, & \text{w.p. } \alpha_t \\ \mathbf{X}_t, & \text{w.p. } 1 - \alpha_t \end{cases}$$

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d
- Initialize \mathbf{X}_0
- Given \mathbf{X}_t , independently generate \mathbf{Y} *symmetric around 0*
- Set

$$\mathbf{X}_{t+1} = \begin{cases} \mathbf{X}_t + \mathbf{Y}, & \text{w.p. } \alpha_t \\ \mathbf{X}_t, & \text{w.p. } 1 - \alpha_t \end{cases}$$

- $\alpha_t = \min \left\{ 1, \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)} \right\}$

The Random Walk Metropolis algorithm

- RWM is a very simple MCMC algorithm
- π is a (possibly unnormalized) density on \mathbb{R}^d
- Initialize \mathbf{X}_0
- Given \mathbf{X}_t , independently generate \mathbf{Y} *symmetric around 0*
- Set

$$\mathbf{X}_{t+1} = \begin{cases} \mathbf{X}_t + \mathbf{Y}, & \text{w.p. } \alpha_t \\ \mathbf{X}_t, & \text{w.p. } 1 - \alpha_t \end{cases}$$

- $\alpha_t = \min \left\{ 1, \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)} \right\}$
- Called *accept/reject step*

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration
- And large $\sigma \Rightarrow \mathbf{X}_t + \mathbf{Y}$ far from the mode

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration
- And large $\sigma \Rightarrow \mathbf{X}_t + \mathbf{Y}$ far from the mode $\Rightarrow \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ small

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration
- And large $\sigma \Rightarrow \mathbf{X}_t + \mathbf{Y}$ far from the mode $\Rightarrow \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ small \Rightarrow Most steps are rejected

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration
- And large $\sigma \Rightarrow \mathbf{X}_t + \mathbf{Y}$ far from the mode $\Rightarrow \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ small \Rightarrow Most steps are rejected \Rightarrow Slow exploration

The Random Walk Metropolis algorithm

- Under mild conditions on instrumental distribution (distribution of increments \mathbf{Y}),

$$\frac{1}{n} \sum_{t=1}^n h(\mathbf{X}_t) \longrightarrow \frac{\int h(\mathbf{x}) \pi(\mathbf{x}) d\mathbf{x}}{\int \pi(\mathbf{x}) d\mathbf{x}} \quad \text{a.s.}$$

- Usually $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- In theory, works for any $\sigma > 0$...
- But small $\sigma \Rightarrow$ Small steps \Rightarrow Slow exploration
- And large $\sigma \Rightarrow \mathbf{X}_t + \mathbf{Y}$ far from the mode $\Rightarrow \frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ small \Rightarrow Most steps are rejected \Rightarrow Slow exploration
- Must choose σ carefully

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$
- Let $\sigma = \frac{\ell}{\sqrt{d}}$ (Large dimension \Rightarrow Small steps)

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$
- Let $\sigma = \frac{\ell}{\sqrt{d}}$ (Large dimension \Rightarrow Small steps)
- Let $\mathbf{X}^{(d)}(t)$ be Markov chain obtained (with step size depending on dimension d)

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$
- Let $\sigma = \frac{\ell}{\sqrt{d}}$ (Large dimension \Rightarrow Small steps)
- Let $\mathbf{X}^{(d)}(t)$ be Markov chain obtained (with step size depending on dimension d)
- And $X_j^{(d)}(t)$ its j -th component

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$
- Let $\sigma = \frac{\ell}{\sqrt{d}}$ (Large dimension \Rightarrow Small steps)
- Let $\mathbf{X}^{(d)}(t)$ be Markov chain obtained (with step size depending on dimension d)
- And $X_j^{(d)}(t)$ its j -th component
- Smaller steps \Rightarrow More steps needed, so accelerate chain:
 $Z_j^{(d)}(t) = X_j^{(d)}(\lfloor dt \rfloor)$

Optimal scaling

- [Roberts et al., 1997] proposed optimal scaling (in simplified framework)
- Assume $\pi(x) = \prod_{i=1}^d f(x_i)$
- Let $\sigma = \frac{\ell}{\sqrt{d}}$ (Large dimension \Rightarrow Small steps)
- Let $\mathbf{X}^{(d)}(t)$ be Markov chain obtained (with step size depending on dimension d)
- And $X_j^{(d)}(t)$ its j -th component
- Smaller steps \Rightarrow More steps needed, so accelerate chain:
 $Z_j^{(d)}(t) = X_j^{(d)}(\lfloor dt \rfloor)$
- Let $d \rightarrow \infty$

Accelerating the Markov chain




Figure: Trace of the accelerated first component of the Markov chain, $Z_1^{(d)}$, for different values of d . The target distribution is multivariate standard Gaussian.

Optimal scaling

- Find that $Z_j^{(d)}$ converges (weakly in Skorokhod topology) to Langevin diffusion with *speed measure* $\nu(\ell) = 2\ell^2 \Phi\left(-\frac{\ell\sqrt{B}}{2}\right)$ and stationary distribution f

Optimal scaling

- Find that $Z_j^{(d)}$ converges (weakly in Skorokhod topology) to Langevin diffusion with *speed measure* $\nu(\ell) = 2\ell^2 \Phi\left(-\frac{\ell\sqrt{B}}{2}\right)$ and stationary distribution f
- B is unknown parameter of f

Optimal scaling

- Find that $Z_j^{(d)}$ converges (weakly in Skorokhod topology) to Langevin diffusion with *speed measure* $\nu(\ell) = 2\ell^2 \Phi\left(-\frac{\ell\sqrt{B}}{2}\right)$ and stationary distribution f
- B is unknown parameter of f
- Find a value $\hat{\ell}$ that maximizes $\nu(\ell)$

Optimal scaling

- Find that $Z_j^{(d)}$ converges (weakly in Skorokhod topology) to Langevin diffusion with *speed measure* $v(\ell) = 2\ell^2 \Phi\left(-\frac{\ell\sqrt{B}}{2}\right)$ and stationary distribution f
- B is unknown parameter of f
- Find a value $\hat{\ell}$ that maximizes $v(\ell)$
- $\hat{\ell}$ is the only value that makes the asymptotic acceptance probability 0.234 (regardless of B)!

Optimal scaling

- Find that $Z_j^{(d)}$ converges (weakly in Skorokhod topology) to Langevin diffusion with *speed measure* $v(\ell) = 2\ell^2 \Phi\left(-\frac{\ell\sqrt{B}}{2}\right)$ and stationary distribution f
- B is unknown parameter of f
- Find a value $\hat{\ell}$ that maximizes $v(\ell)$
- $\hat{\ell}$ is the only value that makes the asymptotic acceptance probability 0.234 (regardless of B)!
- So just simulate a few short runs and tune σ so that acceptance rate is roughly 0.234

The problem with bimodal distributions

- Suppose the target density π (on \mathbb{R}^d) has two distinct modes with a "hole" in between (low density region)

The problem with bimodal distributions

- Suppose the target density π (on \mathbb{R}^d) has two distinct modes with a "hole" in between (low density region)
- Optimal scaling strategy tends to favor local exploration (Relatively small σ)

The problem with bimodal distributions

- Suppose the target density π (on \mathbb{R}^d) has two distinct modes with a "hole" in between (low density region)
- Optimal scaling strategy tends to favor local exploration (Relatively small σ)
- Small steps \Rightarrow Almost impossible to cross the hole in 1 step

The problem with bimodal distributions

- Suppose the target density π (on \mathbb{R}^d) has two distinct modes with a "hole" in between (low density region)
- Optimal scaling strategy tends to favor local exploration (Relatively small σ)
- Small steps \Rightarrow Almost impossible to cross the hole in 1 step
- Almost impossible to accept steps into the hole ($\frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ very small)

The problem with bimodal distributions

- Suppose the target density π (on \mathbb{R}^d) has two distinct modes with a "hole" in between (low density region)
- Optimal scaling strategy tends to favor local exploration (Relatively small σ)
- Small steps \Rightarrow Almost impossible to cross the hole in 1 step
- Almost impossible to accept steps into the hole ($\frac{\pi(\mathbf{X}_t + \mathbf{Y})}{\pi(\mathbf{X}_t)}$ very small)
- Chain gets stuck in a mode and never explores the other one

Solution: A new instrumental distribution

- Previously, $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$

Solution: A new instrumental distribution

- Previously, $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- Now, let

$$\mathbf{Y} \sim \begin{cases} N_d(0, \sigma^2 I_d), & \text{w.p. } 1 - p \\ \mathcal{D}, & \text{w.p. } p \end{cases},$$

where \mathcal{D} is any distribution on \mathbb{R}^d *symmetric around 0*, $p \in (0, 1)$

Solution: A new instrumental distribution

- Previously, $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- Now, let

$$\mathbf{Y} \sim \begin{cases} N_d(0, \sigma^2 I_d), & \text{w.p. } 1 - p \\ \mathcal{D}, & \text{w.p. } p \end{cases},$$

where \mathcal{D} is any distribution on \mathbb{R}^d *symmetric around 0*, $p \in (0, 1)$

- In practice, choose \mathcal{D} to favor switching modes

Solution: A new instrumental distribution

- Previously, $\mathbf{Y} \sim N_d(0, \sigma^2 I_d)$
- Now, let

$$\mathbf{Y} \sim \begin{cases} N_d(0, \sigma^2 I_d), & \text{w.p. } 1 - p \\ \mathcal{D}, & \text{w.p. } p \end{cases},$$

where \mathcal{D} is any distribution on \mathbb{R}^d *symmetric around 0*, $p \in (0, 1)$

- In practice, choose \mathcal{D} to favor switching modes
- Turns the "slowly mixing" chain into a "rapidly mixing" chain ([Guan and Krone, 2007])

- 1 Random Walk Metropolis algorithm
- 2 Weak convergence of our algorithm
 - Framework
 - The limiting processes
- 3 Summary and Future work

Framework

- Assume $\pi(x) = f_1(x_1) \prod_{i=2}^d f(x_i)$

Framework

- Assume $\pi(x) = f_1(x_1) \prod_{i=2}^d f(x_i)$
- Instrumental distribution \mathcal{D} : $Y_1 \sim \mathcal{D}_1$, $Y_2, \dots, Y_d \sim N(0, \sigma^2)$, all independent

Framework

- Assume $\pi(x) = f_1(x_1) \prod_{i=2}^d f(x_i)$
- Instrumental distribution \mathcal{D} : $Y_1 \sim \mathcal{D}_1$, $Y_2, \dots, Y_d \sim N(0, \sigma^2)$, all independent
- Scale both $\sigma = \frac{\ell}{\sqrt{d}}$ and $p = 1 \wedge \frac{\beta}{d}$, but not \mathcal{D}_1

Framework

- Assume $\pi(x) = f_1(x_1) \prod_{i=2}^d f(x_i)$
- Instrumental distribution \mathcal{D} : $Y_1 \sim \mathcal{D}_1$, $Y_2, \dots, Y_d \sim N(0, \sigma^2)$, all independent
- Scale both $\sigma = \frac{\ell}{\sqrt{d}}$ and $p = 1 \wedge \frac{\beta}{d}$, but not \mathcal{D}_1
- Accelerate the chain by a factor of d (like before)

Framework

- Assume $\pi(x) = f_1(x_1) \prod_{i=2}^d f(x_i)$
- Instrumental distribution \mathcal{D} : $Y_1 \sim \mathcal{D}_1$, $Y_2, \dots, Y_d \sim N(0, \sigma^2)$, all independent
- Scale both $\sigma = \frac{\ell}{\sqrt{d}}$ and $p = 1 \wedge \frac{\beta}{d}$, but not \mathcal{D}_1
- Accelerate the chain by a factor of d (like before)
- The one-dimensional accelerated processes $Z_j^{(d)}$ weakly converge

The Langevin diffusion with Metropolis jumps

- Z_L and Z_{LM} are Langevin diffusions with speed measure $v(\ell)$ and stationary distributions f and f_1

The Langevin diffusion with Metropolis jumps

- Z_L and Z_{LM} are Langevin diffusions with speed measure $v(\ell)$ and stationary distributions f and f_1
- At random times T_1, T_2, \dots generated by a Poisson process of rate β , generate $Y(T_i) \stackrel{iid}{\sim} \mathcal{D}_1$

The Langevin diffusion with Metropolis jumps

- Z_L and Z_{LM} are Langevin diffusions with speed measure $v(\ell)$ and stationary distributions f and f_1
- At random times T_1, T_2, \dots generated by a Poisson process of rate β , generate $Y(T_i) \stackrel{iid}{\sim} \mathcal{D}_1$
- At time T_i , Z_{LM} jumps by $Y(T_i)$ with probability $\alpha(\ell, Z_{LM}(T_i^-), Z_{LM}(T_i^-) + Y(T_i))$

The Langevin diffusion with Metropolis jumps

- Z_L and Z_{LM} are Langevin diffusions with speed measure $v(\ell)$ and stationary distributions f and f_1
- At random times T_1, T_2, \dots generated by a Poisson process of rate β , generate $Y(T_i) \stackrel{iid}{\sim} \mathcal{D}_1$
- At time T_i , Z_{LM} jumps by $Y(T_i)$ with probability $\alpha(\ell, Z_{LM}(T_i^-), Z_{LM}(T_i^-) + Y(T_i))$
-

$$\alpha(\ell, x, y) = \Phi \left(\frac{\log \frac{f_1(y)}{f_1(x)} - \frac{\ell^2}{2} B}{\ell \sqrt{B}} \right) + \frac{f_1(y)}{f_1(x)} \Phi \left(\frac{-\log \frac{f_1(y)}{f_1(x)} - \frac{\ell^2}{2} B}{\ell \sqrt{B}} \right)$$

The limiting processes

Theorem

If the chain starts at stationnarity ($\mathbf{X}^{(d)}(0) \sim \pi$), then as $d \rightarrow \infty$,

$$Z_1^{(d)} \Rightarrow Z_{LM}, \quad \text{and for } j \geq 2, \quad Z_j^{(d)} \Rightarrow Z_L.$$

Here, \Rightarrow represents *weak convergence in the Skorokhod topology*.

Optimizing the limiting processes

- Run classic RWM on π and tune σ so that acceptance rate is roughly 0.234

Optimizing the limiting processes

- Run classic RWM on π and tune σ so that acceptance rate is roughly 0.234
- Run RWM on f_1 with instrumental distribution \mathcal{D}_1 and acceptance probability $\alpha(\ell, x, y)$ to estimate acceptance probability of large steps, say λ

Optimizing the limiting processes

- Run classic RWM on π and tune σ so that acceptance rate is roughly 0.234
- Run RWM on f_1 with instrumental distribution \mathcal{D}_1 and acceptance probability $\alpha(\ell, x, y)$ to estimate acceptance probability of large steps, say λ
- Choose p , proportion of accepted large steps will be $\approx p\lambda$

Optimizing the limiting processes

- Run classic RWM on π and tune σ so that acceptance rate is roughly 0.234
- Run RWM on f_1 with instrumental distribution \mathcal{D}_1 and acceptance probability $\alpha(\ell, x, y)$ to estimate acceptance probability of large steps, say λ
- Choose p , proportion of accepted large steps will be $\approx p\lambda$
- Number of times we switch modes $\approx np\lambda$

- 1 Random Walk Metropolis algorithm
- 2 Weak convergence of our algorithm
- 3 Summary and Future work

- Can work for much more general target distributions (align modes with rotation)

- Can work for much more general target distributions (align modes with rotation)
- Find objective way to choose p through non-asymptotics

- Can work for much more general target distributions (align modes with rotation)
- Find objective way to choose p through non-asymptotics
- Can replace the large step distribution \mathcal{D} by any *algorithm* ("something-inside-Metropolis")



Guan, Y. and Krone, S. (2007).

Small-world MCMC and convergence to multi-modal distributions:
From slow mixing to fast mixing.

Ann. Appl. Probab., 17:284–304.



Roberts, G., Gelman, A., and Gilks, W. (1997).

Weak convergence and optimal scaling of random walk Metropolis
algorithms.

Ann. Appl. Probab., 7:110–120.