

Classification




- Qualitative variables take values in an unordered set \mathcal{C} , such as:
 $\text{eye color} \in \{\text{brown}, \text{blue}, \text{green}\}$
 $\text{email} \in \{\text{spam}, \text{ham}\}.$
- Given a feature vector X and a qualitative response Y taking values in the set \mathcal{C} , the classification task is to build a function $C(X)$ that takes as input the feature vector X and predicts its value for Y ; i.e. $C(X) \in \mathcal{C}$.
- Often we are more interested in estimating the *probabilities* that X belongs to each category in \mathcal{C} .

Classification

- Qualitative variables take values in an unordered set \mathcal{C} , such as:
 $\text{eye color} \in \{\text{brown}, \text{blue}, \text{green}\}$
 $\text{email} \in \{\text{spam}, \text{ham}\}.$
- Given a feature vector X and a qualitative response Y taking values in the set \mathcal{C} , the classification task is to build a function $C(X)$ that takes as input the feature vector X and predicts its value for Y ; i.e. $C(X) \in \mathcal{C}$.
- Often we are more interested in estimating the *probabilities* that X belongs to each category in \mathcal{C} .

For example, it is more valuable to have an estimate of the probability that an insurance claim is fraudulent, than a classification fraudulent or not.

Example: Credit Card Default



Can we use Linear Regression?

Suppose for the **Default** classification task that we code

$$Y = \begin{cases} 0 & \text{if No} \\ 1 & \text{if Yes.} \end{cases}$$

Can we simply perform a linear regression of Y on X and classify as **Yes** if $\hat{Y} > 0.5$?

Can we use Linear Regression?

Suppose for the **Default** classification task that we code

$$Y = \begin{cases} 0 & \text{if No} \\ 1 & \text{if Yes.} \end{cases}$$

Can we simply perform a linear regression of Y on X and classify as **Yes** if $\hat{Y} > 0.5$?

- In this case of a binary outcome, linear regression does a good job as a classifier, and is equivalent to *linear discriminant analysis* which we discuss later.
- Since in the population $E(Y|X = x) = \Pr(Y = 1|X = x)$, we might think that regression is perfect for this task.

Can we use Linear Regression?



Suppose for the **Default** classification task that we code

$$Y = \begin{cases} 0 & \text{if No} \\ 1 & \text{if Yes.} \end{cases}$$

Can we simply perform a linear regression of Y on X and classify as **Yes** if $\hat{Y} > 0.5$?

- In this case of a binary outcome, linear regression does a good job as a classifier, and is equivalent to *linear discriminant analysis* which we discuss later.
- Since in the population $E(Y|X = x) = \Pr(Y = 1|X = x)$, we might think that regression is perfect for this task.
- However, *linear* regression might produce probabilities less than zero or bigger than one. *Logistic regression* is more appropriate.

Linear versus Logistic Regression



The orange marks indicate the response Y , either 0 or 1. Linear regression does not estimate $\Pr(Y = 1|X)$ well. Logistic regression seems well suited to the task.

Linear Regression continued

Now suppose we have a response variable with three possible values. A patient presents at the emergency room, and we must classify them according to their symptoms.

$$Y = \begin{cases} 1 & \text{if stroke;} \\ 2 & \text{if drug overdose;} \\ 3 & \text{if epileptic seizure.} \end{cases}$$

This coding suggests an ordering, and in fact implies that the difference between **stroke** and **drug overdose** is the same as between **drug overdose** and **epileptic seizure**.

Linear Regression continued

Now suppose we have a response variable with three possible values. A patient presents at the emergency room, and we must classify them according to their symptoms.

$$Y = \begin{cases} 1 & \text{if stroke;} \\ 2 & \text{if drug overdose;} \\ 3 & \text{if epileptic seizure.} \end{cases}$$

This coding suggests an ordering, and in fact implies that the difference between **stroke** and **drug overdose** is the same as between **drug overdose** and **epileptic seizure**.

Linear regression is not appropriate here.

Multiclass Logistic Regression or *Discriminant Analysis* are more appropriate.

Logistic Regression

Let's write $p(X) = \Pr(Y = 1|X)$ for short and consider using **balance** to predict **default**. Logistic regression uses the form

$$p(X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}}.$$

($e \approx 2.71828$ is a mathematical constant [Euler's number.])

It is easy to see that no matter what values β_0 , β_1 or X take, $p(X)$ will have values between 0 and 1.

Logistic Regression

Let's write $p(X) = \Pr(Y = 1|X)$ for short and consider using **balance** to predict **default**. Logistic regression uses the form

$$p(X) = \frac{e^{\beta_0 + \beta_1 X}}{1 + e^{\beta_0 + \beta_1 X}}.$$

($e \approx 2.71828$ is a mathematical constant [Euler's number.])



It is easy to see that no matter what values β_0 , β_1 or X take, $p(X)$ will have values between 0 and 1.

A bit of rearrangement gives

$$\log \left(\frac{p(X)}{1 - p(X)} \right) = \beta_0 + \beta_1 X.$$

This monotone transformation is called the *log odds* or *logit* transformation of $p(X)$. (by log we mean *natural log*: \ln .)

Linear versus Logistic Regression



Logistic regression ensures that our estimate for $p(X)$ lies between 0 and 1.

Maximum Likelihood

We use maximum likelihood to estimate the parameters.

$$\ell(\beta_0, \beta) = \prod_{i:y_i=1} p(x_i) \prod_{i:y_i=0} (1 - p(x_i)).$$

This *likelihood* gives the probability of the observed zeros and ones in the data. We pick β_0 and β_1 to maximize the likelihood of the observed data.

Maximum Likelihood

We use maximum likelihood to estimate the parameters.

$$\ell(\beta_0, \beta) = \prod_{i:y_i=1} p(x_i) \prod_{i:y_i=0} (1 - p(x_i)).$$

This *likelihood* gives the probability of the observed zeros and ones in the data. We pick β_0 and β_1 to maximize the likelihood of the observed data.

Most statistical packages can fit linear logistic regression models by maximum likelihood. In **R** we use the **glm** function.

	Coefficient	Std. Error	Z-statistic	P-value
Intercept	-10.6513	0.3612	-29.5	< 0.0001
balance	0.0055	0.0002	24.9	< 0.0001

Making Predictions

What is our estimated probability of **default** for someone with a balance of \$1000?

$$\hat{p}(X) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 X}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 X}} = \frac{e^{-10.6513 + 0.0055 \times 1000}}{1 + e^{-10.6513 + 0.0055 \times 1000}} = 0.006$$

Making Predictions

What is our estimated probability of **default** for someone with a balance of \$1000?

$$\hat{p}(X) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 X}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 X}} = \frac{e^{-10.6513 + 0.0055 \times 1000}}{1 + e^{-10.6513 + 0.0055 \times 1000}} = 0.006$$

With a balance of \$2000?

$$\hat{p}(X) = \frac{e^{\hat{\beta}_0 + \hat{\beta}_1 X}}{1 + e^{\hat{\beta}_0 + \hat{\beta}_1 X}} = \frac{e^{-10.6513 + 0.0055 \times 2000}}{1 + e^{-10.6513 + 0.0055 \times 2000}} = 0.586$$

Lets do it again, using **student** as the predictor.

	Coefficient	Std. Error	Z-statistic	P-value
Intercept	-3.5041	0.0707	-49.55	< 0.0001
student[Yes]	0.4049	0.1150	3.52	0.0004

Lets do it again, using **student** as the predictor.

	Coefficient	Std. Error	Z-statistic	P-value
Intercept	-3.5041	0.0707	-49.55	< 0.0001
student[Yes]	0.4049	0.1150	3.52	0.0004

$$\widehat{\Pr}(\text{default}=\text{Yes}|\text{student}=\text{Yes}) = \frac{e^{-3.5041+0.4049 \times 1}}{1 + e^{-3.5041+0.4049 \times 1}} = 0.0431,$$

$$\widehat{\Pr}(\text{default}=\text{Yes}|\text{student}=\text{No}) = \frac{e^{-3.5041+0.4049 \times 0}}{1 + e^{-3.5041+0.4049 \times 0}} = 0.0292.$$

Logistic regression with several variables



$$\log \left(\frac{p(X)}{1 - p(X)} \right) = \beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p$$

$$p(X) = \frac{e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}}{1 + e^{\beta_0 + \beta_1 X_1 + \cdots + \beta_p X_p}}$$

	Coefficient	Std. Error	Z-statistic	P-value
Intercept	-10.8690	0.4923	-22.08	< 0.0001
balance	0.0057	0.0002	24.74	< 0.0001
income	0.0030	0.0082	0.37	0.7115
student[Yes]	-0.6468	0.2362	-2.74	0.0062

Why is coefficient for **student** negative, while it was positive before?


Confounding



- Students tend to have higher balances than non-students, so their marginal default rate is higher than for non-students.
- But for each level of balance, students default less than non-students.
- Multiple logistic regression can tease this out.

Example: South African Heart Disease

- 160 cases of MI (myocardial infarction) and 302 controls (all male in age range 15-64), from Western Cape, South Africa in early 80s.
- Overall prevalence very high in this region: 5.1%.
- Measurements on seven predictors (risk factors), shown in scatterplot matrix.
- Goal is to identify relative strengths and directions of risk factors.
- This was part of an intervention study aimed at educating the public on healthier diets.



Scatterplot matrix of the *South African Heart Disease* data. The response is color coded — The cases (MI) are red, the controls turquoise. **famhist** is a binary variable, with 1 indicating family history of MI.

```
> heartfit<-glm(chd~.,data=heart,family=binomial)
> summary(heartfit)
```

Call:

```
glm(formula = chd ~ ., family = binomial, data = heart)
```

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-4.1295997	0.9641558	-4.283	1.84e-05	***
sbp	0.0057607	0.0056326	1.023	0.30643	
tobacco	0.0795256	0.0262150	3.034	0.00242	**
ldl	0.1847793	0.0574115	3.219	0.00129	**
famhistPresent	0.9391855	0.2248691	4.177	2.96e-05	***
obesity	-0.0345434	0.0291053	-1.187	0.23529	
alcohol	0.0006065	0.0044550	0.136	0.89171	
age	0.0425412	0.0101749	4.181	2.90e-05	***

(Dispersion parameter for binomial family taken to be 1)

Null deviance: 596.11 on 461 degrees of freedom
Residual deviance: 483.17 on 454 degrees of freedom
AIC: 499.17


Case-control sampling and logistic regression

- In South African data, there are 160 cases, 302 controls — $\tilde{\pi} = 0.35$ are cases. Yet the prevalence of MI in this region is $\pi = 0.05$.
- With case-control samples, we can estimate the regression parameters β_j accurately (if our model is correct); the constant term β_0 is incorrect.
- We can correct the estimated intercept by a simple transformation

$$\hat{\beta}_0^* = \hat{\beta}_0 + \log \frac{\pi}{1 - \pi} - \log \frac{\tilde{\pi}}{1 - \tilde{\pi}}$$

- Often cases are rare and we take them all; up to five times that number of controls is sufficient. See next frame

Diminishing returns in unbalanced binary data



Sampling more controls than cases reduces the variance of the parameter estimates. But after a ratio of about 5 to 1 the variance reduction flattens out.

Logistic regression with more than two classes

So far we have discussed logistic regression with two classes. It is easily generalized to more than two classes. One version (used in the R package **glmnet**) has the symmetric form

$$\Pr(Y = k|X) = \frac{e^{\beta_{0k} + \beta_{1k}X_1 + \dots + \beta_{pk}X_p}}{\sum_{\ell=1}^K e^{\beta_{0\ell} + \beta_{1\ell}X_1 + \dots + \beta_{p\ell}X_p}}$$

Here there is a linear function for *each* class.

Logistic regression with more than two classes

So far we have discussed logistic regression with two classes. It is easily generalized to more than two classes. One version (used in the R package **glmnet**) has the symmetric form

$$\Pr(Y = k|X) = \frac{e^{\beta_{0k} + \beta_{1k}X_1 + \dots + \beta_{pk}X_p}}{\sum_{\ell=1}^K e^{\beta_{0\ell} + \beta_{1\ell}X_1 + \dots + \beta_{p\ell}X_p}}$$

Here there is a linear function for *each* class.

(The *mathier* students will recognize that some cancellation is possible, and only $K - 1$ linear functions are needed as in 2-class logistic regression.)

Logistic regression with more than two classes

So far we have discussed logistic regression with two classes. It is easily generalized to more than two classes. One version (used in the R package `glmnet`) has the symmetric form

$$\Pr(Y = k|X) = \frac{e^{\beta_{0k} + \beta_{1k}X_1 + \dots + \beta_{pk}X_p}}{\sum_{\ell=1}^K e^{\beta_{0\ell} + \beta_{1\ell}X_1 + \dots + \beta_{p\ell}X_p}}$$

Here there is a linear function for *each* class.

(The *mathier* students will recognize that some cancellation is possible, and only $K - 1$ linear functions are needed as in 2-class logistic regression.)

Multiclass logistic regression is also referred to as *multinomial regression*.