

MATH 405: Assignment 4

Micah Sherry

March 7, 2025

1. Find the smallest positive solution to the system of congruences.

$$x \equiv 4 \pmod{7}$$

$$x \equiv 5 \pmod{11}$$

$$x \equiv 2 \pmod{16}$$

$$x \equiv 1 \pmod{19}$$

$$M = 7 \cdot 11 \cdot 16 \cdot 19 = 23408$$

$$M_1 = 11 \cdot 16 \cdot 19 = 3344$$

$$M_2 = 7 \cdot 16 \cdot 19 = 2128$$

$$M_3 = 7 \cdot 11 \cdot 19 = 1463$$

$$M_4 = 7 \cdot 11 \cdot 16 = 1232$$

$$M_1 b_1 \equiv 1 \pmod{7}$$

$$5b_1 \equiv 1 \pmod{7}$$

$$1 = 3(5) - 2(7)$$

$$\text{Therefore } b_1 \equiv 3 \pmod{7}$$

$$M_3 b_3 \equiv 1 \pmod{16}$$

$$7b_3 \equiv 1 \pmod{16}$$

$$1 = 7(7) - 3(16)$$

$$\text{Therefore } b_3 \equiv 7 \pmod{16}$$

$$M_2 b_2 \equiv 1 \pmod{11}$$

$$5b_2 \equiv 1 \pmod{11}$$

$$1 = -2(5) - (11)$$

$$\text{Therefore } b_2 \equiv -2 \equiv 9 \pmod{11}$$

$$M_4 b_4 \equiv 1 \pmod{19}$$

$$16b_4 \equiv 1 \pmod{19}$$

$$1 = 6(16) - 5(19)$$

$$\text{Therefore } b_4 \equiv 6 \pmod{19}$$

$$\begin{aligned} x_0 &= 4M_1 b_1 + 5M_2 b_2 + 2M_3 b_3 + 1M_4 b_4 \\ &= 4(3344)(3) + 5(2128)(9) + 2(1463)(7) + 1(1232)(6) \\ &= 163762 \end{aligned}$$

$$x_k = x_0 + kM \text{ (where } k \text{ is some integer)}$$

$$x_{-6} = 23314 \text{ is the smallest positive value for } x.$$

2. Consider the set $\mathbb{Z}_3[i] = \{a + bi \mid a, b \in \mathbb{Z}_3\}$ where $i = \sqrt{-1}$.

- (a) find all the elements of $\mathbb{Z}_3[i]$. How many are there?
 Notice there are 3 choices for a and 3 for b, so $\mathbb{Z}_3[i]$ has 9 elements.
 $0, 1, 2, i, 1+i, 2+i, 2i, 1+2i, 2+2i$
- (b) $1+2i \in \mathbb{Z}_3[i]$ has a multiplicative inverse in find it.

$$\begin{aligned}
 (1+2i)(2+2i) &= \\
 &= 2 + 4i + 2i + 4i^2 \\
 &= 2 - 4 \\
 &= -2 \\
 &= 1
 \end{aligned}$$

Therefore $(2+2i)$ is the multiplicative inverse of $(1+2i)$.

- (c) Classify each nonzero element of $\mathbb{Z}_3[i]$ as a unit, a zero divisor or neither.

Number	Inverse	classification
0	NA	Neither
1	1	unit
2	2	unit
i	2i	unit
1+i	2+i	unit
2+i	1+i	unit
2i	i	unit
1+2i	2i+2	unit
2+2i	1+2i	unit

3. let R be a ring and let S and T be subrings of R . Let $M = S \cap T$. Show that M is a subring of R

Proof. Let S and T be subrings of a Ring R . And Let $M = S \cap T$.

To show M is a subring we need to show that $0, 1 \in M$, M is closed under addition and multiplication, and if $a \in M$ then $-a \in M$

Since S is a subring $0 \in S$. Similarly, T is a subring $0 \in T$. Therefore $0 \in M$.

Since S is a subring $1 \in S$. Similarly, T is a subring $1 \in T$. Therefore $1 \in M$.

Let $a, b \in M$. Since, $a, b \in S$ since S is a subring $a+b \in S$

Similarly, $a, b \in T$ since T is a subring $a+b \in T$

Therefore $a+b \in M$ Thus M is closed under addition.

Let $a, b \in M$. Since, $a, b \in S$ since S is a subring $a \cdot b \in S$

Similarly, $a, b \in T$ since T is a subring $a \cdot b \in T$

Therefore $a \cdot b \in M$ Thus M is closed under multiplication.

Let $a \in M$. Since, $a \in S$ since S is a subring $-a \in S$

Similarly, $a \in T$ since T is a subring $-a \in T$

Therefore if $a \in M$ then $-a \in M$.

Therefore M is a Subring of R

□

4. Let R be a ring (not necessarily commutative).

Let $a, b \in R$, prove that $(a \cdot b)^{-1} = b^{-1} \cdot a^{-1}$

Proof. Let R be a ring and let $a, b \in R$. And assuming a^{-1}, b^{-1} exist. Consider:

$$\begin{aligned}
 (a \cdot b) \cdot (b^{-1} \cdot a^{-1}) &= a \cdot (b \cdot b^{-1}) \cdot a^{-1} && \text{(by associativity of } R) \\
 &= a \cdot 1 \cdot a^{-1} && \text{(by definition of multiplicative inverse)} \\
 &= a \cdot a^{-1} && \text{(by definition of multiplicative identity)} \\
 &= 1 && \text{(by definition of multiplicative inverse)}
 \end{aligned}$$

Therefore $(b^{-1} \cdot a^{-1})$ is the multiplicative inverse of $(a \cdot b)$

□

Extra Credit: let R be a ring (not necessarily commutative).

If for any $a, b \in R$, $(a \cdot b)^{-1} = a^{-1} \cdot b^{-1}$ then show that R is commutative.

Proof. Let $a, b \in R$. Assume a^{-1}, b^{-1} exist and $(a \cdot b)^{-1} = (a^{-1} \cdot b^{-1})$. from proof of problem 4 we have $(a \cdot b)^{-1} = (b^{-1} \cdot a^{-1})$. So,

$$\begin{aligned}(a^{-1} \cdot b^{-1}) &= (b^{-1} \cdot a^{-1}) \\ b^{-1} &= a \cdot (b^{-1} \cdot a^{-1}) && \text{(left multiply by } a) \\ 1 &= b \cdot a \cdot (b^{-1} \cdot a^{-1}) && \text{(left multiply by } b) \\ a &= b \cdot a \cdot b^{-1} && \text{(right multiply by } a) \\ a \cdot b &= b \cdot a && \text{(right multiply by } b)\end{aligned}$$

Therefore since $b \cdot a = a \cdot b$, R is commutative.

□