# Factorization of Polynomials Over the Boolean Semifield

MAA Student Talk

Micah Sherry
MicahJSherry@gmail.com

March 27, 2026

# Background

The Boolean semifield, denoted as $\mathbb{B}$, is an algebraic structure with two operations: logical OR, analogous to addition, and logical AND, analogous to multiplication. $\mathbb{B}$ the only finite semifield that is not a field. Our research explores the factorization of polynomials over the Boolean semifield, denoted as $\mathbb{B}[x]$.

A key challenge in $\mathbb{B}[x]$ is that factorization is not unique. Specifically, we investigate the factorization of a particular class of polynomials in $\mathbb{B}[x]$ namely:

$$f_n(x) = \sum_{i=0}^{n} x^i = x^n + x^{n-1} + \cdots + x + 1$$

Understanding the non-unique factorization of these polynomials is a central focus of our study. This research contributes to a deeper understanding of algebraic structures beyond traditional fields and rings.

# Definitions

- A semifield is an algebraic structure like a field with addition and multiplication. That does not necessarily have additive inverses and does have multiplicative inverses.
- Let $\mathbb{B} = \{0, 1\}$ be the Boolean semifield equipped with logical OR as addition and Logical AND as Multiplication.
- Let $\mathbb{B}[x]$ be the set of all polynomials in $x$ over $\mathbb{B}$.
- Let $f_n(x) \in \mathbb{B}[x]$, be defined as $f_n(x) = x^n + x^{n-1} + \cdots + x + 1$.

# Necessary but Not Sufficient Condition for Factorization

## Theorem

*Let* $f(x) = \sum_{i=0}^{n} f_i x^i$ *with* $f_0 = 1$. *If* $f(x)$ *is factorable, then there exist* $j, k$ *with* $1 \leq j, k \leq n-1$ *and* $j + k = n$ *such that* $f_j = f_k = 1$.

## Proof.

Assume $f(x) = g(x)h(x)$ with $1 \leq \deg(g), \deg(h) \leq n-1$. Let $\deg(g) = j$, $\deg(h) = k$. Then $g(x) = x^j + g'(x) + 1, \quad h(x) = x^k + h'(x) + 1$.

$$\begin{aligned} f(x) &= g(x)h(x) \\ &= (x^j + g'(x) + 1)(x^k + h'(x) + 1) \\ &= x^{j+k} + x^j + x^k + \big(x^j h'(x) + x^k g'(x) + g'(x)h'(x) + g'(x) + h'(x)\big) + 1. \end{aligned}$$

Thus $j + k = n$ and $f_j = f_k = 1$ □

**Remarks.**

$f(x) = x^6 + x^4 + x^2 + x + 1$ satisfies the condition but is not factorable.

$f(x) = x^6 + x^5 + x^2 + 1$ does not satisfy the condition and is not factorable.

# Examples of Possible Factorization

| n | $f_n(x)$ | Factorizations |
|---|---|---|
| 2 | $x^2 + x + 1$ | $(x+1)^2$ |
| 3 | $x^3 + x^2 + x + 1$ | $(x+1)^3$<br>$(x+1)(x^2+1)$ |
| 4 | $x^4 + x^3 + x^2 + x + 1$ | $(x+1)^4$<br>$(x+1)^2(x^2+1)$<br>$(x+1)(x^3+x^2+1)$<br>$(x+1)(x^3+x+1)$ |
| 5 | $x^5 + x^4 + x^3 + x^2 + x + 1$ | $(x+1)^5$<br>$(x+1)^3(x^2+1)$<br>$(x+1)^2(x^3+x^2+1)$<br>$(x+1)^2(x^3+x+1)$<br>$(x+1)^2(x^3+1)$<br>$(x+1)(x^4+x^3+x^2+1)$<br>$(x+1)(x^4+x^2+x+1)$<br>$(x+1)(x^2+1)^2$ |

# Exponential Growth of Factors



# Growth Rate of Factors



# Divisibility of $f_n(x)$

## Lemma

*Lemma 3.1 from the paper 'On the Sequence A079500 and Its Combinatorial Interpretations' [3] states conditions for divisibility in the numbral arithmetic: For each $n, d > 0$, it holds that $[d]$ is a divisor of $[2^n - 1]$ if and only if it ends with the digit 1, and it does not contain any subsequence of 0s having length greater than $n - k$, with $k$ being the length of $[d]$.*

## Theorem

*Let $g(x) \in \mathbb{B}[x]$ such that the constant term of $g(x)$ is 1. If $\deg(g(x)) = k$ and $2k - 1 \leq n$, then $g(x)$ divides $f_n(x)$.*

## Proof.

Let $\theta(h(x))$ be the isomorphism between $\mathbb{B}[x]$ and the numbral arithmetic $\theta(h(x)) = [h(2)]$. Assume $2k - 1 \leq n$ and let $d = \theta(g(x))$ (notice $[d]$ is length $k + 1$). Let $m$ be the maximum length of zeros in $[d]$. Notice $m \leq k - 1$. Notice $\theta(f_n(x)) = 2^{n+1} - 1$. By the above lemma, $d$ divides $2^{n+1} - 1$ because $k - 1 \leq n + 1 - (k + 1)$, which is equivalent to $2k - 1 \leq n$. And by applying $\theta^{-1}$ we see $g(x)$ divides $f_n(x)$. $\square$

# Factorizations Of The Form $(x+1)h(x) = f_n(x)$

## Theorem

$(x+1)h(x) = f_n(x)$ *if and only if* $h(x) = \sum_{i=0}^{n-1} h_i x^i$ *With* $h_{n-1} = h_0 = 1$ *and* $h_i + h_{i-1} = 1$ *for* $0 < i < n-1$.

## Proof.

($\rightarrow$): Assume that $(x+1)h(x) = f_n(x)$ and let $h(x) = \sum_{i=0}^{n-1} h_i x^i$. (Notice $h_{n-1} = h_0 = 1$ otherwise it would contradict the assumption). Now assume for the sake of contradiction that there exists $k < n-1$ such that $h_k = h_{k-1} = 0$. Consider the following:

$$(x+1)h(x) = x\sum_{i=0}^{n-1} h_i x^i + \sum_{i=0}^{n-1} h_i x^i.$$

Notice the $k$ term will have coefficient $h_k + h_{k-1} = 0$, contradicting the original assumption.
($\leftarrow$): Assume $h(x) = \sum_{i=0}^{n-1} a_i x^i$ with $h_{n-1} = h_0 = 1$ and for all $h_i$, $h_i + h_{i-1} = 1$. Consider the following:

$$(x+1)h(x) = \sum_{i=0}^{n-1} a_i x^{i+1} + \sum_{i=0}^{n-1} a_i x^i = a_{n-1}x^n + \sum_{i=1}^{n-1}(a_{i-1} + a_i)x^i + a_0 = f_n(x).$$

Therefore, the theorem holds. $\square$

# Factorizations Of The Form $(x+1)h(x) = f_n(x)(Cont.)$

## Theorem

*Let* $G_{n\,1} = \{h(x) | (x+1)h(x) = f_n(x)\}$ *Let* $F_n$ *be the nth Fibonacci number.* $|G_{n\,1}| = F_n$ *for* $n \geq 1$

## Proof (by Strong induction).

$G_{1\,1} = \{1\}, \quad |G_{1\,1}| = 1 = F_1 \quad \& \quad G_{2\,1} = \{x+1\}, \quad |G_{2\,1}| = 1 = F_2$
Therefore the base cases hold. Assume that $|G_{j\,1}| = F_j$ for all $j \in \{1, 2, \ldots, k\}$ for some $k$.
Let $h(x) \in G_{(k+1)\,1}$. Notice $\deg(h(x)) = k$ and $h_k = 1$ consider the possible values for $h_{k-1}$.

1. Case 1. $h_{k-1} = 1$. In which case $h(x) = x^k + h'(x)$ Where $h'(x) \in G_{k\,1}$.
2. Case 2. $h_{k-1} = 0$, So $h_{k-2} = 1$. In which case $h(x) = x^k + 0x^{k-1} + h''(x)$ Where $h''(x) \in G_{(k-1)\,1}$.

So, $|G_{(k+1)\,1}| = |G_{k\,1}| + |G_{(k-1)\,1}| = F_k + F_{k-1} = F_{k+1}$. Therefore, by the Principle of Mathematical Induction $|G_{n\,1}| = F_n$ for all $n \geq 1$. □

# Factorizations Of The Form $(x^k+1)g(x) = f_n(x)$

## Theorem

*Let* $G_{n\,k} = \{g(x) | (x^k+1)g(x) = f_n(x)\}$, *let* $m = n - 3k + 1$, *let* $q = \lfloor \frac{m}{k} \rfloor$, *and let* $r = m \bmod k$.

$$|G_{n\,k}| = (F_{q+3})^r \times (F_{q+2})^{k-r}$$

## Sketch.

Consider $g(x) = \sum_{i=0}^{n-k} g_i x^i \in G_{nk}$.

$$\begin{aligned}(x^k+1)g(x) &= \sum_{i=0}^{n-k} g_i x^{i+k} + \sum_{i=0}^{n-k} g_i x^i \\ &= \sum_{i=k}^{n} g_{i-k} x^i + \sum_{i=0}^{n-k} g_i x^i \\ &= \sum_{i=n-k+1}^{n} g_i x^i + \sum_{i=k}^{n-k} (g_k + g_{i-k}) x^i + \sum_{i=0}^{k-1} g_i x^i = f_n(x)\end{aligned}$$

Notice this implies $g_i = 1$ for $i \leq k-1$ or $i \geq n-2k+1$ and $g_i + g_{i-k} = 1$ for $k \leq i \leq n-2k$

# Factorizations Of The Form $(x^k + 1)g(x) = f_n(x)$ (Cont.)

## Sketch.

Construct a partition the coefficients $h_i$ where $k \leq i \leq n - 2k$. Let $g_i \in P_j$ if $i \equiv j \mod k$.

| | | |
|---|---|---|
| $P_0$ | $g_k,\ g_{2k},\ \cdots$ | $i \equiv 0 \mod k$ |
| $P_1$ | $g_{k+1},\ g_{2k+1}, \cdots$ | $i \equiv 1 \mod k$ |
| $\vdots$ | $\vdots$ | $\vdots$ |
| $P_0$ | $g_{2k-1},\ g_{3k-1}, \cdots$ | $i \equiv k-1 \mod k$ |

Let $m = n - 3k + 1$, $q = \lfloor \frac{m}{k} \rfloor$, and $r = m \mod k$. The length of $r$ of these partitions is $q + 1$ and $k - r$ have length $q$. It can be shown that the number of options (where $g_i + g_{i-k} = 1$) for $P_j$ of length $q$ is $F_{q+2}$. So, The number of valid polynomials is $(F_{q+3})^r \times (F_{q+2})^{k-r}$ $\square$

# Sources


[1] Francisco Alarcon and Daniel D Anderson.
Commutative semirings and their lattices of ideals.
*Houston J. Math*, 20(4):571–590, 1994.

[2] David Applegate, Marc LeBrun, and Neil JA Sloane.
Dismal arithmetic.
*arXiv preprint arXiv:1107.1130*, 2011.

[3] Andrea Frosini, S Rinaldi, and Pian dei Mantellini.
On the sequence a079500 and its combinatorial interpretations.
*J. Integer Seq*, 9, 2006.

[4] Gal Gross.
*Maximally additively reducible subsets of the integers.*
University of Toronto (Canada), 2019.

[5] Ki Hang Kim and Fred W Roush.
Factorization of polynomials in one variable over the tropical semiring.
*arXiv preprint math/0501167*, 2005.

[6] OEIS Foundation Inc.
A067139: Irreducible elements in or-numbral arithmetic.
The On-Line Encyclopedia of Integer Sequences, 2002.
Available at `https://oeis.org/A067139`.

[7] OEIS Foundation Inc.
A169912: Number of irreducible boolean polynomials of degree n.
The On-Line Encyclopedia of Integer Sequences, 2010.
Available at `https://oeis.org/A169912`.

[8] OEIS Foundation Inc.
A171000: Irreducible boolean polynomials written as binary vectors.
The On-Line Encyclopedia of Integer Sequences, 2010.
Available at `https://oeis.org/A171000`.

# Related Topics

1. Set Addition
   $A + B = \{a + b | a \in A \text{ and } b \in B\}$ where $A, B \subseteq \mathbb{N}$
2. Dismal Arithmetic (base $b \geq 2$) Let $a$ and $b$ be base-$b$ numbers with digits in $\{0, 1, \ldots, b-1\}$.
   1. $a \oplus b$ is given by digitwise max (no carries).
   2. $a \otimes b$ is given by digitwise min, combined as in ordinary multiplication (no carries).
3. Numbral Arithmetic let $a$ and $b$ be binary sequences.
   1. $a + b$ is given by Bitwise OR.
   2. $a \times b$ is given by Shift-and-OR.
4. Polynomials in $x$ over the Tropical semiring