

PATIENT

Age (years)

45

Sex

Male

Diabetes duration (y)

20

Age at onset (derived)

25

Ethnicity (simplified PCE/QRISK3/UKPDS mapping)

White

MEASUREMENTS

HbA1c (%)

8.0

HbA1c (mmol/mol)

63,93

Systolic BP (mmHg)

130

Diastolic BP (mmHg)

78

Total chol (mmol/L)

4,8

HDL (mmol/L)

1,4

TG (mmol/L)

1,2

Measured LDL (mmol/L, optional; required if TG ≥4.5)

Otherwise estimated from lipids

eGFR (mL/min/1.73m²)

95

Albuminuria

Normal

BMI (kg/m²)

25

☐ Current smoker

☒ Regular physical activity (≥30 min/day, Steno)

☐ On BP-lowering treatment

☐ On a statin

☐ Atrial fibrillation

☐ Retinopathy

☐ Prior CVD

☐ Family history of premature CVD

CALIBRATION

SCORE2 risk region (drives SCORE2 family)

High (Central Europe, incl. Czechia)

Exploratory T1D cumulative-hazard multiplier (default 1)


1,0

Derived: LDL (estimated) 2.9 mmol/L · non-HDL 3.4 mmol/L · TC:HDL 3.43 (unitless) · onset age 25.00


CALCULATOR OUTPUTS — MODEL-SPECIFIC ENDPOINTS

Outputs describe different endpoints. Common analytic bands are not treatment thresholds.


TYPE 1 DIABETES–SPECIFIC — 10-YEAR ENDPOINTS




TYPE 1 DIABETES–SPECIFIC — NATIVE FIVE-YEAR ENDPOINT



TYPE-2-DERIVED (OFF-LABEL IN T1D)



GENERAL / MIXED-POPULATION MODELS



T1D-specific

Type-2-derived

General-population

 · vertical markers appear only on 10-year/adapted outputs and denote common analytic boundaries. Bars stop at 40%; numerical labels show the full calculated risk.

► [Methods, implementation checks & coefficients](#)