1A.

 $5_{10} = 00101_2$ 

-10<sub>10</sub> in signed 5-bit binary:  $10_{10} = 01010_2$ , invert it then +1<sub>2</sub>  $\rightarrow$  -10<sub>10</sub> = 10101<sub>2</sub> + 1<sub>2</sub> = 10110<sub>2</sub>

1B.

5 \* -10 = -50, or 5 \* 10 = 50, then 50 \* -1 = -50.

5 = 00101, 10 = 01010.

01010	01010	01010	01010	01010	01010	
00101	00010 1	00001 01	00000 101	00000 0101	00000 00101	
01010	00000 0	01010 00	00000 000	00000 0000	00000 00000	

0000010000
01010
000000
0101000
00000000
000000000
000000000
0000110010

Find two's compliment: invert the result:  $0000110010_2 \rightarrow 1111001101_2$ , +  $1_2 = 1111001110_2$ .

## 1C-1. Zero extension from 5-bit to 8-bit:

 $5_{10} = 00101_2$ . Add zeros in the front of MSB until it is 8-bit:  $00101_2 \rightarrow 00000101_2$ .

 $-10_{10} = 10110_2$ . Add zeros in the front of MSB until it is 8-bit:  $10110_2 \rightarrow 00010110_2$ .

## 1C-2. Signed extension from 5-bit to 8-bit:

 $5_{10} = 00101_2$ . Number is positive, add zeros in the front of MSB until it is 8-bit:

 $00101_2 \rightarrow 00000101_2$ .

 $-10_{10}$  =  $10110_2$ . Number is negative, add ones in the front of MSB until it is 8-bit:

 $10110_2 \rightarrow 11110110_2$ .

2A.

F(x, y, z) = x + y + z'

X	Υ	Z	F(x, y, z)
0	0	0	1
0	0	1	0
0	1	0	1
0	1	1	1
1	0	0	1
1	0	1	1
1	1	0	1
1	1	1	1

### 2B.

F(x, y)	, z)	= x'y'	+	yΖ
---------	------	--------	---	----

X	Υ	Z	F(x, y, z)
0	0	0	1
0	0	1	1
0	1	0	0
0	1	1	1
1	0	0	0
1	0	1	0
1	1	0	0
1	1	1	1

#### 3A.

$$x'z + y + xy' = x + y + z$$

$$x'z + (x + y)(y + y') = x + y + z$$

$$x'z + (x + y)(1) = x + y + z$$

$$x'z + (x + y) = x + y + z$$

$$x'z + x + y = x + y + z$$

$$(x + x')(x + z) + y = x + y + z$$

$$(1)(x + z) + y = x + y + z$$

$$x + y + z = x + y + z$$

#### 3B.

$$abc' + bc'd' + bc + c'd = b + c'd$$

$$abc' + (bc'd' + c'd) + bc = b + c'd$$

$$abc' + (c'(bd' + d)) + bc = b + c'd$$

$$abc' + (c'((b + d)(d' + d))) + bc = b + c'd$$

$$abc' + (c'((b + d)(1))) + bc = b + c'd$$

$$abc' + (c'(b + d)) + bc = b + c'd$$

$$abc' + bc' + c'd + bc = b + c'd$$

$$b(ac' + c' + c) + c'd = b + c'd$$

$$b(ac' + (c' + c)) + c'd = b + c'd$$

$$b(ac' + (1)) + c'd = b + c'd$$

$$b(1) + c'd = b + c'd$$

$$b + c'd = b + c'd$$

4A.  $f(A, B, C, D) = \sum m(0, 1, 5, 7, 8, 10, 14, 15)$ 

cd				
ab	00	01	11	10
00	1)		3	
01	U	[1]	1	6
11	8		1/	
10	(1)			1/

0000 0001 0111 1110 1000 0101 1111 1010

B'C'D' + A'C'D + BCD + ACD' = f(A, B, C, D)

<u>Prime implicants</u>: **B'C'D'**, **A'C'D**, **BCD**, **ACD'**, all groups listed are in their maximum groupings. <u>Essential prime implicants</u>: There are no essential prime implicants, because there are no 1s in the vertical groupings not covered by the horizontal groupings.

4B.  $f(W, X, Y, Z) = \sum m(2, 4, 9, 12, 15) + d(3, 5, 6, 13)$ 

	YZ				
WX		00	01	11	10
	00			X	1
	01	1)	X	) (	X
	11	1	M	1)	(
	10		V		
0100	1101	1 1	101	001	0
1100	1001	1 1:	111	001	1

0101

1101

XY' + WY'Z + WXZ + W'X'Y = f(W, X, Y, Z)

<u>Prime implicants</u>: **XY', WY'Z, WXZ, W'X'Y**, because those are the maximum groupings available. A group of 4 followed by 3 groups of 2 are the maximum groupings available with adding addition terms (the red groups) to the answer.


Essential prime implicants: XY', WY'Z, WXZ, W'X'Y are all essential prime implicants. 1100 in XY' expressed in any other groupings without decreasing the grouping's size, so it is an essential prime implicant. 1001 in WY'Z cannot be grouped by any other way, so that is an EPI. 1111 in WXZ cannot be grouped in any other way, it is also an EPI. 0010 in W'X'Y' can be grouped with 0110 to form another group. You can select either group to be the essential prime implicant, but doing so will make the other group non-essential, so I chose to group it with 0011 to make W'X'Y' an EPI. Additionally, 0110 and 0011 are don't cares, so if you choose to group with either of them, you can treat the non-grouped one as 0, eliminating the need to group it.

5. Take a 4-bit number, return true if number is between (inclusive) 5 and 12, and if it's even.

# Truth table:

Decimal	W	Х	Υ	Z	f(W, X, Y, Z)
0	0	0	0	0	0
1	0	0	0	1	0
2	0	0	1	0	0
3	0	0	1	1	0
4	0	1	0	0	0
5	0	1	0	1	0
6	0	1	1	0	1
7	0	1	1	1	0
8	1	0	0	0	1
9	1	0	0	1	0
10	1	0	1	0	1
11	1	0	1	1	0
12	1	1	0	0	1
13	1	1	0	1	0
14	1	1	1	0	0
15	1	1	1	1	0

# K-Map:



1100 0110 1000 1000 1010

WY'Z'+W'XYZ'+WX'Z' = f(W, X, Y, Z)

Schematic diagram:

