

Automated Faunal Activity Monitoring Report

BNG-aligned supporting evidence — not a statutory Biodiversity Gain Plan

This report adopts the structure of a Biodiversity Gain Plan to present species-monitoring evidence. It is not a statutory Biodiversity Gain Plan and does not calculate biodiversity units. Camera-trap detections are presented as evidence of faunal activity only.

Document control

Field	Value
Site name	Not recorded
Monitoring point / camera ID	Not recorded
OS grid reference	Not recorded — required
Monitoring period	2026-06-28 to 2026-07-16 (local dates, Europe/London)
Report version	2.0
Date generated	2026-09-17
Prepared by	Not provided
Reviewed by	Not provided
Data source	Dedicated recipient mailbox — TrapTracker RT alert emails
Upstream classifier / model version	Not recorded — not transmitted in the alert
BioCLIP model version	Not recorded

Executive summary

Between 2026-06-28 and 2026-07-16, one fixed feeder-facing monitoring point recorded 475 alert events across 4 automated species classifications. The system was confirmed operational for 434.9 hours (100% of covered time). The most frequent automated classification was *Columba palumbus* (96.4% of events), with 3 less-frequent labels recorded less often. BioCLIP independently corroborated only 3% of classifications (460 of 475 not corroborated), and no competent ecologist has verified the identifications, so the species-level results are provisional. This report is a monitoring baseline for future comparison; it does not measure abundance, habitat condition, or site-wide biodiversity, and it calculates no biodiversity units. It is not a statutory Biodiversity Gain Plan.

Headline summary

TOTAL ALERT EVENTS 475 raw count, not animals	PROVISIONAL SPECIES LABELS 4 automated, unverified	CONFIRMED OPERATIONAL HOURS 434.9 over known status only	OPERATIONAL COVERAGE 100% of covered time
BIOCLIP NOT CORROBORATED 97% 460 of 475			

Evidence quality

BioCLIP did not independently corroborate 460 of 475 upstream classifications

(~97%). Lack of corroboration does not necessarily indicate an incorrect classification, but the species-level results remain provisional until image review.

Evidence-quality measure	Value
Total upstream classifications	475
BioCLIP exact-label corroborated	15
Not corroborated	460
Exact-label agreement rate	3%
Exact-label non-agreement rate	97%
Classifications human-reviewed	Not available
Overall status	Provisional

Corroboration method: BioCLIP's single best (top-1) label is resolved to a scientific binomial and compared for exact equality with the upstream label's binomial (both mapped through the shared alias table, which absorbs known synonyms); a match is 'corroborated', anything else is 'not corroborated'. The comparison is label-only — the 0.5 low-confidence flag does not enter it — and BioCLIP is an automated model, not human or ecological verification (see Appendix B1).

1. Submission and site identity

- **Report date (generated):** 2026-09-17
- **Monitoring window:** 2026-06-28 to 2026-07-16 (local dates, Europe/London)
- **Site / monitoring point:** a single fixed, feeder-facing trail camera at one location. The detection data carries no camera identity or coordinates (none is transmitted with the alerts), so the site is described from the known deployment, not asserted from the data.
- **OS grid reference (camera location): NOT RECORDED — required.** No surveyed grid reference is present in the detection data; a real submission requires one, and none is fabricated here.
- **Monitoring-point ID:** Not recorded. **Camera orientation:** Not recorded. **Deployment start / end:** Not recorded. (None of these is transmitted with the alert; none is inferred.)
- **Feeder-facing status:** yes — the camera faces a feeder (known deployment context, not asserted from the data).
- **Site / camera-position map:** Not available — no map or camera-position diagram is available to the report generator.
- **Monitoring area — what it covers and does not:** the field of view of one fixed camera pointed at a feeder. It records what passes that single point within the camera's frame and trigger. It does **not** cover the wider site, other habitat features, or anything outside the camera's field of view; its temporal coverage is bounded (see §5).

2. Provenance and competence

Unlike a statutory Biodiversity Gain Plan, this report does not include certification by a competent ecologist. Instead, this section explains which aspects of the report were generated automatically and which require human verification.

- **No competent ecologist has verified the species identifications in this report.** Every identification is a machine output routed to a human for review, not certified by one.

- **Identifications are automated:** the upstream classifier assigns the species label, and an independent BioCLIP taxonomic model provides a cross-check. Both are model outputs.
- **Cross-check outcome (derived from this window):** BioCLIP did **not corroborate 460 of 475 (97%)** of the upstream labels (see the Evidence-quality panel). A label that was not corroborated is not necessarily incorrect, but it is not an independently corroborated one either.
- **Non-target identifications are provisional:** identifications of less-frequent (non-dominant) species are especially uncertain and must be checked against the actual images before any use.

3. Baseline status

- This monitoring window **establishes a baseline**. The relevant date is the window start (**2026-06-28**); the first record in the window forms the initial baseline observation.
- **No prior baseline exists** to measure against. This report therefore states current activity-monitoring evidence only.
- **No change is computed** — no trend is presented as habitat change, and no difference of any kind is calculated. A later monitoring window could be compared against this baseline, but that comparison is not made here.

4. Monitoring period and survey effort

The monitoring period and effort behind the counts — the same figures as the standard monitoring report.

- **Requested window:** 2026-06-28 to 2026-07-16 (local dates, Europe/London) (**456h**).
- **Coverage (first-to-last detection span):** 2026-06-28 09:24 → 2026-07-16 12:15 UTC — a span of **434.9h** (not the calendar range those dates touch).
- **Proven system downtime within coverage: 0.0h.**
- **Confirmed operational: 434.9 of 434.9 hours (100%).**

Counts elsewhere in this report are raw over this effort, not divided by operational hours; with one monitoring point and no time-of-day control an events-per-hour rate would not be sound as an abundance measure.

Effort-standardised indices (*indices of recorded camera activity, not abundance — see the calculation definitions in Appendix B1*):

Index	Value
Alert events per 100 confirmed operational hours	109.2
Active days per 10 calendar days in the coverage span	9.5
Confirmed operational hours	434.9
Confirmed downtime hours	0.0
Operational coverage	100%

5. Survey constraints

The survey constraints on this monitoring evidence — the appropriate place to record them, gathered once, here.

- **Single camera, single location, feeder-facing** — detections are what passed one fixed point, not a site-level or transect survey.
- **Automated identifications** — unverified; see §2.
- **Monitoring outages** — no delivery silence in this window was evidenced as an outage; any quiet stretch is undetermined (a real lull and undetected downtime cannot be told apart).

- **Short window** — a 19-day window in summer; not a full-year or multi-season survey.
- **Time-of-day coverage** — detections span 05:00–23:00 local; no detections fall outside that span in this window, so hours beyond it are unrepresented, not evidenced as inactive.

6. Automated species classification results

These are **automated, provisional** classifications, not confirmed presences. **No detection count is a biodiversity value**, and counts are alert EVENTS (one alert-rule × image match) — not animals, not abundance, and not a community or biodiversity assessment. Scientific names are the model's labels pending image review.


Automated classification labels (N = 475 alert events):

Species label	Alert events	Share	Active days	BioCLIP corroborated	Not corroborated	Human-reviewed	Final status
<i>Columba palumbus</i>	458	96.4%	18	15	443	Not available	Provisional
<i>Corvus corone</i>	8	1.7%	4	0	8	Not available	Provisional
<i>Sciurus carolinensis</i>	7	1.5%	2	0	7	Not available	Provisional
<i>Felis catus</i>	2	0.4%	1	0	2	Not available	Provisional
Total (N)	475	100%	—	15	460	—	—

Non-dominant counts (17 events across 3 smaller labels) are **provisional** — with *n* this small a few misclassifications visibly change the shares (see §2). 'Not available' human-review reflects that no human-review record exists in the system; adding one would populate that column and the final status.

Activity and monitoring figures

Daily alert events by species (coverage span; a hatched baseline marks a day AFFECTED by proven downtime — outages are often partial-day, so this does not mean the whole day was down — distinct from a genuinely quiet day):




■ Columba palumbus ■ Corvus corone ■ Sciurus carolinensis ■ Felis catus ■ day affected by proven downtime


Daily monitoring status (one cell per day; the hatched share of a cell is the portion of that day affected by proven downtime — often partial — distinct from no-data):



Activity by hour of day (local; hours with no bar had no detections in this window):



Species composition of alert events (automated classifications; horizontal bars with exact counts so minor labels stay readable — a pie would hide them):



Methodology — detection-event definition

One **alert event** is a single upstream alert: one (alert-rule × image) match delivered by the detector for one triggering image. It is a *record of a classification event*, not a count of animals: one animal can raise many alert events, and two animals in one frame raise a single event.

Temporally adjacent images may therefore represent the **same animal or visit**. **Alert events have not been grouped into independent visits**. Repeated alerts may therefore represent the same animal or visit, and the totals in this report are raw alert-event counts, not counts of independent encounters.

Interpretation of automated classifications

Inferences the detection data supports, for the monitoring operator — read as automated-classification ACTIVITY at one monitored point, not confirmed presence (§6).

- **Dominant automated classification — *Columba palumbus*.** 458 of 475 events (96.4%) on 18 of 19 days with data: the upstream classifier assigned this label to the overwhelming majority of the alert events in this window.
- **Infrequent but recurring — *Corvus corone* on 4 days (8 events) and *Sciurus carolinensis* on 2 days (7 events).** Low counts, but spread across multiple distinct days rather than single episodes: recurring, low-frequency automated classifications across multiple days.
- **Management-relevant, provisional — *Felis catus*.** The most management-relevant detection in this window — a potential predator at a feeder — but it rests on only 2 events with an unverified identification (§2), so treat it as provisional pending image review.

These describe automated classifications and their recorded activity at a single monitored point — not confirmed presence, abundance, or wider site biodiversity.

Evidence limitations

The limitations on this evidence, consolidated. Short reminders elsewhere refer back here.

- **One fixed monitoring point** — a single camera at one location; no wider-site coverage.
- **Feeder-facing** — detections are what passed one feeder, not a site-level or transect survey.
- **Automated, unverified identifications** — labels are model outputs; no competent-ecologist verification.

- **Limited independent corroboration** — BioCLIP did not corroborate 460 of 475 labels (97%); results are provisional pending image review.
- **System downtime** — no proven downtime was evidenced in this window; absence during downtime is not evidence of animal absence.
- **Short monitoring window** — 19 days in summer; not a full-year or multi-season survey.
- **Incomplete time-of-day representation** — detections span 05:00–23:00 local; hours beyond that span are unrepresented, not evidenced as inactive.
- **Counts are alert events, not animals** — not abundance and not a biodiversity value.
- **Possible repeated detections of the same visit** — events are not grouped into independent encounters (see the detection-event definition).
- **No habitat-condition assessment** — a camera does not assess habitat condition.
- **No biodiversity-unit calculation** — this is not a statutory Biodiversity Gain Plan.

Recommended monitoring actions

Actionable next steps. Priority follows documented rules (High = addresses the provisional-quality risk or missing required metadata; Medium = routine continuation; Low = conditional). No named person is assigned; the responsible role is the monitoring operator.

Recommended action	Evidence / reason	Trigger	Responsible role	Priority	Status
Human-review the non-dominant detections	Provisional low-count labels (<i>Corvus corone</i> , <i>Sciurus carolinensis</i> and <i>Felis catus</i>)	Before acting on any label	Monitoring operator	High	Open
Record the camera OS grid reference	Grid reference is required and not recorded	Before the next submission	Monitoring operator	High	Open
Human-review labels not corroborated by BioCLIP	BioCLIP did not corroborate most upstream labels	Before relying on any species-level result	Monitoring operator	High	Open
Continue monitoring across seasons	This is a short single-season window	Ongoing	Monitoring operator	Medium	Open
Compare equivalent future windows against this baseline	A baseline is established (§3)	At each future monitoring window	Monitoring operator	Medium	Open
Inspect the feeder and camera after unexpected composition changes	Composition change may be a system artefact, not ecology	On a marked change in the label mix	Monitoring operator	Medium	Open
Undertake a habitat survey where habitat condition is required	A camera does not assess habitat condition	If condition data is needed	Monitoring operator	Low	Open
Review repeated <i>Felis catus</i> classifications before considering mitigation	Provisional predator classification at a feeder	Only after confirmed ID and repeated records	Monitoring operator	High	Open

7. Data sharing

This monitoring data could, in principle, be shared with the **Local Environmental Records Centre (LERC)** for the area. However, the species labels in this report are **automated and**

unverified, and are **not ready for external ecological recording** as they stand: any submission would first require **human verification of the identifications and appropriate preparation** by a competent recorder. This is a statement of possibility only; this report shares no data and takes no action.

Appendix A — Representative image evidence

Representative-image evidence was unavailable during report generation (no readable image files were accessible to the report generator). Each non-dominant classification would otherwise show a frame with its timestamp, upstream label and confidence, BioCLIP result and confidence, and human-review status.

Appendix B — Definitions, metadata and data-quality rules

B1 Calculation definitions

- **Alert events per 100 confirmed operational hours** = $100 \times (\text{alert events}) \div (\text{confirmed operational hours})$. Confirmed operational hours = coverage span (first-to-last detection, clamped to the window) – proven downtime.
- **Active days per 10 calendar days in the coverage span** = $10 \times (\text{calendar days with } \geq 1 \text{ detection}) \div (\text{calendar days in the coverage span})$. The denominator is calendar days, not confirmed-operational days.
- **Operational coverage** = confirmed operational hours $\div$ coverage-span hours.
- **BioCLIP corroboration method** = BioCLIP's single best (top-1) label is resolved to a scientific binomial and compared for **exact equality** with the upstream label's binomial; both are mapped through the shared alias table, which absorbs known synonyms. A match is **corroborated**, anything else is **not corroborated** (never asserted as an incorrect label). The comparison is label-only: the 0.5 low-confidence flag is a separate display flag and does **not** enter the corroboration join, and top-1 (not top-k) is used. BioCLIP is an automated model — corroboration is not human or ecological verification.

B2 Proven downtime — dates and durations

No proven downtime was evidenced in this window.

B3 Model and threshold metadata

- **BioCLIP model:** Not recorded
- **Upstream classifier / model version:** Not recorded (not transmitted in the alert).
- **Low-confidence flag threshold:** 0.5
- **Upstream confidence:** stored as delivered — 2 dp, best-per-label only.

B4 Data-quality status rules

- Overall status = **Verified** if every classification is human-reviewed, **Partially verified** if some are, otherwise **Provisional**.
- No human-review record exists in the system, so the status is **Provisional** and per-row human-review shows 'Not available'.

B5 Event-grouping rule

- Alert events are **not** grouped into independent visits/encounters; totals are raw alert-event counts. Temporally adjacent events may be the same animal or visit.

B6 Report generation

- **Report version:** 2.0
- **Generated:** 2026-09-17 (Europe/London)

- **Data source:** dedicated recipient mailbox (TrapTracker RT alert emails).