

Detection report — all monitored species

Window: 2026-06-28 to 2026-07-16 (local dates, Europe/London)

Alert events: 475


Summary

Over the 19-day window from 2026-06-28 to 2026-07-16, a total of 475 alert events were recorded across all monitored species. The busiest days were 2026-06-29 (84), 2026-07-02 (68), and 2026-07-07 (49). The first activity was recorded on 2026-06-28 (21), and the last on 2026-07-16 (10). There were no alert events recorded on 2026-07-06. The overall trend across the window was a decline in alert events.

What this supports: strong evidence of sustained, near-daily *Columba palumbus* activity at a single feeder-facing camera — recorded on 18 of 19 days with data (2026-06-28 to 2026-07-16), busiest 29 Jun–7 Jul 2026; 3 other species also detected (unconfirmed).

Day-by-day alert-event counts

The 475 dated event(s); this table sums to 475. Counts are alert events, not animals — see honesty notes below.



The bars are the SAME per-day counts as the table below — discrete daily alert-event counts, not a trend line. No values are derived, smoothed, or interpolated; every bar equals its table row.

Date (local dates, Europe/London)	Alert events
2026-06-28	21
2026-06-29	84
2026-06-30	46
2026-07-01	18
2026-07-02	68
2026-07-03	26
2026-07-04	28
2026-07-05	3
2026-07-06	0
2026-07-07	49
2026-07-08	19
2026-07-09	17
2026-07-10	21

Date (local dates, Europe/London)	Alert events
2026-07-11	18
2026-07-12	11
2026-07-13	19
2026-07-14	10
2026-07-15	7
2026-07-16	10
Total (dated)	475

Monitoring effort

The requested window spans **456h**. Of that, the dataset's coverage — from the first to the last detection delivered (2026-06-28 09:24 → 2026-07-16 12:15 UTC), a span of **434.9h** (not the calendar range those dates touch) — carries the data; the remaining **21h** fall outside it, before the first or after the last detection, where monitoring status is **unknown**, not zero activity.

Within that span, proven system downtime accounts for **0.0h**, so the system was **confirmed operational for 434.9 of 434.9 hours (100%)**.


*The daily counts above are **raw over this effort** — deliberately NOT divided by operational hours. With one camera, one location, and no control for time-of-day, an events-per-hour rate would not be a sound figure; the uptime is given so the counts can be read against the effort behind them, not turned into a rate here.*

How these events were dated

475 event(s) had no decodable capture time and are placed by alert send time instead, flagged in the data rather than dropped.

Time-of-day distribution

Dated alert events by hour of capture, local time (Europe/London). N = 475. Events, not animals.



Detections concentrate in the 05:00–23:59 window, peaking at 13:00, with none overnight — consistent with a diurnal species and daytime-only capture.

Hours (local)	Alert events
00:00–02:59	0
03:00–05:59	10
06:00–08:59	125
09:00–11:59	112
12:00–14:59	86

Hours (local)	Alert events
15:00–17:59	43
18:00–20:59	83
21:00–23:59	16
Total	475

This is detection TIMING under whatever the camera's trigger/schedule was — not a behavioural rate, and still uncorrected for effort-by-hour (the system was not equally up every hour). Events, not animals.

Period-over-period comparison

No comparable prior period exists in the data — the previous equal-length period (**9–27 Jun 2026**) is entirely before the earliest dated all monitored species record held (2026-06-28), so no change can be computed. (This window: **28 Jun–16 Jul 2026**, 475 dated alert events.)

*Compared on **dated** alert events only. This window's figure (**475**) is the day-by-day table's dated total; the prior figure (**0**) is the same measure over the previous period. Undated events (no usable timestamp) and unmapped detections are excluded from BOTH — they cannot be placed in a period — and are disclosed separately.*

Species breakdown


Every dated event in the window, grouped by species key. The counts below sum to the day-by-day table's dated total (475). Alert events, not animals; local dates, Europe/London.

Species / key	Alert events	Active days
<i>Columba palumbus</i>	458	18
<i>Corvus corone</i>	8	4
<i>Sciurus carolinensis</i>	7	2
<i>Felis catus</i>	2	1
Total	475	

*This all-species total counts **every** key. If any unmapped: (unrecognised upstream tokens) or non-biological classes (e.g. Person, Car) are present, they are included here — whereas a single-species report **excludes** them from its count and discloses them separately. So a per-species figure need not sum to this total without them. This window contains none, so every row here is a mapped species.*

Species composition of alert events

Composition of detection EVENTS, not of animals or abundance — one event is one (alert-rule × image) match; repeat visits and multiple animals in frame are not resolved (Decision 4). N = 475 dated events.



- *Columba palumbus* — 458 (96.4%)
- *Corvus corone* — 8 (1.7%)
- *Sciurus carolinensis* — 7 (1.5%)
- *Felis catus* — 2 (0.4%)

Species IDs are automated and mostly unconfirmed: the BioCLIP cross-check did not confirm 460 of 475 (97%) — see 'Data-quality flags'. Treat the non-dominant counts (17 events across 3 smaller species) as **PROVISIONAL** — with *n* this small, a few misclassifications visibly change the shares.

Species	Alert events	Share of N
<i>Columba palumbus</i>	458	96.4%
<i>Corvus corone</i>	8	1.7%
<i>Sciurus carolinensis</i>	7	1.5%
<i>Felis catus</i>	2	0.4%
Total (N)	475	100%

Single camera, single location, feeder-facing: this measures what was detected at one point, not the site's community composition.

Non-target detections — for review

These species IDs are automated and unconfirmed (the cross-check did not confirm 460 of 475, 97%; see 'Data-quality flags'). Management-relevant detections below — a predator or a non-native species — must be checked against the actual images before any predator- or invasive-management decision. Do not act on the automated label alone.

Sciurus carolinensis — 7 events · 🐿 grey squirrel — non-native/invasive in the UK: - 04 Jul 2026 15:39 (local) - 04 Jul 2026 15:56 (local) - 04 Jul 2026 15:58 (local) - 04 Jul 2026 17:43 (local) - 10 Jul 2026 05:50 (local) - 10 Jul 2026 20:13 (local) - 10 Jul 2026 20:13 (local)

Felis catus — 2 events · 🐈 domestic cat — predation signal at a feeder: - 04 Jul 2026 14:24 (local) - 04 Jul 2026 14:31 (local)

Corvus corone — 8 events: - 29 Jun 2026 08:47 (local) - 29 Jun 2026 14:36 (local) - 30 Jun 2026 08:05 (local) - 30 Jun 2026 10:11 (local) - 30 Jun 2026 10:50 (local) - 03 Jul 2026 08:58 (local) - 04 Jul 2026 12:50 (local) - 04 Jul 2026 15:57 (local)

Cross-check summary

The cross-check did not corroborate 460 of 475 detections (97%) — BioCLIP's own top-1 identification was not the upstream species (460 disagree). This is a dataset-wide quality fact; the per-detection rationales are in the individual per-species reports.

Taxonomic distance of the disagreements — how far BioCLIP's identification sat from the upstream label:

BioCLIP's read vs the upstream label	Events	Share of disagreements
same genus, different species	49	11%
same family, different genus	38	8%
same order, different family	2	0%
same class, different order	157	34%
different class of animal	45	10%
BioCLIP read a plant, fungus or other non-animal	169	37%

System downtime

No delivery silence in this window was followed by back-dated captures, the only evidence that distinguishes an outage from a genuinely quiet period. Any quiet stretch here is therefore **undetermined** — a real lull and undetected downtime cannot be told apart from this data.

What these figures mean

- Each of the **475 alert events** here is an alert-event count, not an animal — one alert event is one (alert rule x image) match at best-per-label; two animals in one frame register as a single event, and other species in frame, sub-threshold detections, and repeat visits are not represented (Decision 4). So this is 475 events, not 475 animals.
- Each of the **475 dated events** in the table above is placed by CAMERA CAPTURE TIME, decoded from the image filename and converted from Europe/London local time to UTC (Decision 3, amended: capture time is the preferred temporal key where available; alert send time is still stored, never overwritten). Capture time was validated against the timestamp burned into the image pixels. Where no capture time could be decoded the alert send time is used instead and

the row is flagged. Send time is a poor proxy for capture time — usually close, but delivery is not guaranteed: an outage or backlog flushes a queue of older captures under one send timestamp, mis-dating events by whole days through delivery latency rather than midnight-boundary rounding. Note that the Europe/London -> UTC conversion uses real DST rules (zoneinfo), not a fixed offset, but it is VALIDATED ONLY WITHIN BST: the evaluated dataset (2026-06-28..2026-07-16) predates the October DST change, so autumn/winter (GMT) captures, and local times that are ambiguous or non-existent across a DST transition, remain unvalidated and are flagged per row.

- Every event here (**475 events** in total) was ingested by a parser whose NOMINAL alert format was validated field-for-field against a real captured alert email on 2026-07-16 (Decision 7); edge-case variants (zero or missing attachments, malformed body, test emails) remain validated against the reconstruction only.
- This report found **no prior equal-length period with data** to weigh the window against; a period-over-period change is the difference between two alert-event counts for the two dated windows named above — not evidence of a trend, a seasonal pattern, or a change in animal abundance; with the little history held here it describes only those two specific windows (Change 2). The earliest dated record held for this class is 2026-06-28.