

Detection report — wood pigeon (*Columba palumbus*)

Window: 2026-06-28 to 2026-07-16 (local dates, Europe/London)

Alert events: 458


Summary

During the monitoring window from 2026-06-28 to 2026-07-16, a total of 458 alert events were recorded, with a noticeable decline in activity towards the end of the window. The busiest days were 2026-06-29 (82), 2026-07-02 (68), and 2026-07-07 (49). The first activity was recorded on 2026-06-28 (21), while the last activity was on 2026-07-16 (10). There was one zero-activity day, which was 2026-07-06.

What this supports: strong evidence of sustained, near-daily *Columba palumbus* activity at a single feeder-facing camera — recorded on 18 of 19 days with data (2026-06-28 to 2026-07-16), busiest 29 Jun–7 Jul 2026.

Day-by-day alert-event counts

The 458 dated event(s); this table sums to 458. Counts are alert events, not animals — see honesty notes below.



The bars are the SAME per-day counts as the table below — discrete daily alert-event counts, not a trend line. No values are derived, smoothed, or interpolated; every bar equals its table row.

Date (local dates, Europe/London)	Alert events
2026-06-28	21
2026-06-29	82
2026-06-30	43
2026-07-01	18
2026-07-02	68
2026-07-03	25
2026-07-04	20
2026-07-05	3
2026-07-06	0
2026-07-07	49
2026-07-08	19
2026-07-09	17
2026-07-10	18

Date (local dates, Europe/London)	Alert events
2026-07-11	18
2026-07-12	11
2026-07-13	19
2026-07-14	10
2026-07-15	7
2026-07-16	10
Total (dated)	458

Monitoring effort

The requested window spans **456h**. Of that, the dataset's coverage — from the first to the last detection delivered (2026-06-28 09:24 → 2026-07-16 12:15 UTC), a span of **434.9h** (not the calendar range those dates touch) — carries the data; the remaining **21h** fall outside it, before the first or after the last detection, where monitoring status is **unknown**, not zero activity.

Within that span, proven system downtime accounts for **0.0h**, so the system was **confirmed operational for 434.9 of 434.9 hours (100%)**.


*The daily counts above are **raw over this effort** — deliberately NOT divided by operational hours. With one camera, one location, and no control for time-of-day, an events-per-hour rate would not be a sound figure; the uptime is given so the counts can be read against the effort behind them, not turned into a rate here.*

How these events were dated

458 event(s) had no decodable capture time and are placed by alert send time instead, flagged in the data rather than dropped.

Time-of-day distribution

Dated alert events by hour of capture, local time (Europe/London). N = 458. Events, not animals.



Detections concentrate in the 05:00–23:59 window, peaking at 13:00, with none overnight — consistent with a diurnal species and daytime-only capture.

Hours (local)	Alert events
00:00–02:59	0
03:00–05:59	9
06:00–08:59	122
09:00–11:59	110
12:00–14:59	82

Hours (local)	Alert events
15:00–17:59	38
18:00–20:59	81
21:00–23:59	16
Total	458

This is detection TIMING under whatever the camera's trigger/schedule was — not a behavioural rate, and still uncorrected for effort-by-hour (the system was not equally up every hour). Events, not animals.

Period-over-period comparison

No comparable prior period exists in the data — the previous equal-length period (**9–27 Jun 2026**) is entirely before the earliest dated wood pigeon record held (2026-06-28), so no change can be computed. (This window: **28 Jun–16 Jul 2026**, 458 dated alert events.)

*Compared on **dated** alert events only. This window's figure (**458**) is the day-by-day table's dated total; the prior figure (**0**) is the same measure over the previous period. Undated events (no usable timestamp) and unmapped detections are excluded from BOTH — they cannot be placed in a period — and are disclosed separately.*

Notable observations

Computed from the stored detection records — no model involvement. Same basis as the table above: alert events (not animals), local dates, Europe/London, capture time where it was decoded.

Busiest day: 29 Jun 2026 — 82 alert events.

Quietest active day: 5 Jul 2026 — 3 alert events.

Days with no wood pigeon alert events: 1 of 19 in the window — but they do not all mean the same thing: - **1 undetermined** — no proven outage, but nothing of ANY species arrived either, so a real absence and undetected downtime cannot be told apart.

Non-target activity: 17 alert events from 3 other species in the same window. - **Corvus corone** — 8 alert events on 29 Jun, 30 Jun, 3 Jul, 4 Jul 2026 - **Sciurus carolinensis** — 7 alert events on 4, 10 Jul 2026 - **Felis catus** — 2 alert events on 4 Jul 2026

System downtime: none detected — no delivery silence in this window was followed by back-dated captures, which is the only evidence that distinguishes an outage from a genuinely quiet period.

wood pigeon not detected, system status unconfirmed: 2 periods with no alert of ANY species and no proven outage, so a real absence and undetected downtime cannot be told apart. Reported as undetermined rather than resolved by assumption. - 05 Jul 07:06 → 07 Jul 12:20 UTC (53.2h) - 14 Jul 19:51 → 15 Jul 22:30 UTC (26.6h)

Cross-check agreement is reported under 'Data-quality flags' below, where the per-detection rationales live.

Unverified model descriptions (not observations)

None available for these events. The computed highlights above are unaffected.

Data-quality flags

The cross-check did not corroborate 443 of 458 detections (97%) — BioCLIP's own top-1 identification was not the upstream species. The verdict is species-level: a near miss in the same genus is still a disagreement, so the distance below is what says how serious each one is. - **same genus, different species (46, 10% of disagreements)** - 2026-06-28: upstream

'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Columba livia domestica' is a different taxon (same_genus) - 2026-06-28: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Columba oenas' is a different taxon (same_genus) - **same family, different genus (38, 9% of disagreements)** - 2026-06-29: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Streptopelia dusumieri' is a different taxon (same_family) - 2026-06-29: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Streptopelia dusumieri' is a different taxon (same_family) - **same class, different order (154, 35% of disagreements)** - 2026-06-28: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Ptilonorhynchus violaceus' is a different taxon (same_class) - 2026-06-28: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Apus apus' is a different taxon (same_class) - **different class of animal (39, 9% of disagreements)** - 2026-06-29: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP 'Oryctolagus cuniculus domesticus' -> Oryctolagus cuniculus (other_animal) - 2026-06-29: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Nebrioporus nipponicus' is a different taxon (other_animal) - **BioCLIP read a plant, fungus or other non-animal (166, 37% of disagreements)** - 2026-06-28: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Amanita populiphila' is a different taxon (non_animal) - 2026-06-28: upstream 'ColumbaPalumbus' -> Columba palumbus, but BioCLIP's top-1 'Poa pratensis' is a different taxon (non_animal) **Enrichment not run (VLM on 458 event(s))**: ingested without enrichment (e.g. via store), not a model failure.

What these figures mean

- Each of the **458 alert events** here is an alert-event count, not an animal — one alert event is one (alert rule x image) match at best-per-label; two animals in one frame register as a single event, and other species in frame, sub-threshold detections, and repeat visits are not represented (Decision 4). So this is 458 events, not 458 animals.
- Each of the **458 dated events** in the table above is placed by CAMERA CAPTURE TIME, decoded from the image filename and converted from Europe/London local time to UTC (Decision 3, amended: capture time is the preferred temporal key where available; alert send time is still stored, never overwritten). Capture time was validated against the timestamp burned into the image pixels. Where no capture time could be decoded the alert send time is used instead and the row is flagged. Send time is a poor proxy for capture time — usually close, but delivery is not guaranteed: an outage or backlog flushes a queue of older captures under one send timestamp, mis-dating events by whole days through delivery latency rather than midnight-boundary rounding. Note that the Europe/London -> UTC conversion uses real DST rules (zoneinfo), not a fixed offset, but it is VALIDATED ONLY WITHIN BST: the evaluated dataset (2026-06-28..2026-07-16) predates the October DST change, so autumn/winter (GMT) captures, and local times that are ambiguous or non-existent across a DST transition, remain unvalidated and are flagged per row.
- Every event here (**458 events** in total) was ingested by a parser whose NOMINAL alert format was validated field-for-field against a real captured alert email on 2026-07-16 (Decision 7); edge-case variants (zero or missing attachments, malformed body, test emails) remain validated against the reconstruction only.
- This report found **no prior equal-length period with data** to weigh the window against; a period-over-period change is the difference between two alert-event counts for the two dated windows named above — not evidence of a trend, a seasonal pattern, or a change in animal abundance; with the little history held here it describes only those two specific windows (Change 2). The earliest dated record held for this class is 2026-06-28.