

Ancient Anorthosites on Mars: Widespread in the Lower Crust

Michael S. Phillips¹
Christina E. Viviano²
A. Deanne Rogers³
Linae Larson¹
Livio Tornabene⁴
Alexander Trowbridge⁵
Jeffrey E. Moersch⁶
Harry Y. McSween Jr.⁶

June 4, 2025

¹Lunar and Planetary Laboratory, University of Arizona, Tucson, AZ, USA.

²Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA.

³Department of Geosciences, Stony Brook University, Stony Brook, NY, USA.

⁴Department of Earth Sciences, Western University, London, ON, Canada.

⁵Southwest Research Institute, San Antonio, Texas, United States

⁶Department of Earth, Environmental, and Planetary Sciences, University of Tennessee, Knoxville, TN, USA.

Abstract

We report extensive detections of feldspathic, likely anorthositic, lithologies within outcrops uplifted by the Argyre basin-forming impact on Mars. The radial position of feldspathic outcrops from the basin center indicates that they are stratigraphically above olivine-bearing material and below low-calcium pyroxene-bearing rocks, and the inferred depths suggest that feldspathic lithologies are present in the lower martian crust in the circum-Argyre region. This stratigraphy bears striking resemblance to that observed at the Hellas basin rim and indicates that feldspathic rocks could be a pervasive component of the lower crust across the entire southern highlands of Mars. These results are consistent with evidence from geophysical data suggesting a lower-density lower crust, but likely not sufficient to fully account for the discrepancy in density. Importantly, these feldspathic rocks are present alongside pyroxene- and olivine-bearing units, a suite of compositions commonly associated with layered mafic intrusions (LMI). We suggest that a process early in Mars' history produced large volumes of melt that cooled slowly to form LMI-like complexes and that these complexes represent an aspect of primary martian crust. The melt-producing process ostensibly affected the whole southern hemisphere, and possibly the globe. These observations offer direct evidence to support and refine existing models of early crust formation, addressing key questions regarding crustal density variations and formation mechanisms that have previously lacked widespread observational constraints.

Main

In contrast to Earth and Venus, Mars preserves a surface rock record dating back to ~ 4.55 Ga (Humayun et al., 2013; Kruijer et al., 2020). The preservation and exposure of ancient rocks (≥ 4.1 Ga, pre-Noachian) at the surface provides a unique opportunity to study early crustal formation through remote sensing and surface investigations. Understanding how and when evolved rocks first formed on terrestrial planets remains a major open question in planetary geology. Since the processes that formed Mars' primary crust share more similarities with Earth- and Venus-sized bodies than with lunar-sized bodies (Elkins-Tanton, 2012), studying the oldest Martian crust provides critical insights into early crustal evolution with potential for comparative planetology.

Noteworthy among the ancient rocks of Mars are compositions apparently dominated by plagioclase feldspar (Fig. 1). Ancient feldspathic and silicic rocks measured *in situ* provide clear evidence that crustal differentiation occurred early in Mars' history (e.g., Stolper et al., 2013; Sautter et al., 2015), although the formation mechanism for these rocks was likely fractional crystallization of basaltic magmas (Udry et al., 2018; Ostwald et al., 2022) rather than a plate-tectonic-style process (Sautter et al., 2015, 2016) akin to the formation of continental crust on Earth. Whether such differentiation occurred only at isolated locations or globally is an open question. Here, we present evidence for a widespread, plagioclase-rich component to the lower martian crust that is likely buried throughout the entire southern highlands. Our results suggest a hemispheric, if not global, process that resulted in differentiated magmas over ~ 4.1 billion years ago on Mars.

Though increasingly recognized as important constituents of the Martian crust, feldspathic rocks are relatively rare compared to basaltic materials based on orbital observations (Ehlmann and Edwards, 2014; McSween, 2015; McSween et al., 2009, 2023). Many detections of plagioclase-rich compositions come from isolated exposures in Noachian and older terrains in the southern highlands, interpreted as possible intrusive bodies (Quantin et al., 2012; Carter and Poulet, 2013; Wray et al., 2013; Sautter et al., 2016; Viviano-Beck et al., 2017; Payré et al., 2022). Other plagioclase-rich occurrences are linked to likely volcanic units found in Hesperian-aged terrains, such as Syrtis Major (Wray et al., 2013; Eggers et al., 2021; Rogers and Farrand, 2022), western Arcadia Planitia (Farrand et al., 2021), and the Eridania basin (Michalski et al., 2023).

Other studies have revealed more extensive occurrences of plagioclase-rich outcrops. Phillips et al. (2022) identified widespread plagioclase-rich outcrops across the northern rim of the Hellas basin. Viviano-Beck et al. (2017) initially reported, and Flahaut et al. (2023) corroborated, the presence of extensive plagioclase-rich compositions in the lower walls of Valles Marineris. The Valles Marineris exposures are plausibly Noachian or older, and are interpreted to be intrusive bodies such as sills or other hypabyssal intrusions (Viviano-Beck et al., 2017), although some evidence may suggest they could alternatively be effusive volcanic rocks (Flahaut et al., 2023). Additionally, nine feldspar-rich sites were identified in the Terra Sirenum/Cimmeria region by Payré et al. (2022), an area with anomalous geochemical and magnetic signatures and suggested to be an ancient crustal block (Bouley et al., 2020). Therefore, though plagioclase-rich rocks are found across the southern highlands of Mars, it appears that a diversity of petrogenetic processes—including plutonic and volcanic activity, and possibly impact-related processes (Grieve et al., 1991; Hurwitz and Kring, 2014; Vaughan and Head, 2014; Koeppel et al., 2020)—have contributed to their formation from the pre-Noachian through the early Hesperian Epochs. This work focuses on the most ancient (pre-Noachian) plagioclase-rich outcrops.

The lower crust of Mars has been hypothesized to be feldspar-rich to explain its apparent low density based on geochemical and geophysical data (Baratoux et al., 2014; Goossens et al., 2017; Knapmeyer-Endrun et al., 2021). Despite the growing number of observations of plagioclase-rich rocks, it remains unclear from surface remote-sensing observations whether a lower-crust plagioclase-rich component is present globally. Here, we report results from an investigation into outcrops uplifted from depth by the Argyre basin-forming impact to test the hypothesis that the martian crust contains a global feldspathic component. Following the identification of abundant plagioclase-rich outcrops along the northern Hellas rim (Phillips et al., 2022), we find that plagioclase-rich outcrops are also ubiquitous in Argyre-uplift materials and share many similarities with those discovered along the northern Hellas rim, supporting the hypothesis that these materials are widespread in the lower martian crust.

Plagioclase-rich outcrops sandwiched between mantle and upper crust

We investigated the composition of uplifted material surrounding Argyre using ~ 180 m/pixel, 72-band multispectral mapping data (MRDRv4, Seelos et al. 2023) from the Compact Recon-

naissance Imaging Spectrometer for Mars (CRISM, Murchie et al., 2007) instrument onboard the Mars Reconnaissance Orbiter (Zurek and Smrekar, 2007). Reflectance spectroscopy is a useful tool for identifying many silicate minerals, including major rock-forming minerals such as olivine, pyroxene, Fe-bearing plagioclase, and oxides (Adams and Filice, 1967; Adams, 1968; Clark and Roush, 1984; Burns, 1993).

We examined 40 CRISM MRDR tiles (Extended Data Table 1) for the presence of primary rock forming silicate minerals. Of the 1349 outcrops identified to contain signatures of primary rock forming minerals, 476 showed absorption features consistent with olivine + high-calcium pyroxene (HCP), 140 were consistent with type 1 olivine, 320 with type 2 olivine, 282 with low-calcium pyroxene (LCP), and 131 were consistent with plagioclase (Fig. 2; see Methods for details on spectral categorization). Type 1 olivine is interpreted as more Mg-rich or smaller grained and type 2 olivine is more Fe-rich or larger grained (Viviano et al., 2014; Ody et al., 2013).

The emplacement mechanism of uplifted massifs resulting from basin-forming impacts, such as the one that formed Argyre, is primarily attributed to structural uplift (Tanaka and Leonard, 1995; Johnson et al., 2016) and to a lesser extent ejected blocks (Johnson et al., 2016). Structural uplift refers to the combined effects of collapse of a large central uplift – analogous to the formation of peak ring structures in complex craters (Johnson et al., 2016) – and isostatic rebound and crustal thinning from transient crater formation (Tanaka and Leonard, 1995), which leads to an extensional tectonic regime that causes normal faulting. The relative stratigraphy of material uplifted this way would be preserved (with some disruption) as opposed to the inverted stratigraphy expected for an overturned flap (Kenkmann et al., 2014). Ejected blocks of material would also maintain the trend of deeper material emplaced closer to the center of the basin, although more mixing of materials originating from different depths would occur than for structurally uplifted massifs (Extended Data Fig. 1, Trowbridge et al. *submitted*). Uplifted materials closer to the basin center would thus tend to be stratigraphically below those further from the basin center. In this work, the distance of each outcrop from the basin center is used as a proxy for pre-uplift stratigraphy (Fig. 2).

Olivine-bearing units (type 1, type 2, and olivine + HCP) vastly dominate the compositions exposed within approximately 720 km of the Argyre basin center (Fig. 2). It is likely that some of these olivine-bearing compositions represent uplifted mantle material (Extended Data Fig. 1; Koeppen and Hamilton, 2008; Ody et al., 2013). Among the olivine-bearing categories, olivine + HCP outcrops are concentrated lowest along the rim (based on area-weighted average distance calculations, Extended Data Fig. 2) where possible mantle material is expected, raising the possibility of wehrlite/lherzolite in the upper martian mantle (McGetchin and Smith, 1978; Treiman, 2025; Bernadet et al., 2025). Beyond ~ 720 km, LCP is the most abundant mineral category. The position of LCP-dominated compositions is suggestive of upper pre-Noachian crust (Buczkowski et al., 2010; McNeil et al., 2025). The material between these two compositions is interpreted as a comminuted mixture of mantle, lower-, and upper-pre-Noachian crust.

The plagioclase detections described here are concentrated between ~ 330 and 890 km from the basin center and are absent beyond 890 km. The highest relative proportion of plagioclase

class ($\sim 29\%$) occurs between 430 and 470 km from the basin center. The overall pattern revealed by the distribution of mineral categories with radial distance (Fig. 2) suggests a clear transition from compositions dominated by olivine at lower stratigraphic positions to those dominated by LCP in the upper stratigraphic positions (Extended Data Fig. 2). This transition occurs between ~ 450 and 810 km distance from the Argyre basin center. Plagioclase-rich rocks appear to be positioned lower in the crust alongside the stratigraphically lower olivine-bearing rocks, some of which may represent mantle materials, and below the stratigraphically higher LCP-bearing rocks (Fig. 2, Extended Data Fig. 2). The gross stratigraphy observed in the circum-Argyre uplifted materials is consistent with the stratigraphy observed in the Hellas basin rim (Phillips et al., 2022) and with the well-preserved stratigraphic relationships exposed in the lower walls of Valles Marineris (Viviano-Beck et al., 2017). The similarity of the observed stratigraphy at Argyre to the intact stratigraphy in Valles Marineris lends support to the interpretation of stratigraphic relationships in the circum-Argyre region, despite the complex emplacement processes associated with the basin-forming impact.

Local contexts, visual appearance and color analysis

Three general categories were identified for outcrops when considering their local geomorphological context: massifs, crater rims, and valley walls. Circum-Argyre massifs are the principal target of interest for our study because they directly represent pre-Argyre rocks excavated by the impact process. We include crater-rim outcrops in our analysis because impacts on the Argyre rim could uplift material that was originally uplifted by Argyre. Exposures within valley walls are also of interest because valleys cut into the circum-Argyre material and thereby expose buried rock.

Plagioclase-rich outcrops on massifs typically appear as light-toned, ridge-like capping units overlying other mineral detections even within the same massifs (Extended Data Figs. 3 and 4). Many plagioclase-rich outcrops exhibit massive, homogeneous light-toned appearances that resemble terrestrial massif anorthosites or thick anorthositic members of layered mafic intrusions (Fig. 1, Extended Data Fig. 3; Ashwal 1993). Massive textured outcrops also appear as rounded or angular blocks (Extended Data Fig. 3) reminiscent of the “mega breccia” reported in the circum-Isidis region by Scheller and Ehlmann (2020); however, the “mega breccia” along the Argyre rim are larger in scale. Thermal Emission Imaging System (THEMIS; Christensen et al. 2004) thermal inertia (Ferguson et al., 2006) and nighttime infrared data (Edwards et al., 2011) over massif outcrops indicate high thermal inertia values relative to surrounding units, suggesting they are well-consolidated exposures rather than loose, unconsolidated materials.

The spatial relationships of neighboring outcrops (defined here as outcrops within 5-km distance of one another) were assessed using a graph network analysis to understand patterns in local associations and relative elevations (Extended Data Fig. 4). The most common neighbor of plagioclase-bearing outcrops (aside from other plagioclase-bearing outcrops) are type 2 olivine and LCP (23 and 22 counts each), followed by type 1 olivine (12 counts), and olivine + HCP (9 counts). The proximal relationships indicate a spatial association among plagioclase, (Fe) olivine and LCP (that is, these compositions commonly appear on or near

the same massifs) and that plagioclase is locally higher in elevation than its mafic neighbors (Extended Data Fig. 4). This may evidence local (kilometer-scale) stratigraphic relationships that are distinct from the basin-wide distribution of minerals that likely represents the overall pre-Argyre crustal stratigraphy.

The plagioclase-bearing outcrops associated with craters that we identified in this study are found exclusively on crater rims, not within crater floors, suggesting they represent uplifted material rather than later volcanic or sedimentary infill or local differentiated impact melt or breccias. Valley walls primarily expose material in the outer regions of the Argyre rim. Though LCP dominates these exposures, plagioclase and olivine are detected in valleys lower down on the rim, consistent with the basin-wide stratigraphic relationship observed in the region.

Analysis of High Resolution Imaging Science Experiment (HiRISE; McEwen et al., 2010) infrared, red, blue-green (IRB) color data reveal that plagioclase-rich outcrops consistently appear as higher reflectance, light-toned materials relative to surrounding terrain (Fig. 1, Extended Data Figs. 3, 5 and 6). Dark subtraction methods (Tornabene et al., 2018) were applied to three HiRISE images (ESP_020799_1385, ESP_031269_1395, and ESP_069064_1390) to remove additive atmospheric residuals in the data and assess the colors associated with CRISM-identified compositions. Outcrops categorized as plagioclase with CRISM show lower BG/IR and Red/IR band ratio values than neighboring mafic outcrops (Extended Data Fig. 5), attributable to an absorption feature present with LCP and olivine but not present with plagioclase that overlaps with the HiRISE IR channel (centered at ~ 900 nm). Light-toned outcrops that were either too small to be resolved by CRISM MRDR data or not covered by the data were also investigated with HiRISE color parameters and show the same relative trend among light- and dark-toned material within the same scene (Extended Data Figs. 5 and 6), which suggests that the CRISM-based detections of plagioclase-rich outcrops are a minimum estimate of the full distribution of such outcrops in the circum-Argyre region.

Possible lithology of plagioclase-bearing outcrops

The plagioclase-rich outcrops are characterized by a broad absorption feature centered around $1.25 \mu\text{m}$, indicative of Fe^{2+} -bearing plagioclase feldspar (Adams and Filice, 1967; Adams et al., 1973; Adams and Goullaud, 1978). This spectral signature is observed if plagioclase is the dominant mineral visible at the surface; small amounts ($\sim 15\%$) of mafic minerals like pyroxene or olivine can mask the plagioclase absorption feature (Crown and Pieters, 1987; Pompilio et al., 2007; Serventi et al., 2013, 2015; Cheek and Pieters, 2014). However, specific, grain-size dependent, combinations of plagioclase with other minerals may also yield a reflectance spectrum dominated by the $1.25\text{-}\mu\text{m}$ absorption feature of plagioclase (“spectrally feldspathic” rocks), which means the $1.25\text{-}\mu\text{m}$ feature of Fe-plagioclase is not sufficient to classify the lithology of the host rock (Rogers and Nekvasil, 2015; Barthez et al., 2023; Vannier et al., 2024). Supplementary Note 1 of Phillips et al. (2022) presents a full discussion of this topic.

THEMIS thermal infrared data provide additional constraints on the composition of spectrally feldspathic outcrops. We assessed nine plagioclase outcrops across four THEMIS scenes

(Extended Data Table 2; see Methods for details on processing) as well as 23 type 1 olivine, 15 type 2 olivine, 50 HCP + olivine, and 19 LCP outcrops within the same THEMIS scenes. Two metrics that track the silica content of target lithologies were calculated: the weighted absorption center (WAC; Smith et al. 2013; Amador and Bandfield 2016) and cubic spline fit minimum (CSFM; Rogers and Nekvasil 2015). Plagioclase-bearing outcrops show WAC values ranging from $\sim 9.9 - 10.2 \mu\text{m}$ with a $10.1 \mu\text{m}$ median and CSFM values ranging from $\sim 9.3 - 11.0$ with a median of $\sim 9.5 \mu\text{m}$. Plotting WAC versus CSFM further elucidates the spectral character of the outcrops. Fig. 3 shows WAC vs. CSFM for THEMIS observations compared to reference spectral library data of felsic, intermediate, mafic, ultramafic, anorthositic and gabbroic rocks. The THEMIS data for plagioclase-bearing outcrops align with intermediate, gabbroic, and anorthositic compositions, but not with felsic compositions.

The reflectance and thermal infrared spectral characteristics of plagioclase-bearing outcrops, their close spatial association with pyroxene and olivine outcrops at $\leq 5\text{-km}$ spatial scales, higher relative position compared to olivine and pyroxene in local contexts on average, and the dark and light-toned layering observed within individual plagioclase-bearing units are consistent with a differentiated magmatic complex. The preferred interpretation of these rocks is that they represent layered mafic intrusion (LMI) style complexes (Fig. 4). Anorthositic layers in LMIs form through fractional crystallization and flotation of plagioclase in a cooling, typically hypabyssal, mafic magma body (Ashwal, 1993). The modest percentage of plagioclase-rich outcrops at any radial distance from Argyre (interquartile range = $1.3 - 7.1\%$ of total detections at distances where plagioclase is present) is consistent with the typical proportion of anorthositic rocks in terrestrial LMIs (Ashwal, 1993, 2021). However, we cannot completely rule out contributions from previous impact-basin melt bodies, which may also undergo differentiation, concentrate minerals and form cumulate rocks akin to LMI.

Alternatively, the light-toned, massive appearance of many plagioclase-rich outcrops in the Argyre region could be interpreted as evidence for massif-type anorthosites (Ashwal, 1993; Arndt, 2013; Keller et al., 2024). Massif-type anorthosites on Earth are distinct from anorthosite formation as layers in LMIs. Terrestrial massif-type anorthosites form large, relatively homogeneous bodies that can extend over thousands of square kilometers, whereas anorthositic components of LMIs typically form discrete layers that can be hundreds of kilometers in extent (e.g., Bushveld Complex, Latypov et al. 2020) and kilometers thick (e.g., Mineral Lake intrusion, Wisconsin; Ashwal 1993). Hypotheses for massif-type anorthosite formation involve an elevated crustal temperature gradient, extensive, long-lived mafic magmas, and density-driven extraction and ascent of plagioclase crystal mushes (Emslie et al., 1994; Keller et al., 2024). The tectonic setting for massif-type anorthosites is rifting at a prior convergent zone, a setting unlikely for Mars, and the associated rock types are mangerite (feldspar and orthopyroxene), charnockite (quartz, feldspar, and orthopyroxene), and granitoids (Emslie et al., 1994; Hamilton et al., 2010; Arndt, 2013; Keller et al., 2024). This suite of rocks, referred to as AMCG, are not observed in the circum-Argyre region, further leading us to favor differentiation from basaltic melts into LMI-style intrusions rather than a massif-type anorthosite interpretation for the observations.

Hypotheses for the origin of the stratigraphy exposed at the Argyre rim

The observed mineralogical distribution around the Argyre basin provides insight into the nature and origin of Mars’ earliest crust. The striking similarities between plagioclase-rich outcrops exposed at both Argyre and Hellas basins—separated by over 4,000 km—suggest they represent the same component of the lower Martian crust. Spectral, morphological, and contextual characteristics of these outcrops are consistent across both regions, including their association with olivine and pyroxene-bearing units, their light-toned appearance, presence of both layered and massive textures, and their relative stratigraphic position along the basin rims (Phillips et al., 2022). The spatial extent indicates these compositions are prevalent beneath much of the southern highlands, if not globally. The creation of such a widespread and compositionally consistent crustal component implies vast volumes of magma derived from a similar source. We favor two scenarios that could explain the generation of such enormous melt volumes across the southern hemisphere: 1) overturn of a post-magma-ocean cumulate pile (Borg and Draper, 2003; Elkins-Tanton et al., 2005); 2) melt and magma generation from basin-forming and/or frequent impacts (Koeppel et al., 2020; Manske et al., 2021; Black and Marchi, 2024); or alternatively 3) partial melting of the southern hemisphere’s lower crust (Bonnet Gibet et al., 2025; Bernadet et al., 2025).

Magma Ocean and Mantle Processes.

A whole mantle magma ocean would have formed a gravitationally unstable cumulate mantle with Mg-rich cumulates below Fe-rich cumulates (Elkins-Tanton et al., 2005). Subsequent mantle overturn would have generated large volumes of melt through decompression. These melts would have ascended and cooled slowly in the crust, consistent with formation of differentiated igneous complexes through fractional crystallization. A degree-one overturn geometry (Borg and Draper, 2003; Elkins-Tanton et al., 2005; Debaille et al., 2008) would have resulted in similar crustal compositions across the southern hemisphere, which more easily explains our observations than a higher degree overturn geometry (Scheinberg et al., 2014) that would have produced spatially variable crustal compositions across the southern highlands.

Impact Melt Differentiation.

A second hypothesis involves the differentiation of a single massive, or many overlapping, impact melt sheet(s). The Borealis basin-forming impact, which is a leading hypothesis for the creation of Mars’ crustal dichotomy (Don and Steven, 1984; Andrews-Hanna et al., 2008; Marinova et al., 2008; Nimmo et al., 2008), would have generated an enormous volume of melt. Differentiation of this melt sheet could have produced anorthositic rocks through plagioclase flotation (Koeppel et al., 2020; Manske et al., 2021), later uplifted by the Argyre and Hellas impacts. This scenario is particularly intriguing given that the impact melt is expected to concentrate primarily within the impact basin (Marinova et al., 2008; Ballantyne

et al., 2023) and the feldspathic outcrops are found in the southern hemisphere. This raises the possibility that if a large impact formed the crustal dichotomy, it may have impacted the southern hemisphere rather than the northern (Reese et al., 2010; Golabek et al., 2011; Leone et al., 2014; Ballantyne et al., 2023; Cheng et al., 2024), creating a localized magma ocean in the southern hemisphere.

The cumulative effect of high frequency bombardment of early Mars would likely have created massive volumes of melt (both impact melt and crustal partial melts) even in the absence of a single, large, dichotomy-forming impact (Koeppel et al., 2020; Manske et al., 2021; Black and Marchi, 2024). The expected fate of such melts is ascent to a depth of neutral buoyancy in the crust, contributing to a compositional stratification of the early crust (Black and Marchi, 2024). Additionally, differentiation of surface melts could contribute to smaller scale compositional layering (Koeppel et al., 2020).

Crustal Melting.

A third possibility is that evolved compositions formed through partial melting of hydrothermally altered mafic crust (Bonnet Gibet et al., 2025; Bernadet et al., 2025). We disfavor this scenario as an explanation for the pre-Noachian feldspathic crust because silicic (i.e., felsic) rocks would be the expected product of the partial melting—compositions not observed in this study.

Two plausible scenarios for crustal melting have recently been proposed. The first involves crustal thickness differences between Martian hemispheres. In the southern highlands, where crust is thicker, higher concentrations of heat-producing elements create steeper temperature gradients. This thermal insulation, combined with thinner lithosphere and higher mantle melt fractions beneath these regions, enhances temperatures at the crustal base, promoting partial melting at depths up to approximately 15 km (Bonnet Gibet et al., 2022, 2025). The resulting evolved magmas would ascend, potentially undergo assimilation and fractional crystallization, and crystallize at various crustal depths—a process sharing characteristics with silicic rock formation in Iceland (Jónasson, 2007; Martin and Sigmarsson, 2007, 2010).

The second mechanism involves interaction between basaltic melts and serpentinized primordial crust (Bernadet et al., 2025), plausible for early Mars given evidence for widespread alteration (Bibring et al., 2006; Bishop et al., 2008; Ehlmann et al., 2009; Ehlmann and Edwards, 2014; Wordsworth et al., 2017; Viviano et al., 2023; Carter et al., 2023; Royer et al., 2024) and the possibility of a dense steam atmosphere persisting for several million years post-accretion (Abe, 2011; Scherf and Lammer, 2020). In this model, water both enables serpentinization of ultramafic protocrust and significantly lowers rock melting temperatures. Thermodynamic modeling indicates a minimum water content of ~ 100 ppm in the bulk silicate Mars would be necessary (Bernadet et al., 2025). Through shallow (< 15 km) fluid-assisted reactions between serpentinite and basaltic melts, felsic compositions could form. Models suggest this process might generate up to $\sim 40\%$ of Martian felsic crust, particularly during two periods: very early Martian history (4.4-4.5 Ga) and during the Hesperian Epoch (Bernadet et al., 2025).

However, the spectral characteristics of our detections are more consistent with intermediate/gabbroic/anorthositic rocks than with silicic/felsic compositions. Though partial melting of possibly hydrated crust may explain some Hesperian-aged spectrally feldspathic detections (Farrand et al., 2021; Rogers and Farrand, 2022; Michalski et al., 2023), it seems less applicable to the pre-Noachian compositions presented here. For the few silicic rocks encountered *in situ* on Mars, fractional crystallization of basaltic magmas may better explain their formation than hydrous crustal melting (Christensen et al., 2005; Udry et al., 2018; Ostwald et al., 2022).

The presence of Fe^{2+} in plagioclase, which enables its spectral identification, provides additional petrogenetic information. Fe^{2+} incorporation into plagioclase is enhanced under reducing conditions, with high melt polymerization (viz., higher SiO_2), low water activity, high $\text{Fe}^{Tot}/\text{Al}_2\text{O}_3$, and in high temperature melts that rapidly ascend to cool at low pressures (Wenk and Wilde, 1973; Longhi et al., 1976; Smith and Brown, 1988; Phinney, 1992; Ashwal, 1993; Lundgaard and Tegner, 2004). Though the presence of Fe^{2+} in plagioclase is not diagnostic of the petrogenetic environment, favorable conditions for Fe^{2+} substitution into plagioclase include dry, high temperature, rapidly ascending melts that undergo fractional crystallization at low pressures and allow for the extraction of plagioclase as floatation layers. Such conditions are more consistent with either a mantle overturn scenario or impact-generated melts than with partial melting of hydrated crust.

The above mechanisms need not be mutually exclusive. The early Martian crust likely formed through a complex interplay of processes operating at different spatial and temporal scales. Cumulate mantle overturn may have produced voluminous melts that solidified into differentiated igneous complexes while concurrent impact events generated substantial near-surface melt across the planet.

In the elevated geothermal and water-rich environment of early Mars, surface volcanics and shallow crustal materials would have undergone hydrothermal alteration (Viviano and Phillips, 2019). After subsequent burial and diagenesis, these altered materials could have experienced partial melting, producing intermediate to evolved compositions. The spatial distribution of these evolved compositions would likely be heterogeneous, with concentrations in regions of thicker crust or areas with significant hydrothermal activity. Such a multi-stage petrogenetic history is likely necessary to explain the diverse suite of feldspathic materials observed across Mars (Carter and Poulet, 2013; Wray et al., 2013; Rogers and Nekvasil, 2015; Viviano-Beck et al., 2017; Farrand et al., 2021; Payré et al., 2022, 2024; Phillips et al., 2022; Flahaut et al., 2023; Michalski et al., 2023).

Implications for crustal formation, density, and remnant magnetism

The presence of anorthositic rocks and associated mafic lithologies in ancient crustal materials uplifted by the Argyre impact has implications for understanding Mars' early crustal evolution. The similar detection of plagioclase-rich materials at both Argyre and Hellas (Phillips et al., 2022) suggests this was a hemispheric, if not global, component of Mars' lower crust. This supports models of early Martian evolution involving hemisphere-scale or global-scale production of magma, though the specific mechanism remains uncertain. These

observations show crustal differentiation can occur without plate tectonics – possibly via fractional crystallization of large mafic magma bodies – a process which may be applicable to early Earth (Kranendonk, 2010).

Though plagioclase-rich rocks are present, they constitute only about 4% by area of the pre-Argyre crust mapped with CRISM data in this study. If this proportion is representative of the true proportion of plagioclase-rich rocks in the lower Martian crust, then it is unlikely that anorthositic rocks can explain the overall lower density of Mars’ lower crust inferred from geochemical and geophysical data (Baratoux et al., 2014; Goossens et al., 2017; Knapmeyer-Endrun et al., 2021; Drilleau et al., 2023; Joshi et al., 2023). Additional factors such as increased porosity, alteration, or hydration, may be necessary to explain the lower crustal density (Ehlmann et al., 2010; Bandfield et al., 2013; Sun and Milliken, 2015; Wordsworth et al., 2017; Goossens et al., 2017; Scheller and Ehlmann, 2020; Wieczorek et al., 2022; Carrasco et al., 2023). Alternatively, the olivine- and LCP-bearing compositions detected in our study may contain a significant feldspar component that is not reflected in their spectral signatures due to the masking effect of mafic minerals (e.g., Extended Data Fig. 7). Such a scenario would lead to an underestimation of the total feldspathic component present in Argyre uplift materials. Indeed, the character of the THEMIS spectra suggest that some of these rocks follow a more gabbroic than ultramafic trend (Fig. 3). However, further detailed analyses are necessary to determine whether troctolite or leucogabbro/leuconorite are present in the uplifted materials surrounding Argyre (see Extended Data Fig. 7).

The igneous complexes suggested by our observations may also have implications for Mars’ remnant crustal magnetism. Layered mafic intrusions and massif anorthosites on Earth often contain significant quantities of iron-oxide minerals such as magnetite and titanomagnetite, which can carry strong remanent magnetization. The shallow differentiated igneous complexes that may have formed during Mars’ early history, when the planet had an active dynamo, could contain the minerals necessary to carry the strong crustal magnetic anomalies observed in the southern highlands.

Conclusions

Detection of widespread plagioclase-rich outcrops within material uplifted by the Argyre basin-forming impact provides compelling evidence for the presence of feldspathic, possibly anorthositic, rocks as a component of Mars’ lower crust. The association of these plagioclase-rich materials with olivine and pyroxene-bearing outcrops suggests they represent components of ancient, differentiated complexes of either igneous or impact origin (or both) that formed during Mars’ early crustal development.

Observations of crustal compositions indicative of LMI-like complexes support models of early Martian evolution involving large-scale melting (Fig. 4), though the specific mechanism—whether mantle overturn, impact melting, or other processes—remains open to debate. The presence of similar plagioclase-rich materials uplifted by both the Argyre and Hellas impacts suggests their formation required a hemispheric or global process rather than isolated activity. This evidence provides constraints for refining theoretical and computational models of early Mars crust formation and offers insights into differentiation processes on terrestrial

planets lacking plate tectonics.

Missions targeting ancient crustal materials could provide additional constraints on their composition, age, and formation mechanisms. For example, detections of sulfide, spinel and oxide horizons in association with layered intrusive outcrops would support an interpretation that these rocks represent differentiated complexes. The Perseverance rover at Jezero crater has recently observed feldspar dominated (Bedford et al., 2024; Schmidt et al., 2025) and spinel-bearing rocks (Mansbach et al., 2024; Bedford et al., 2024; Lawson et al., 2025) that could represent material uplifted by the Isidis-forming impact, warranting further investigation.

Methods

Reflectance Spectroscopic Analyses

We used the Compact Reconnaissance Imaging Spectrometer for Mars (CRISM; Murchie et al., 2007) onboard the Mars Reconnaissance Orbiter to identify plagioclase-rich outcrops in the circum-Argyre region. CRISM is a visible and near-infrared imaging spectrometer that provides high-resolution reflectance spectral data across a wavelength range of approximately 400 to 4000 nm. The CRISM data used in this study are the 72-band (410 – 3923 nm), 180 m/pixel Map-Projected Multispectral Reduced Data Record (MRDR) v4 data (Seelos et al., 2023; Murchie, 2024). The MRDR map tiles were analyzed using the Spectral Cube Analysis Tool (Phillips, 2024). Regions of interest were drawn using combinations of CRISM spectral parameters (Viviano et al., 2014) and spectra were then averaged from the ROIs and ratioed to in-scene spectrally bland regions – a common technique in CRISM analysis to remove scene-dependent instrument and atmospheric artifacts in the data. Interpretations of mineralogy present in each spectrum were made through visual comparison to reference spectral libraries (Minerals identified with CRISM Analysis [MICA], Viviano et al. 2014; USGS Spectral Library v7, Kokaly et al. 2017).

Plagioclase was identified based on the presence of a broad absorption feature centered at approximately 1.25- μm (Fig. 3). Plagioclase spectra were differentiated from olivine spectra based on a reflectance peak near 900 nm that is present in plagioclase but absent in olivine. This diagnostic feature highlights the importance of using "joined" CRISM reflectance data products (that include both short and long detector channels) to identify plagioclase spectra. Pyroxene detections were categorized as low-calcium pyroxene (LCP) and high-calcium pyroxene (HCP) based on the minimum positions of the ~ 1 - and ~ 2 - μm absorption bands (Viviano et al., 2014). Where the pyroxene category presented as mixtures of olivine and pyroxene, pyroxene composition was determined solely by the presence and position of the 2- μm band. Olivine detections were categorized as "Type 1" and "Type 2" olivine (Ody et al., 2013) based on a k-means classification 5 band parameters (band center, band depth, band asymmetry, slope between 1 and 1.3 μm , and full width half max) that quantify properties of the ~ 1 - μm olivine absorption feature (Extended Data Fig. 8). The complete python code and data to replicate the classification are attached to this article. Type 1 olivine is con-

sidered more forsteritic and Type 2 olivine spectra indicate either more fayalitic or coarser grain-size olivine; a coarse-grained low-iron olivine is spectrally similar to a fine-grained high-iron olivine (e.g., Mustard et al., 2005), which motivates the compositionally agnostic “Type 1” and “Type 2” designations (Ody et al., 2013). Spectra with a broad 1- μm olivine absorption and a broad 2- μm absorption consistent with HCP were categorized as Olivine + HCP (Extended Data Fig. 7). Spectra were further categorized as Low, Moderate, or High confidence based on the strength and clarity of absorption features present on a qualitative basis (Fig. 3). Geologic units from Dohm et al. (2015) were used to assign each mineral unit with an associated geologic context: either uplifted by the Argyre impact (Argyre basin and rim materials, Argyre basin members) or uplifted by later large impacts (old and young crater materials) that excavated Argyre rim materials (Extended Data Table 3).

Thermal Emissivity Analyses

In addition to reflectance spectroscopy, 100 m/pixel, 9-band thermal emissivity data from the Thermal Emission Imaging System (THEMIS) onboard the 2001 Mars Odyssey orbiter (Christensen et al., 2004) were used to constrain the silica content of select spectrally feldspathic outcrops. THEMIS data with the warmest available surface temperatures were processed following the methods of Bandfield et al. (2004). In the absence of alteration, the wavelength of minimum emissivity is negatively correlated with the silica content of a rock (Lyon, 1965). Locations analyzed were ensured to indicate no evidence for alteration (hydration bands) in the co-located CRISM spectra. Felsic, mafic, and ultramafic rocks show minima at $\sim 9.6\ \mu\text{m}$, $\sim 10.5\ \mu\text{m}$, and $11\ \mu\text{m}$, respectively (Fig. 3). Notably, anorthositic compositions can display minima consistent with mafic/gabbroic compositions due to their similar silica contents (Fig. 3).

To assess the silica content of outcrops identified with CRISM, we calculated two metrics for the THEMIS emissivity minimum position: the cubic spline fit minimum (CSFM) and the weighted absorption center (WAC). The CSFM method fits a cubic spline to the THEMIS emissivity data for bands 3 – 8 ($\lambda = 7.89 - 12.56\ \mu\text{m}$) and returns the minimum of the cubic spline fit values (Rogers and Nekvasil, 2015). The WAC method calculates the wavelength value at which the cumulative absorption area equals half the total absorption area (Vincent and Thomson, 1972; Smith et al., 2013; Amador and Bandfield, 2016). The WAC metric has less variance than the CSFM and is more robust against idiosyncrasies in wavelength minimum values that may arise from a cubic spline fit; however, both measurements correlate inversely with the total silica content of the rock (Vincent and Thomson, 1972; Walter and Salisbury, 1989; Rogers and Nekvasil, 2015; Amador and Bandfield, 2016).

The relationship between CSFM and WAC can further elucidate the spectral character of rocks, particularly for anorthositic and gabbroic compositions compared to intermediate rocks. We calculated WAC and CSFM values for THEMIS-convolved reference spectra from the Arizona State University (ASU) spectral library (ASU), RELAB database (Brown University), Ward’s spectral library (Bryson), and from Dunn (2021). CSFM values show a wider range than WAC values (Fig. 3B). For felsic and intermediate rocks, CSFM values are lower than the corresponding WAC values for the same rock. For mafic, ultramafic,

gabbroic, anorthositic, and pyroxenitic rocks the CSFM–WAC values show an approximate 1:1 relationship with some CSFM values being higher and some lower than the corresponding WAC values. For middling WAC values (~ 10.1 to $10.6 \mu\text{m}$) that could be interpreted as multiple rock types (e.g., intermediate, anorthositic, gabbroic), a relatively high CSFM value would indicate a more anorthositic or gabbroic rock (Fig. 3B).

Assessment of Color, Geomorphological and Thermophysical Properties

We used the 5 m/pixel Context Camera (CTX) global mosaic (Malin et al., 2007; Dickson et al., 2018) and 0.25 m/pixel HiRISE images (McEwen et al., 2010) to assess the color and texture of outcrops. For HiRISE IRB color analysis, dark subtraction methods were used after Tornabene et al. (2018) to correct for additive atmospheric residuals in the scenes that affect the spectral shape of the data. HiRISE scene ESP_031269_1395 overlaps with CRISM-identified plagioclase, LCP, and HCP + olivine outcrops. Regions of interest (ROIs) were drawn from areas overlapping the CRISM-detections to investigate the HiRISE spectral character and support interpretations of similarly colored outcrops in nearby HiRISE scenes (ESP_020799_1385, ESP_031269_1395) as representative of plagioclase-bearing or mafic-bearing compositions (LCP or olivine $\pm$ HCP).

Thermal inertia properties were examined using THEMIS global thermal inertia mosaic (Ferguson et al., 2006; Edwards et al., 2009) and nighttime infrared data (Edwards et al., 2011) to distinguish between relatively consolidated and unconsolidated materials.

Data availability

All data used in this study are available from public repositories. CRISM data are available from the Planetary Data System (PDS) Geosciences Node. THEMIS, CTX, and HiRISE data are available from their respective PDS nodes. Processed data products, including interpreted spectral detection maps with associated metadata and THEMIS convolved reference spectra, as well as code used to produce the figures in this publication, are available at <https://doi.org/10.5281/zenodo.15596017>.

Acknowledgments

This research utilizes spectra acquired with the NASA RELAB facility at Brown University. We are grateful to Dr. James Dohm for sharing GIS files from his 2015 published map of the Argyre basin (Dohm et al., 2015). This work was funded by NASA Mars Data Analysis Program grant no. 80NSSC20K0938 awarded to M. Phillips. [Thank you to reviewers once reviewed].

References

- Abe, Y. (2011). Protoatmospheres and surface environment of protoplanets. *Earth, Moon, and Planets*, 108:9–14.
- Adams, J. B. (1968). Lunar and martian surfaces - petrologic significance of absorption bands in the near-infrared. *Science*, 159:1453–1455.
- Adams, J. B., Bell, P. M., Conel, J. E., Mao, H. K., McCord, T. B., and Nash, D. B. (1973). Visible and near-infra-red transmission and reflectance measurements of the luna 20 soil. *Geochimica et Cosmochimica Acta*, 37:731–743.
- Adams, J. B. and Filice, A. L. (1967). Spectral reflectance 0.4 to 2.0 microns of silicate rock powders. *Journal of Geophysical Research*, 72.
- Adams, J. B. and Goullaud, L. H. (1978). Plagioclase feldspars: Visible and near infrared diffuse reflectance spectra as applied to remote sensing. In *Lunar and Planetary Science Conference*, pages 2901–2909.
- Amador, E. S. and Bandfield, J. L. (2016). Elevated bulk-silica exposures and evidence for multiple aqueous alteration episodes in nili fossae, mars. *Icarus*, 276:39–51.
- Andrews-Hanna, J. C., Zuber, M. T., and Banerdt, W. B. (2008). The borealis basin and the origin of the martian crustal dichotomy. *Nature*, 453:1212.
- Arndt, N. (2013). The formation of massif anorthosite: Petrology in reverse. *Geoscience Frontiers*, 4:195–198.
- Ashwal, L. D. (1993). *Anorthosites*. Springer-Verlag, Berlin.
- Ashwal, L. D. (2021). Anorthosites. In Alderton, D. and Elias, S. A., editors, *Encyclopedia of Geology*, pages 130–144. Academic Press, Oxford, second edition.
- Ballantyne, H. A., Jutzi, M., Golabek, G. J., Mishra, L., Cheng, K. W., Rozel, A. B., and Tackley, P. J. (2023). Investigating the feasibility of an impact-induced martian dichotomy. *Icarus*, 392:115395.
- Bandfield, J. L., Edwards, C. S., Montgomery, D. R., and Brand, B. D. (2013). The dual nature of the martian crust: Young lavas and old clastic materials. *Icarus*, 222:188–199.
- Bandfield, J. L., Rogers, D., Smith, M. D., and Christensen, P. R. (2004). Atmospheric correction and surface spectral unit mapping using thermal emission imaging system data. *Journal of Geophysical Research: Planets*, 109.
- Baratoux, D., Samuel, H., Michaut, C., Toplis, M. J., Monnereau, M., Wieczorek, M., Garcia, R., and Kurita, K. (2014). Petrological constraints on the density of the martian crust. *Journal of Geophysical Research: Planets*, 119:1707–1727.

- Barthez, M., Flahaut, J., Guitreau, M., Ito, G., and Pik, R. (2023). Understanding vnir plagioclase signatures on mars through petrographic, geochemical, and spectral characterization of terrestrial feldspar-bearing igneous rocks. *Journal of Geophysical Research: Planets*, 128:e2022JE007680.
- Bedford, C. et al. (2024). Investigating the source of unusual kaolinite-spinel float rocks in jezero crater, mars, and their implications for mars crustal processes. In *GOLDSCHMIDT*.
- Bernadet, J., Borisova, A. Y., Guitreau, M., Safonov, O. G., Asimow, P., Nédélec, A., Bohrsen, W. A., Kosova, S. A., and de Parseval, P. (2025). Making continental crust on water-bearing terrestrial planets. *Science Advances*, 11:eads6746.
- Bibring, J.-P. et al. (2006). Global mineralogical and aqueous mars history derived from omega/mars express data. *Science*, 312:400–404.
- Bishop, J. L. et al. (2008). Phyllosilicate diversity and past aqueous activity revealed at mawrth vallis, mars. *Science*, 321:830–833.
- Black, B. A. and Marchi, S. (2024). Buoyant impact partial melts on ancient mars. *Journal of Geophysical Research: Planets*, 129:e2023JE008040.
- Bonnet Gibet, V., Michaut, C., Bodin, T., Wieczorek, M., and Dubuffet, F. (2025). Martian highlands differentiation concomitant to dichotomy formation. *Journal of Geophysical Research: Planets*, 130:e2024JE008486.
- Bonnet Gibet, V., Michaut, C., Wieczorek, M., and Lognonné, P. (2022). A positive feedback between crustal thickness and melt extraction for the origin of the martian dichotomy. *Journal of Geophysical Research: Planets*, 127:e2022JE007472.
- Borg, L. E. and Draper, D. S. (2003). A petrogenetic model for the origin and compositional variation of the martian basaltic meteorites. *Meteoritics & Planetary Science*, 38:1713–1731.
- Bouley, S. et al. (2020). A thick crustal block revealed by reconstructions of early mars highlands. *Nature Geoscience*, 13:105–109.
- Buczkowski, D. L., Murchie, S., Clark, R., Seelos, K., Seelos, F., Malaret, E., and Hash, C. (2010). Investigation of an argyre basin ring structure using mars reconnaissance orbiter/compact reconnaissance imaging spectrometer for mars. *Journal of Geophysical Research: Planets*, 115.
- Burns, R. G. (1993). *Mineralogical Applications of Crystal Field Theory*. Cambridge University Press, Cambridge.
- Carrasco, S., Knapmeyer-Endrun, B., Margerin, L., Xu, Z., Joshi, R., Schimmel, M., Stutzmann, É., Charalambous, C., Lognonné, P., and Banerdt, W. B. (2023). Constraints for the martian crustal structure from rayleigh waves ellipticity of large seismic events. *Geophysical Research Letters*, 50.

- Carter, J. and Poulet, F. (2013). Ancient plutonic processes on mars inferred from the detection of possible anorthositic terrains. *Nature Geoscience*, 6:1008.
- Carter, J., Riu, L., Poulet, F., Bibring, J.-P., Langevin, Y., and Gondet, B. (2023). A mars orbital catalog of aqueous alteration signatures (mocaas). *Icarus*, 389:115164.
- Cheek, L. C. and Pieters, C. M. (2014). Reflectance spectroscopy of plagioclase-dominated mineral mixtures: Implications for characterizing lunar anorthosites remotely. *American Mineralogist*, 99:1871–1892.
- Cheng, K. W., Rozel, A. B., Golabek, G. J., Ballantyne, H. A., Jutzi, M., and Tackley, P. J. (2024). Mars’s crustal and volcanic structure explained by southern giant impact and resulting mantle depletion. *Geophysical Research Letters*, 51:e2023GL105910.
- Christensen, P. R. et al. (2004). The thermal emission imaging system (themis) for the mars 2001 odyssey mission. *Space Science Reviews*, 110:85–130.
- Christensen, P. R. et al. (2005). Evidence for magmatic evolution and diversity on mars from infrared observations. *Nature*, 436:504.
- Clark, R. N. and Roush, T. L. (1984). Reflectance spectroscopy: Quantitative analysis techniques for remote sensing applications. *Journal of Geophysical Research: Solid Earth*, 89:6329–6340.
- Crown, D. A. and Pieters, C. M. (1987). Spectral properties of plagioclase and pyroxene mixtures and the interpretation of lunar soil spectra. *Icarus*, 72:492–506.
- Debaille, V., Yin, Q. Z., Brandon, A. D., and Jacobsen, B. (2008). Duration of a magma ocean and subsequent mantle overturn in mars: Evidence from nakhlites. In *Lunar and Planetary Science Conference XXXIX*, page Abstract #1837.
- Dickson, J. L., Kerber, L., Fassett, C. I., and Ehlmann, B. (2018). A global, blended ctx mosaic of mars with vectorized seam mapping: A new mosaicking pipeline using principles of non-destructive image editing. In *Lunar and Planetary Science Conference XLIX*, page Abstract #2480.
- Dohm, J. M. et al. (2015). Geological and hydrological histories of the argyre province, mars. *Icarus*, 253:66–98.
- Don, E. W. and Steven, W. S. (1984). The martian hemispheric dichotomy may be due to a giant impact. *Nature*, 309:138.
- Drilleau, M., Beucler, É., Shi, J., Knapmeyer-Endrun, B., Garcia, R. F., Ansan, V., Kim, D., Lognonné, P., and Banerdt, W. B. (2023). Structure of the martian crust below insight from surface waves and body waves generated by nearby meteoroid impacts. *Geophysical Research Letters*, 50:e2023GL104601.
- Dunn, T. L. (2021). Thermal emission spectra of terrestrial alkaline rocks.

625 Edwards, C. S., Bandfield, J. L., Christensen, P. R., and Fergason, R. L. (2009). Global
626 distribution of bedrock exposures on mars using themis high-resolution thermal inertia.
627 *Journal of Geophysical Research: Planets*, 114.

628 Edwards, C. S., Nowicki, K. J., Christensen, P. R., Hill, J., Gorelick, N., and Murray, K.
629 (2011). Mosaicking of global planetary image datasets: 1. techniques and data processing
630 for thermal emission imaging system (themis) multi-spectral data. *Journal of Geophysical*
631 *Research: Planets*, 116.

632 Eggers, G. L., Wray, J. J., and Dufek, J. (2021). Compositional mapping of the nili patera
633 feldspathic unit: Extent and implications for formation. *Journal of Geophysical Research:*
634 *Planets*, 126.

635 Ehlmann, B. L. and Edwards, C. S. (2014). Mineralogy of the martian surface. *Annual*
636 *Review of Earth and Planetary Sciences*, 42:291–315.

637 Ehlmann, B. L. et al. (2009). Identification of hydrated silicate minerals on mars using mro-
638 crism: Geologic context near nili fossae and implications for aqueous alteration. *Journal*
639 *of Geophysical Research: Planets*, 114.

640 Ehlmann, B. L., Mustard, J. F., and Murchie, S. L. (2010). Geologic setting of serpentine
641 deposits on mars. *Geophysical Research Letters*, 37.

642 Elkins-Tanton, L. T. (2012). Magma oceans in the inner solar system. *Annual Review of*
643 *Earth and Planetary Sciences*, 40:113–139.

644 Elkins-Tanton, L. T., Hess, P. C., and Parmentier, E. M. (2005). Possible formation of
645 ancient crust on mars through magma ocean processes. *Journal of Geophysical Research:*
646 *Planets*, 110.

647 Emslie, R. F., Hamilton, M. A., and Thériault, R. J. (1994). Petrogenesis of a mid-proterozoic
648 anorthosite-mangerite-charnockite-granite (amcg) complex: Isotopic and chemical evi-
649 dence from the nain plutonic suite. *The Journal of Geology*, 102:539–558.

650 Farrand, W. H., Rice, J. W., Chuang, F. C., and Rogers, A. D. (2021). Spectral and
651 geological analyses of domes in western arcadia planitia, mars: Evidence for intrusive
652 alkali-rich volcanism and ice-associated surface features. *Icarus*, 357:114111.

653 Fergason, R. L., Christensen, P. R., and Kieffer, H. H. (2006). High-resolution thermal
654 inertia derived from the thermal emission imaging system (themis): Thermal model and
655 applications. *Journal of Geophysical Research: Planets*, 111.

656 Fergason, R. L., Hare, T. M., and Laura, J. (2018). Hrsc and mola blended digital elevation
657 model at 200m v2. Digital elevation model, Astrogeology PDS Annex, U.S. Geological
658 Survey.

659 Flahaut, J., Payet, V., Fueten, F., Guitreau, M., Barthez, M., Ito, G., and Allemand, P.
660 (2023). New detections of feldspar-bearing volcanic rocks in the walls of valles marineris,
661 mars. *Geophysical Research Letters*, 50:e2022GL100772.

- Golabek, G. J., Keller, T., Gerya, T. V., Zhu, G., Tackley, P. J., and Connolly, J. A. D. (2011). Origin of the martian dichotomy and tharsis from a giant impact causing massive magmatism. *Icarus*, 215:346–357.
- Goossens, S., Sabaka, T. J., Genova, A., Mazarico, E., Nicholas, J. B., and Neumann, G. A. (2017). Evidence for a low bulk crustal density for mars from gravity and topography. *Geophysical Research Letters*, 44:7686–7694.
- Grieve, R. A. F., Stöffler, D., and Deutsch, A. (1991). The sudbury structure: Controversial or misunderstood? *Journal of Geophysical Research: Planets*, 96:22753–22764.
- Hamilton, M. A., Scoates, J. S., and Rämö, O. T. (2010). The petrology of anorthosites, related granitic rocks, and uht assemblages: a tribute to ronald f. emslie. *The Canadian Mineralogist*, 48:705–710.
- Humayun, M. et al. (2013). Origin and age of the earliest martian crust from meteorite nwa 7533. *Nature*, 503:513–516.
- Hurwitz, D. M. and Kring, D. A. (2014). Differentiation of the south pole–aitken basin impact melt sheet: Implications for lunar exploration. *Journal of Geophysical Research: Planets*, 119:1110–1133.
- Johnson, B. C. et al. (2016). Formation of the orientale lunar multiring basin. *Science*, 354:441–444.
- Jónasson, K. (2007). Silicic volcanism in iceland: Composition and distribution within the active volcanic zones. *Journal of Geodynamics*, 43:101–117.
- Joshi, R., Knapmeyer-Endrun, B., Mosegaard, K., Wieczorek, M. A., Igel, H., Christensen, U. R., and Lognonné, P. (2023). Joint inversion of receiver functions and apparent incidence angles to determine the crustal structure of mars. *Geophysical Research Letters*, 50:e2022GL100469.
- Keller, D. S., Lee, C.-T. A., Peck, W. H., Monteleone, B. D., Martin, C., Vervoort, J. D., and Bolge, L. (2024). Mafic slab melt contributions to proterozoic massif-type anorthosites. *Science Advances*, 10:eadn3976.
- Kenkmann, T., Poelchau, M. H., and Wulf, G. (2014). Structural geology of impact craters. *Journal of Structural Geology*, 62:156–182.
- Knapmeyer-Endrun, B. et al. (2021). Thickness and structure of the martian crust from insight seismic data. *Science*, 373:438–443.
- Koeppel, A. H. D., Black, B. A., and Marchi, S. (2020). Differentiation in impact melt sheets as a mechanism to produce evolved magmas on mars. *Icarus*, 335:113422.
- Koeppen, W. C. and Hamilton, V. E. (2008). Global distribution, composition, and abundance of olivine on the surface of mars from thermal infrared data. *Journal of Geophysical Research: Planets*, 113.

698 Kokaly, R. et al. (2017). Usgs spectral library version 7 data.

699 Kranendonk, M. J. V. (2010). Two types of archean continental crust: Plume and plate
700 tectonics on early earth. *American Journal of Science*, 310:1187–1209.

701 Kruijer, T. S., Borg, L. E., Wimpenny, J., and Sio, C. K. (2020). Onset of magma ocean
702 solidification on mars inferred from mn-cr chronometry. *Earth and Planetary Science
703 Letters*, 542:116315.

704 Latypov, R., Chistyakova, S., Costin, G., Namur, O., Barnes, S., and Kruger, W. (2020).
705 Monomineralic anorthosites in layered intrusions are indicators of the magma chamber
706 replenishment by plagioclase-only-saturated melts. *Scientific Reports*, 10:3839.

707 Lawson, P. R. et al. (2025). Adaptive sampling with pixl on the mars *Perseverance* rover.
708 *Icarus*, 429:116433.

709 Leone, G., Tackley, P. J., Gerya, T. V., May, D. A., and Zhu, G. (2014). Three-dimensional
710 simulations of the southern polar giant impact hypothesis for the origin of the martian
711 dichotomy. *Geophysical Research Letters*, 41:8736–8743.

712 Longhi, J., Walker, D., and Hays, J. F. (1976). Fe and mg in plagioclase. In *Lunar and
713 Planetary Science Conference Proceedings*, volume 1, pages 1281–1300.

714 Lundgaard, K. L. and Tegner, C. (2004). Partitioning of ferric and ferrous iron between
715 plagioclase and silicate melt. *Contributions to Mineralogy and Petrology*, 147:470–483.

716 Lyon, R. J. P. (1965). Analysis of rocks by spectral infrared emission (8 to 25 microns).
717 *Economic Geology*, 60:715–736.

718 Malin, M. C. et al. (2007). Context camera investigation on board the mars reconnaissance
719 orbiter. *Journal of Geophysical Research: Planets*, 112.

720 Mansbach, E. N. et al. (2024). Likely ferromagnetic minerals identified by the perseverance
721 rover and implications for future paleomagnetic analyses of returned martian samples.
722 *Authorea*.

723 Manske, L., Marchi, S., Plesa, A.-C., and Wünnemann, K. (2021). Impact melting upon
724 basin formation on early mars. *Icarus*, 357:114128.

725 Marinova, M. M., Aharonson, O., and Asphaug, E. (2008). Mega-impact formation of the
726 mars hemispheric dichotomy. *Nature*, 453:1216–1219.

727 Martin, E. and Sigmarsson, O. (2007). Crustal thermal state and origin of silicic magma in
728 iceland: the case of torfajökull, ljósufjöll and snæfellsjökull volcanoes. *Contributions to
729 Mineralogy and Petrology*, 153:593–605.

730 Martin, E. and Sigmarsson, O. (2010). Thirteen million years of silicic magma production
731 in iceland: Links between petrogenesis and tectonic settings. *Lithos*, 116:129–144.

732 McEwen, A. S. et al. (2010). The high resolution imaging science experiment (hirise) during
733 mro’s primary science phase (psp). *Icarus*, 205:2–37.

734 McGetchin, T. R. and Smith, J. R. (1978). The mantle of mars: Some possible geological
735 implications of its high density. *Icarus*, 34:512–536.

736 McNeil, J. D., Fawdon, P., Balme, M. R., Coe, A. L., Cuadros, J., and Turner, S. M. R.
737 (2025). Dichotomy retreat and aqueous alteration on noachian mars recorded in highland
738 remnants. *Nature Geoscience*, 18:124–132.

739 McSween, H. Y. (2015). Petrology on mars. *American Mineralogist*, 100:2380–2395.

740 McSween, H. Y., Head III, J. W., Rogers, A. D., and Schmidt, M. E. (2023). Assessing
741 global trends in mars magma compositions using ground truth. *Meteoritics & Planetary
742 Science*, 58:1306–1317.

743 McSween, H. Y., Taylor, G. J., and Wyatt, M. B. (2009). Elemental composition of the
744 martian crust. *Science*, 324:736–739.

745 Michalski, J., Rogers, A., Edwards, C., Cowart, A., and Xiao, L. (2023). Diverse volcanism
746 and crustal evolution on early mars. *In Review preprint*.

747 Murchie, S. (2024). Crism data product software interface specification. Technical report,
748 PDS Geosciences Node, Washington University.

749 Murchie, S. et al. (2007). Compact reconnaissance imaging spectrometer for mars (crism)
750 on mars reconnaissance orbiter (mro). *Journal of Geophysical Research: Planets*, 112.

751 Mustard, J. F., Poulet, F., Gendrin, A., Bibring, J.-P., Langevin, Y., Gondet, B., Mangold,
752 N., Bellucci, G., and Altieri, F. (2005). Olivine and pyroxene diversity in the crust of
753 mars. *Science*, 307:1594–1597.

754 Nimmo, F., Hart, S. D., Korycansky, D. G., and Agnor, C. B. (2008). Implications of an
755 impact origin for the martian hemispheric dichotomy. *Nature*, 453:1220–1223.

756 Ody, A., Poulet, F., Bibring, J.-P., Loizeau, D., Carter, J., Gondet, B., and Langevin, Y.
757 (2013). Global investigation of olivine on mars: Insights into crust and mantle composi-
758 tions. *Journal of Geophysical Research: Planets*, 118:234–262.

759 Ostwald, A., Udry, A., Payré, V., Gazel, E., and Wu, P. (2022). The role of assimilation and
760 fractional crystallization in the evolution of the mars crust. *Earth and Planetary Science
761 Letters*, 585:117514.

762 Payré, V., Salvatore, M. R., and Edwards, C. S. (2022). An evolved early crust exposed on
763 mars revealed through spectroscopy. *Geophysical Research Letters*, 49:e2022GL099639.

764 Payré, V., Udry, A., and Fraeman, A. A. (2024). Igneous diversity of the early martian crust.
765 *Minerals*, 14:452.

766 Phillips, M. (2024). Spectral cube analysis tool: A python program for analyzing multi- and
767 hyperspectral images. In *Lunar and Planetary Institute*.

768 Phillips, M. S., Viviano, C. E., Moersch, J. E., Rogers, A. D., McSween, H. Y., and Seelos,
769 F. P. (2022). Extensive and ancient feldspathic crust detected across north hellas rim,
770 mars: Possible implications for primary crust formation. *Geology*, 50:1182–1186.

771 Phinney, W. C. (1992). Partition coefficients for iron between plagioclase and basalt as a
772 function of oxygen fugacity: Implications for archean and lunar anorthosites. *Geochimica
773 et Cosmochimica Acta*, 56:1885–1895.

774 Pompilio, L., Sgavetti, M., and Pedrazzi, G. (2007). Visible and near-infrared reflectance
775 spectroscopy of pyroxene-bearing rocks: New constraints for understanding planetary sur-
776 face compositions. *Journal of Geophysical Research: Planets*, 112.

777 Quantin, C., Flahaut, J., Clenet, H., Allemand, P., and Thomas, P. (2012). Composition
778 and structures of the subsurface in the vicinity of valles marineris as revealed by central
779 uplifts of impact craters. *Icarus*, 221:436–452.

780 Reese, C. C., Orth, C. P., and Solomatov, V. S. (2010). Impact origin for the martian crustal
781 dichotomy: Half emptied or half filled? *Journal of Geophysical Research: Planets*, 115.

782 Rogers, A. D. and Farrand, W. H. (2022). Spectral evidence for alkaline rocks and compo-
783 sitional diversity among feldspathic light-toned terrains on mars. *Icarus*, 376:114883.

784 Rogers, A. D. and Nekvasil, H. (2015). Feldspathic rocks on mars: Compositional constraints
785 from infrared spectroscopy and possible formation mechanisms. *Geophysical Research
786 Letters*, 42:2619–2626.

787 Royer, C. et al. (2024). Intense alteration on early mars revealed by high-aluminum rocks
788 at jezero crater. *Communications Earth & Environment*, 5:1–13.

789 Sautter, V. et al. (2015). In situ evidence for continental crust on early mars. *Nature
790 Geoscience*, 8:605–609.

791 Sautter, V. et al. (2016). Magmatic complexity on early mars as seen through a combination
792 of orbital, in-situ and meteorite data. *Lithos*, 254–255:36–52.

793 Scheinberg, A., Elkins-Tanton, L. T., and Zhong, S. J. (2014). Timescale and morphology
794 of martian mantle overturn immediately following magma ocean solidification. *Journal of
795 Geophysical Research: Planets*, 119:454–467.

796 Scheller, E. L. and Ehlmann, B. L. (2020). Composition, stratigraphy, and geological history
797 of the noachian basement surrounding the isidis impact basin. *Journal of Geophysical
798 Research: Planets*, 125:e2019JE006190.

799 Scherf, M. and Lammer, H. (2020). Did mars possess a dense atmosphere during the first
800 ~ 400 million years? *Space Science Reviews*, 217:2.

801 Schmidt, M. E. et al. (2025). Diverse and highly differentiated lava suite in jezero crater,
802 mars: Constraints on intracrustal magmatism revealed by mars 2020 pixl. *Science Ad-*
803 *vances*, 11:eadr2613.

804 Seelos, F. P. et al. (2023). The crism investigation in mars orbit: Overview, history, and
805 delivered data products. *Icarus*, page 115612.

806 Serventi, G., Carli, C., and Sgavetti, M. (2015). Spectral variability of plagioclase–mafic
807 mixtures (3): Quantitative analysis applying the mgm algorithm. *Icarus*, 254:34–55.

808 Serventi, G., Carli, C., Sgavetti, M., Ciarniello, M., Capaccioni, F., and Pedrazzi, G. (2013).
809 Spectral variability of plagioclase–mafic mixtures (1): Effects of chemistry and modal
810 abundance in reflectance spectra of rocks and mineral mixtures. *Icarus*, 226:282–298.

811 Smith, J. V. and Brown, W. L. (1988). *Feldspar Minerals*. Springer, Berlin, Heidelberg.

812 Smith, M. R., Bandfield, J. L., Cloutis, E. A., and Rice, M. S. (2013). Hydrated silica on
813 mars: Combined analysis with near-infrared and thermal-infrared spectroscopy. *Icarus*,
814 223:633–648.

815 Stolper, E. M. et al. (2013). The petrochemistry of jake_m: A martian mugearite. *Science*,
816 341:1239463.

817 Sun, V. Z. and Milliken, R. E. (2015). Ancient and recent clay formation on mars as revealed
818 from a global survey of hydrous minerals in crater central peaks. *Journal of Geophysical*
819 *Research: Planets*, 120:2293–2332.

820 Tanaka, K. L. and Leonard, G. J. (1995). Geology and landscape evolution of the hellas
821 region of mars. *Journal of Geophysical Research: Planets*, 100:5407–5432.

822 Tornabene, L. L. et al. (2018). Image simulation and assessment of the colour and spatial
823 capabilities of the colour and stereo surface imaging system (cassis) on the exomars trace
824 gas orbiter. *Space Science Reviews*, 214:18.

825 Treiman, A. H. (2025). The nakhlite martian meteorites are igneous, but not cumulates. In
826 *LPI*, page 1472.

827 Udry, A., Gazel, E., and McSween Jr., H. Y. (2018). Formation of evolved rocks at gale
828 crater by crystal fractionation and implications for mars crustal composition. *Journal of*
829 *Geophysical Research: Planets*, 123:1525–1540.

830 Vannier, H., Horgan, B., Greenberger, R., and Eddy, M. (2024). Assessing plagioclase
831 feldspar detectability limits in whole-rock vnir spectra: Possible lower detection limits
832 and application to mars. In *LPI*.

833 Vaughan, W. M. and Head, J. W. (2014). Impact melt differentiation in the south pole-aitken
834 basin: Some observations and speculations. *Planetary and Space Science*, 91:101–106.

835 Vincent, R. K. and Thomson, F. (1972). Spectral compositional imaging of silicate rocks.
836 *Journal of Geophysical Research (1896-1977)*, 77:2465–2472.

837 Viviano, C. E., Beck, A. W., Murchie, S. L., Dapremont, A. M., and Seelos, F. P. (2023).
838 Heterogeneity of the noachian crust of mars using crism multispectral mapping data.
839 *Geophysical Research Letters*, 50:e2022GL102711.

840 Viviano, C. E. et al. (2014). Revised crism spectral parameters and summary products based
841 on the currently detected mineral diversity on mars. *Journal of Geophysical Research:*
842 *Planets*, 119:1403–1431.

843 Viviano, C. E. and Phillips, M. S. (2019). Hydrothermal alteration and large impact basins
844 on mars. In *50th LPSC*, page Abstract #2824, Houston, TX. LPI.

845 Viviano-Beck, C. E., Murchie, S. L., Beck, A. W., and Dohm, J. M. (2017). Compositional
846 and structural constraints on the geologic history of eastern tharsis rise, mars. *Icarus*,
847 284:43–58.

848 Walter, L. S. and Salisbury, J. W. (1989). Spectral characterization of igneous rocks in the
849 8-to 12- μ m region. *Journal of Geophysical Research: Solid Earth*, 94:9203–9213.

850 Wenk, H.-R. and Wilde, W. R. (1973). Chemical anomalies of lunar plagioclase, described by
851 substitution vectors and their relation to optical and structural properties. *Contributions*
852 *to Mineralogy and Petrology*, 41:89–104.

853 Wieczorek, M. A. et al. (2022). Insight constraints on the global character of the martian
854 crust. *Journal of Geophysical Research: Planets*, 127:e2022JE007298.

855 Wordsworth, R., Kalugina, Y. N., Lokshtanov, S., Vigasin, A. A., Ehlmann, B. L., Head,
856 J. W., Sanders, C., and Wang, H. (2017). Transient reducing greenhouse warming on early
857 mars. *Geophysical Research Letters*, 44:665–671.

858 Wray, J. J., Hansen, S. T., Dufek, J., Swayze, G. A., Murchie, S. L., Seelos, F. P., Skok,
859 J. R., Irwin, R. P., and Ghiorso, M. S. (2013). Prolonged magmatic activity on mars
860 inferred from the detection of felsic rocks. *Nature Geoscience*, 6:1013–1017.

861 Zurek, R. W. and Smrekar, S. E. (2007). An overview of the mars reconnaissance orbiter
862 (mro) science mission. *Journal of Geophysical Research: Planets*, 112.




Figure 1: Top panel shows the global distribution of outcrops showing a plagioclase absorption feature in visible to near-infrared reflectance spectra. Results from this study in the circum-Argyre region are shown as yellow squares. Bottom left panel shows an example of possible layering between dark and light-toned units (HiRISE scene ESP_020799_1385). Bottom right panel shows an example light-toned massive textured outcrops with plagioclase reflectance signatures (yellow dots, mint green = alteration $\pm$ plagioclase) from the CTX global mosaic (Dickson et al., 2018).




Figure 2: Detections of high-Ca pyroxene + Olivine, type 1 and 2 olivine, plagioclase, and low-Ca pyroxene made with CRISM MRDR v4 data displayed on MOLA/HRSC shaded elevation base map (Ferguson et al., 2018). The bottom right panel shows the spatial distribution of each category as a function of radial distance from the basin center binned at 20 km intervals. The top histogram is normalized to the area of geological units from Dohm et al. (2015) investigated within each ring, and the bottom histogram is normalized by the total area of detections with each ring to elucidate relative trends in the data.




Figure 3: Reflectance and emissivity spectral character of outcrops surrounding Argyre. A) Average CRISM spectra (ratioed I/F, offset for clarity) for each category separated by low (L), moderate (M) and high (H) confidence based on absorption strength and clarity. B) Weighted absorption center (WAC) and cubic spline fit minimum (CSFM) values calculated on THEMIS-convolved reference library data (see methods for details) of felsic, intermediate, mafic, ultramafic, gabbroic, anorthositic, and pyroxenitic rocks. Anorthositic and gabbroic rocks show a trend of higher CSFM values compared to WAC values and pyroxenitic and ultramafic rocks tend to have closer to 1:1 correspondence between CSFM and WAC values. C) WAC and CSFM values calculated on ROIs detected with CRISM (THEMIS IDs: I09614002, I09639003, I17888007, I18025006) with regions from reference library spectra for interpretation. Outcrops show the highest overlap with intermediate, gabbroic, and anorthositic compositions.




Figure 4: Conceptual model for the pre-Noachian crust exposed by the Argyre impact that could explain the observed stratigraphy at both regional (10s to 100s of kilometers) and local (1–5 kilometers) spatial scales. A primitive source of possible lherzolitic or wehrlitic composition produces basaltic melt extracts that undergo fractional crystallization at lower pressures to form layered-mafic-intrusion style complexes, likely containing oxide and spinel horizons. Low-Ca pyroxene rich volcanics form a secondary upper crust.