

# Kernelized Wasserstein Natural Gradient

Michael Arbel <sup>1</sup>   Arthur Gretton <sup>1</sup>   Wuchen Li <sup>2</sup>   Guido  
Montúfar <sup>3</sup>

<sup>1</sup>Gatsby Computational Neuroscience Unit, UCL, London

<sup>2</sup>University of California, Los Angeles

<sup>3</sup>Max Planck Institute for Mathematics in the Sciences, Leipzig

November 20, 2019

# Outline

- ▶ General problem and Motivation
- ▶ Wasserstein Natural Gradient
- ▶ Kernelized Wasserstein Natural Gradient
- ▶ Experiments

## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad Z \sim \nu$$

Consider the learning problem:

## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad Z \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(Z)) \, \mathrm{d}\nu(Z)$$

## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad Z \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(\mathbf{Z})) \, \mathrm{d}\nu(Z)$$

Latent variable



## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad Z \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(Z)) \, \mathrm{d}\nu(Z)$$

Latent variable



Image

## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad (Z, Y) \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(Z), Y) \, \mathrm{d}\nu(Z, Y)$$

## Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad (Z, Y) \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(\mathbf{Z}), Y) \, \mathrm{d}\nu(Z, Y)$$



Image




# Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad (Z, Y) \sim \nu$$

Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(\mathbf{Z}), \mathbf{Y}) \, \mathrm{d}\nu(\mathbf{Z}, \mathbf{Y})$$



# Motivation: General problem

Given a model  $\rho_\theta$  of the form:

$$X \sim \rho_\theta \iff X = h_\theta(Z), \quad (Z, Y) \sim \nu$$


Consider the learning problem:

$$\min_{\theta} \mathcal{L}(\rho_\theta) := \int \ell(h_\theta(Z), Y) \, d\nu(Z, Y)$$

Image

Label

Logit



# First order methods

Large scale models  $\Rightarrow$  SGD

$$\theta_{t+1} = \theta_t - \gamma \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

# First order methods

Large scale models  $\Rightarrow$  SGD

$$\theta_{t+1} = \theta_t - \gamma \widehat{\nabla \mathcal{L}}(\rho_{\theta_t}) \leftarrow \text{Euclidean gradient}$$

# First order methods

Large scale models  $\Rightarrow$  SGD

$$\theta_{t+1} = \theta_t - \gamma \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$



- ▶ Fast training (using autodiff and GPUs)
- ▶ Often gives impressive results

# First order methods

Large scale models  $\Rightarrow$  SGD

$$\theta_{t+1} = \theta_t - \gamma \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

- ▶ Fast training (using autodiff and GPUs)
- ▶ Often gives impressive results





# First order methods: Challenges

Large scale models  $\Rightarrow$  SGD

$$\theta_{t+1} = \theta_t - \gamma \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$



- ▶ Sensitive to parametrization
- ▶ Can fail miserably when the problem is ill-conditioned



# Ill-conditioned problem

## Definition (Ill-conditioned problem)

A problem  $\min_{\theta} \mathcal{L}(\rho_{\theta})$  is ill-conditioned if the hessian  $H\mathcal{L}(\rho_{\theta})$  at a local optimum  $\theta^*$  has a high condition number:  $\kappa := \frac{\lambda_{\max}}{\lambda_{\min}}$





## Second order methods


Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \hat{\mathbf{G}}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

## Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \hat{G}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$



Euclidean gradient

# Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \widehat{\mathbf{G}}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

Preconditioner



The diagram illustrates the components of the second-order method equation. A red arrow points from the 'Preconditioner' label to the  $\widehat{\mathbf{G}}(\theta_t)^{-1}$  term in the equation. Another red arrow points from the 'Euclidean gradient' label to the  $\widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$  term.

Euclidean gradient

# Second order methods


Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \widehat{G}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

Second order information



Euclidean gradient



# Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \widehat{G}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

Second order information

- When the density of  $\rho_{\theta}$  is available ...

Euclidean gradient

## Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \widehat{G}(\theta_t)^{-1} \widehat{\nabla} \mathcal{L}(\rho_{\theta_t})$$

Second order information

- ▶ When the density of  $\rho_\theta$  is available ...
- ▶ Can choose  $\widehat{G}(\theta_t)$  as an estimator of the fisher information matrix:

Euclidean gradient

$$G_F(\theta_t) = \int \nabla \rho_{\theta_t}(x) \nabla \rho_{\theta_t}(x)^\top \rho_{\theta_t}(x) dx$$
$$\nabla^F \mathcal{L}(\rho_\theta) := G_F(\theta_t)^{-1} \nabla \mathcal{L}(\rho_{\theta_t})$$

## Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \hat{G}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

Second order information

- ▶ When the density of  $\rho_\theta$  is available ...
- ▶ Can choose  $\hat{G}(\theta_t)$  as an estimator of the fisher information matrix:

Euclidean gradient

$$G_F(\theta_t) = \int \nabla \rho_{\theta_t}(x) \nabla \rho_{\theta_t}(x)^\top \rho_{\theta_t}(x) dx$$
$$\nabla^F \mathcal{L}(\rho_\theta) := G_F(\theta_t)^{-1} \nabla \mathcal{L}(\rho_{\theta_t})$$

- ▶  $\nabla^F \mathcal{L}(\rho_\theta)$  called the *Fisher Natural gradient*.

# Second order methods

Ill-conditioned problem  $\Rightarrow$  Second order methods!!

$$\theta_{t+1} = \theta_t - \gamma \hat{G}(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$$

Second order information

- ▶ When the density of  $\rho_\theta$  is available ...

- ▶ Can choose  $\hat{G}(\theta_t)$  as an estimator of the fisher information matrix:



$$G_F(\theta_t) = \int \nabla \rho_{\theta_t}(x) \nabla \rho_{\theta_t}(x)^\top \rho_{\theta_t}(x) dx$$
$$\nabla^F \mathcal{L}(\rho_\theta) := G_F(\theta_t)^{-1} \nabla \mathcal{L}(\rho_{\theta_t})$$

- ▶  $\nabla^F \mathcal{L}(\rho_\theta)$  called the *Fisher Natural gradient*.
- ▶ Robust to parametrization.



Euclidean gradient



## Second order methods: Robustness to parametrization



## Second order methods: Robustness to parametrization



## Second order methods: Robustness to parametrization

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \nabla^F \mathcal{L}(\rho_{\theta_t})$

- Have a change of variables  $\psi = \Psi(\theta)$  and write  $\tilde{\rho}_\psi = \rho_\theta$ .

## Second order methods: Robustness to parametrization

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \nabla^F \mathcal{L}(\rho_{\theta_t})$

- ▶ Have a change of variables  $\psi = \Psi(\theta)$  and write  $\tilde{\rho}_\psi = \rho_\theta$ .
- ▶ Continuous-time limit of the Fisher natural gradient:

$$\begin{aligned}\dot{\theta}_t &= -\nabla^F \mathcal{L}(\rho_{\theta_t}), & \theta_0 \\ \dot{\psi}_t &= -\nabla^F \mathcal{L}(\tilde{\rho}_{\psi_t}), & \psi_0 = \Psi(\theta_0)\end{aligned}$$

## Second order methods: Robustness to parametrization

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \nabla^F \mathcal{L}(\rho_{\theta_t})$

- ▶ Have a change of variables  $\psi = \Psi(\theta)$  and write  $\tilde{\rho}_\psi = \rho_\theta$ .
- ▶ Continuous-time limit of the Fisher natural gradient:

$$\begin{aligned}\dot{\theta}_t &= -\nabla^F \mathcal{L}(\rho_{\theta_t}), & \theta_0 \\ \dot{\psi}_t &= -\nabla^F \mathcal{L}(\tilde{\rho}_{\psi_t}), & \psi_0 = \Psi(\theta_0)\end{aligned}$$

- ▶ Robustness to parametrization  $\Rightarrow \psi_t = \Psi(\theta_t)$

## Second order methods: Robustness to parametrization

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \nabla^F \mathcal{L}(\rho_{\theta_t})$

- ▶ Have a change of variables  $\psi = \Psi(\theta)$  and write  $\tilde{\rho}_\psi = \rho_\theta$ .
- ▶ Continuous-time limit of the Fisher natural gradient:

$$\begin{aligned}\dot{\theta}_t &= -\nabla^F \mathcal{L}(\rho_{\theta_t}), & \theta_0 \\ \dot{\psi}_t &= -\nabla^F \mathcal{L}(\tilde{\rho}_{\psi_t}), & \psi_0 = \Psi(\theta_0)\end{aligned}$$

- ▶ Robustness to parametrization  $\Rightarrow \psi_t = \Psi(\theta_t)$
- ▶ Doesn't hold for the euclidean gradient in general!

## Second order methods: Challenges

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \hat{\mathbf{G}}_F(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$

---

<sup>1</sup>[Grosse and Martens, 2016, George et al., 2018]

## Second order methods: Challenges

Fisher Natural gradient descent:  $\theta_{t+1} = \theta_t - \gamma \hat{\mathbf{G}}_F(\theta_t)^{-1} \widehat{\nabla \mathcal{L}}(\rho_{\theta_t})$

- ▶ Computational: Expensive to store and invert  $\hat{\mathbf{G}}_F(\theta)$  at every iteration .
- ▶ Prior works proposed cheap approximations of  $\hat{\mathbf{G}}_F(\theta)$  <sup>1</sup>.
- ▶ Requires to know the density  $\rho_\theta$ .

---

<sup>1</sup>[Grosse and Martens, 2016, George et al., 2018]



# Contributions

A second order method based on the Wasserstein natural gradient which is:

- ▶ Robust to parametrization
- ▶ Doesn't require access to the density of the model
- ▶ Trades-off between accuracy and computational cost
- ▶ Comes with convergence rates.

# General recipe for natural gradients

1. Choose a distance/divergence  $d$  defined on the model  $\rho_\theta$ :

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + d(\rho_{\theta_t+u}, \rho_{\theta_t})$$

2. Second order expansion of  $d$  at  $\rho_\theta$ :

$$d(\rho_{\theta_t+u}, \rho_{\theta_t}) \simeq \frac{1}{2} u^\top G(\theta_t) u$$

3. Solve:

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2} u^\top G(\theta_t) u$$

# Recipe for Fisher natural gradients

1. Choose the  $KL$  as a divergence

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + KL(\rho_{\theta_t+u} | \rho_{\theta_t})$$

2. Second order expansion of  $KL$  at  $\rho_{\theta}$ :

$$KL(\rho_{\theta_t+u}, \rho_{\theta_t}) \simeq \frac{1}{2} u^\top G_F(\theta_t) u$$

3. Solve:

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2} u^\top G_F(\theta_t) u$$

# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- Well defined even when  $\rho_\theta$  doesn't admit a density

# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- Well defined even when  $\rho_\theta$  doesn't admit a density

$$W_2^2(\delta_{\theta_1}, \delta_{\theta_2}) = \|\theta_1 - \theta_2\|^2$$

# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$


- $W_2$  has a nice geometric interpretation:

# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:




# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:






# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:




# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:




# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:




# Wasserstein Natural Gradient: Step 1

1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:



# Wasserstein Natural Gradient: Step 1


1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:

**Continuity equation:**  $\partial_t \nu_t + \text{div}(\nu_t \phi_t) = 0$

**Boundary conditions:**  $\nu_0 = p, \quad \nu_1 = q$



# Wasserstein Natural Gradient: Step 1


1. Choose  $d(\rho_\theta, \rho_{\theta+u}) = \frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2\lambda} W_2^2(\rho_{\theta_t+u}, \rho_{\theta_t})$$

- $W_2$  has a nice geometric interpretation:

**Continuity equation:**  $\partial_t \nu_t + \text{div}(\nu_t \phi_t) = 0$

**Boundary conditions:**  $\nu_0 = p, \quad \nu_1 = q$



Benamou-Brenier formula<sup>2</sup>:  $W_2^2(p, q) := \inf_{(\nu_t, \phi_t)} \int_0^1 \int \|\phi_t(x)\|^2 d\nu_t(x)$

---

<sup>2</sup>[Benamou and Brenier, 2000]

# General recipe for natural gradients

1. Choose a distance/divergence  $d$  defined on the model  $\rho_\theta$ :

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + d(\rho_{\theta_t+u}, \rho_{\theta_t})$$

2. Second order expansion of  $d$  at  $\rho_\theta$ :

$$d(\rho_{\theta_t+u}, \rho_{\theta_t}) \simeq \frac{1}{2} u^\top G(\theta_t) u$$

3. Solve:

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2} u^\top G(\theta_t) u$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$




# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$


**Continuity equation:**  $\partial_t \nu_t + \text{div}(\nu_t \phi_t) = 0$

**Boundary conditions:**  $\nu_0 = p, \quad \nu_1 = q$




# Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$




# Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$




# Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$




# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$




# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$



# Wasserstein Natural Gradient: Step 2


## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$



# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

**Elliptic equation:**  $\nabla \rho_\theta^\top u + \text{div}(\rho_\theta \phi) = 0$






# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

**Elliptic equation:**  $\nabla \rho_\theta^\top u + \text{div}(\rho_\theta \phi) = 0$




$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \inf_{\phi} \int \|\phi(x)\|^2 d\rho_\theta(x)$$

# Wasserstein Natural Gradient: Step 2

## 2. Taylor expansion of $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

**Elliptic equation:**  $\nabla \rho_\theta^\top u + \text{div}(\rho_\theta \phi) = 0$



$$W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \int \|\phi(x)\|^2 d\rho_\theta(x)$$

$\phi$  constrained to be 'almost' a gradient of a real valued function.

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla \rho_\theta^\top u + \text{div}(\rho_\theta \phi) = 0$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla_{\rho_\theta}^\top u + \text{div}(\rho_\theta \phi) = 0$$

What does  $\nabla_{\rho_\theta}$  mean?

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla \rho_\theta^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$

What does  $\nabla \rho_\theta$  mean?

$$\text{Implicit model} \Rightarrow \mathbb{E}_{\rho_\theta}[f(x)] = \int f(h_\theta(z)) d\nu(z)$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla \rho_\theta^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$

What does  $\nabla \rho_\theta$  mean?

$$\text{Implicit model} \Rightarrow \mathbb{E}_{\rho_\theta}[f(x)] = \int f(h_\theta(z)) d\nu(z)$$

Taylor expansion in  $u$ :

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla \rho_\theta^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$

What does  $\nabla \rho_\theta$  mean?

$$\text{Implicit model} \Rightarrow \mathbb{E}_{\rho_\theta}[f(x)] = \int f(h_\theta(z)) d\nu(z)$$

Taylor expansion in  $u$ :

$$\mathbb{E}_{\rho_{\theta+u}}[f(x)] - \mathbb{E}_{\rho_\theta}[f(x)] \simeq \int \nabla_x f(h_\theta(z)) \nabla h_\theta(z)^\top u d\nu(z)$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x) \\ \nabla_{\rho_\theta}^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$

What does  $\nabla_{\rho_\theta}$  mean?

$$\text{Implicit model} \Rightarrow \mathbb{E}_{\rho_\theta}[f(x)] = \int f(h_\theta(z)) d\nu(z)$$

Taylor expansion in  $u$ :

$$\mathbb{E}_{\rho_{\theta+u}}[f(x)] - \mathbb{E}_{\rho_\theta}[f(x)] \simeq \underbrace{\int \nabla_x f(h_\theta(z)) \nabla h_\theta(z)^\top u d\nu(z)}_{\nabla_{\rho_\theta}(f)^\top u}$$



## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x)$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \frac{1}{2} \int \|\phi(x)\|^2 d\rho_\theta(x)$$

Variational expression for elliptic equations:

$$\nabla_{\rho_\theta}^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$



$$\phi \in \arg \sup_{f \in C_c^\infty(\Omega)} \nabla_{\rho_\theta}(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta}[\|\nabla f(x)\|^2]$$

## Wasserstein Natural Gradient: Step 2

2. Taylor expansion of  $\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u})$

$$\frac{1}{2} W_2^2(\rho_\theta, \rho_{\theta+u}) \simeq \sup_{f \in C_c^\infty(\Omega)} \nabla \rho_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

Variational expression for elliptic equations:

$$\nabla \rho_\theta^\top u + \operatorname{div}(\rho_\theta \phi) = 0$$



$$\phi \in \arg \sup_{f \in C_c^\infty(\Omega)} \nabla \rho_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla f(x)\|^2]$$

# General recipe for natural gradients

1. Choose a distance/divergence  $d$  defined on the model  $\rho_\theta$ :

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + d(\rho_{\theta_t+u}, \rho_{\theta_t})$$

2. Second order expansion of  $d$  at  $\rho_\theta$ :

$$d(\rho_{\theta_t+u}, \rho_{\theta_t}) \simeq \frac{1}{2} u^\top G(\theta_t) u$$

3. Solve:

$$\min_u \nabla \mathcal{L}(\rho_{\theta_t})^\top u + \frac{1}{2} u^\top G(\theta_t) u$$

## Wasserstein Natural Gradient: Step 3

3. Solve:

$$\min_u \nabla \mathcal{L}(\rho_\theta)^\top u + \sup_{f \in C_c^\infty(\Omega)} \nabla \rho_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

## Wasserstein Natural Gradient: Step 3

3. Solve:

$$\min_u \sup_{f \in C_c^\infty(\Omega)} (\nabla \mathcal{L}(\rho_\theta) + \nabla \rho_\theta(f))^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

## Wasserstein Natural Gradient: Step 3

3. Solve:

$$\min_u \sup_{f \in C_c^\infty(\Omega)} \overbrace{(\nabla \mathcal{L}(\rho_\theta) + \nabla \rho_\theta(f))}^{\mathcal{U}_\theta(f)}^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

## Wasserstein Natural Gradient: Step 3

3. Solve:

$$\min_u \sup_{f \in C_c^\infty(\Omega)} \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$



# Kernelized Wasserstein Natural Gradient

$$\min_u \sup_{f \in C_c^\infty(\Omega)} \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

# Kernelized Wasserstein Natural Gradient

$$\min_u \sup_{f \in C_c^\infty(\Omega)} \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

- Functional optimization: hard in general

# Kernelized Wasserstein Natural Gradient

$$\min_u \sup_{f \in \mathcal{H}} \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2]$$

- ▶ Functional optimization: hard in general
- ▶ Replace  $C_c^\infty(\Omega)$  by a nicer space: an RKHS  $\mathcal{H}^3$ .

# Kernelized Wasserstein Natural Gradient

$$\min_u \sup_{f \in \mathcal{H}} \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2] + \frac{1}{2} (\epsilon \|u\|^2 - \lambda \|f\|_{\mathcal{H}}^2)$$

- ▶ Functional optimization: hard in general
- ▶ Replace  $C_c^\infty(\Omega)$  by a nicer space: an RKHS  $\mathcal{H}$ .
- ▶ Regularization terms  $\Rightarrow$  Ensure strong convexity/concavity

# Kernelized Wasserstein Natural Gradient

$$\sup_{f \in \mathcal{H}} \min_u \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2] + \frac{1}{2} (\epsilon \|u\|^2 - \lambda \|f\|_{\mathcal{H}}^2)$$

- ▶ Functional optimization: hard in general
- ▶ Replace  $C_c^\infty(\Omega)$  by an RKHS  $\mathcal{H}$  with kernel  $k^4$ .
- ▶ Regularization terms  $\Rightarrow$  Ensure strong convexity/concavity
- ▶ Minimax Theorem<sup>5</sup>  $\Rightarrow$  Exchange order of min and sup.

---

<sup>4</sup>[Mroueh et al., 2019]

<sup>5</sup>[Ekeland and Témam, 1999]

# Kernelized Wasserstein Natural Gradient

$$\sup_{f \in \mathcal{H}} \min_u \mathcal{U}_\theta(f)^\top u - \frac{1}{2} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2] + \frac{1}{2} (\epsilon \|u\|^2 - \lambda \|f\|_{\mathcal{H}}^2)$$

- ▶ Functional optimization: hard in general
- ▶ Replace  $C_c^\infty(\Omega)$  by an RKHS  $\mathcal{H}$  with kernel  $k^4$ .
- ▶ Regularization terms  $\Rightarrow$  Ensure strong convexity/concavity
- ▶ Minimax Theorem<sup>5</sup>  $\Rightarrow$  Exchange order of min and sup.
- ▶ Optimal descent direction  $u^*$  given by:

$$u^* = -\frac{1}{\epsilon} \mathcal{U}_\theta(f^*)$$

$$f^* = \arg \min_{f \in \mathcal{H}} \mathbb{E}_{\rho_\theta} [\|\nabla_x f(x)\|^2] + \frac{1}{\epsilon} \|\mathcal{U}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

---

<sup>4</sup>[Mroueh et al., 2019]

<sup>5</sup>[Ekeland and T  man, 1999]

## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

► Representer theorem  $\Rightarrow$  Optimal solution  $\hat{f}^*$  of the form:

$$\hat{f}^* = \sum_{n=1}^N \sum_{i=1}^d \beta_{n,i} \partial_i k(X_n, \cdot)$$



## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

- Representer theorem  $\Rightarrow$  Optimal solution  $\hat{f}^*$  of the form:

$$\hat{f}^* = \sum_{n=1}^N \sum_{i=1}^d \beta_{n,i} \partial_i k(X_n, \cdot)$$

- $\beta_{n,i}$  obtained by solving a linear system of size  $Nd$ !!

## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

- Nystrom projections <sup>6</sup>  $\Rightarrow$  Reduce computational cost:

$$\hat{f}_M^* = \sum_{m=1}^M \alpha_m \partial_{i_m} k(\tilde{X}_m, \cdot)$$

---

<sup>6</sup>[Rudi et al., 2015, Sutherland et al., 2017]

## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

- Nystrom projections <sup>6</sup>  $\Rightarrow$  Reduce computational cost:

$$\hat{f}_M^* = \sum_{m=1}^M \alpha_m \partial_{i_m} k(\tilde{X}_m, \cdot)$$

$M$  sub-samples from  $(X_i)_{1 \leq i \leq N}$

---

<sup>6</sup>[Rudi et al., 2015, Sutherland et al., 2017]

## KWNG: Sample based version

Have some i.i.d. samples  $(Z_n)_{1 \leq n \leq N}$  from  $\nu$  and  $X_n = h_\theta(Z_n)$ :

$$\hat{u}^* = -\frac{1}{\epsilon} \hat{\mathcal{U}}_\theta(\hat{f}^*),$$

$$\hat{f}^* = \arg \min_{f \in \mathcal{H}} \frac{1}{N} \sum_{n=1}^N \|\nabla_x f(X_n)\|^2 + \frac{1}{\epsilon} \|\hat{\mathcal{U}}_\theta(f)\|^2 + \lambda \|f\|_{\mathcal{H}}^2$$

- Nystrom projections <sup>6</sup>  $\Rightarrow$  Reduce computational cost:

$$\hat{f}_M^* = \sum_{m=1}^M \alpha_m \partial_{i_m} k(\tilde{X}_m, \cdot)$$

Randomly sampled from  $\{1, \dots, d\}$

$M$  sub-samples from  $(X_i)_{1 \leq i \leq N}$

---

<sup>6</sup>[Rudi et al., 2015, Sutherland et al., 2017]

## KWNG: Sample based version


After some further calculations:

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (TT^\top + \lambda \epsilon K + \epsilon CC^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

## KWNG: Sample based version

After some further calculations:

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (TT^\top + \lambda \epsilon K + \epsilon \mathbf{C} \mathbf{C}^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$


$$C_{m,(n,i)} = \frac{1}{\sqrt{N}} \partial_{i_m} \partial_{i+d} k(\tilde{X}_m, X_n)$$

# KWNG: Sample based version

After some further calculations:

$$K_{m,m'} = \partial_{i_m} \partial_{i_{m'}+d} k(\tilde{X}_m, \tilde{X}_{m'})$$

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (T T^\top + \lambda \epsilon K + \epsilon C C^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$C_{m,(n,i)} = \frac{1}{\sqrt{N}} \partial_{i_m} \partial_{i+d} k(\tilde{X}_m, X_n)$$

# KWNG: Sample based version

After some further calculations:

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - \boxed{T^\top} (TT^\top + \lambda \epsilon \boxed{K} + \epsilon \boxed{C} C^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$K_{m,m'} = \partial_{i_m} \partial_{i_{m'}+d} k(\tilde{X}_m, \tilde{X}_{m'})$

$C_{m,(n,i)} = \frac{1}{\sqrt{N}} \partial_{i_m} \partial_{i+d} k(\tilde{X}_m, X_n)$

$T := \nabla \tau(\theta)$  with  $\tau(\theta)_m = \frac{1}{N} \sum_{n=1}^N \partial_{i_m} k(\tilde{X}_m, h_\theta(Z_n))$



# Theory: Consistency and convergence rates

## Theorem

Let  $\delta$  be such that  $0 \leq \delta \leq 1$ . Under smoothness assumptions on the model characterized by some constant  $c \geq 0$ , for  $N$  large enough,  $M \sim (dN^{\frac{2+c}{4+c}} \log(N))$ ,  $\lambda \sim N^{\frac{1}{2b+1}}$  and  $\epsilon \lesssim N^{-\frac{1}{4+c}}$ , it holds with probability at least  $1 - \delta$  that:

$$\|\widehat{\nabla^W \mathcal{L}(\theta)} - \nabla^W \mathcal{L}(\theta)\|^2 = \mathcal{O}\left(N^{-\frac{2}{4+c}}\right).$$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (TT^\top + \epsilon CC^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (TT^\top + \epsilon CC^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$T_m = \frac{1}{N} \sum_{n=1}^N \nabla_{\theta} \partial_{i_m} k(Y_m, h_{\theta}(Z_n))$$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - T^\top (TT^\top + \epsilon CC^\top)^\dagger T \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$T = CB, \quad B_n = \nabla_\theta h_\theta(Z_n)$$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - B^\top C^\top (CBB^\top C^\top + \epsilon CC^\top)^\dagger CB \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$T = CB, \quad B_n = \nabla_\theta h_\theta(Z_n)$$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^W \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - B^\top C^\top (C B B^\top C^\top + \epsilon C C^\top)^\dagger C B \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$T = C B, \quad B_n = \nabla_\theta h_\theta(Z_n)$$

'Simplify'  $C$ :

$$\tilde{T} = S^\dagger U^\top T, \quad P = S^\dagger S$$

where  $C C^\top = U S U^\top$

## KWNG: Ridgeless version

Additional structure when  $\lambda = 0$ :

$$\widehat{\nabla^w \mathcal{L}(\theta)} = \frac{1}{\epsilon} \left( I - \tilde{T}^\top (\tilde{T} \tilde{T}^\top + \epsilon P)^\dagger \tilde{T} \right) \widehat{\nabla \mathcal{L}(\theta)}$$

$$T = CB, \quad B_n = \nabla_\theta h_\theta(Z_n)$$


'Simplify'  $C$ :

$$\tilde{T} = S^\dagger U^\top T, \quad P = S^\dagger S$$

where  $CC^\top = USU^\top$

# Experimental evaluation: Synthetic models


Hyper-spheres:  $X = a + rZ$ ,  $Z \sim \mathbb{S}_d$





# Experimental evaluation: Synthetic models

Gaussians:  $X = a + rZ$ ,  $Z \sim \mathcal{N}(0, I)$




## Experimental evaluation: Sensitivity to the choice of the kernel

- ▶ Gaussian kernel  $k(x, y) = \exp(-\frac{\|x-y\|^2}{\sigma})$

# Experimental evaluation: Sensitivity to the choice of the kernel

- Gaussian kernel  $k(x, y) = \exp(-\frac{\|x-y\|^2}{\sigma})$





## Experimental evaluation: Optimization trajectory

- ▶ Gaussian model for  $\rho_\theta$
- ▶ Loss functional  $\mathcal{L}(\rho_\theta) = W_2^2(\rho_\theta, \rho_{\theta^*})$ .

# Experimental evaluation: Optimization trajectory

- ▶ Gaussian model for  $\rho_\theta$
- ▶ Loss functional  $\mathcal{L}(\rho_\theta) = W_2^2(\rho_\theta, \rho_{\theta^*})$ .



# Experimental evaluation: Classification task



Well-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) \, \mathrm{d}\nu(Z, Y)$$

# Experimental evaluation: Classification task

Well-conditioned problem:



$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$



# Experimental evaluation: Classification task

Well-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$







# Experimental evaluation: Classification task

Well-conditioned problem:



$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$



# Experimental evaluation: Classification task

Well-conditioned problem:



$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$



# Experimental evaluation: Classification task

Well-conditioned problem:



$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$



# Experimental evaluation: Classification task

Well-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(h_{\theta}(Z), Y) d\nu(Z, Y)$$



# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) \, \mathrm{d}\nu(Z, Y)$$


$U$  is a diagonal matrix with  $\kappa = 10^7$

# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$




# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$




# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$







# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$




# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$




# Experimental evaluation: Classification task

Ill-conditioned problem:

$$\min_{\theta} \mathcal{L}(\rho_{\theta}) := \int \ell(Uh_{\theta}(Z), Y) d\nu(Z, Y)$$

$U$  is a diagonal matrix with  $\kappa = 10^7$



# Conclusion

## Summary of contributions

- ▶ Proposed to use the Wasserstein natural gradient for ill-conditioned problems
- ▶ A new algorithm to estimate the Wasserstein natural gradient
- ▶ Convergence rate: trade-off between computational complexity and statistical accuracy

## Future work:

- ▶ Consistency result for the ridgeless version <sup>7</sup>
- ▶ Potential application in RL ( implicit policy for RL <sup>8</sup> )

---


<sup>7</sup>[Liang et al., 2017]

<sup>8</sup>[Tang and Agrawal, 2019]

Thank you !

# Ablation study

- ▶ Choice of the damping matrix  $D(\theta)$
- ▶ Choice of the kernel (gaussian vs rational quadratic)





Benamou, J.-D. and Brenier, Y. (2000).

A computational fluid mechanics solution to the monge-kantorovich mass transfer problem.

*Numerische Mathematik*, 84(3):375–393.



Ekeland, I. and Téman, R. (1999).

*Convex Analysis and Variational Problems*.

Society for Industrial and Applied Mathematics,  
Philadelphia, PA, USA.



George, T., Laurent, C., Bouthillier, X., Ballas, N., and Vincent, P. (2018).

Fast Approximate Natural Gradient Descent in a  
Kronecker-factored Eigenbasis.

*arXiv:1806.03884 [cs, stat]*.

arXiv: 1806.03884.



Grosse, R. and Martens, J. (2016).

A Kronecker-factored Approximate Fisher Matrix for  
Convolution Layers.

*In Proceedings of the 33rd International Conference on*