

Natural Land Cover Improves the COVID-19 Health Outcomes

Abstract:

Coronavirus disease 2019 (COVID-19) poses special challenges for societies, as the disease causes millions of deaths. Although the direct prevention measures affect the prevalence and mortality the most, the other indirect factors, including natural environments and economics, could not be neglected. Assessing whether natural land cover impacts COVID-19 health outcomes is an urgent and crucial public health topic. Here, we examine the relationships between natural land cover and the prevalence and mortality of COVID-19 in the United States. A 1% increase of open water or deciduous forest is associated with a 0.004-death and 0.163-confirmed-case, or 0.006-death and 0.099-confirmed-case decrease in every 1,000 people. Converting them into monetary value, for the mortality, a 1% increase in open water, deciduous forest, or evergreen forest in a county is equivalent to a 212-, 313-, or 219-USD increase in household income in the long term. Moreover, for the prevalence, a 1% change in open water, deciduous forest, or mixed forest is worth a 382-, 230-, or 650-USD increase in household income. Furthermore, a rational development intensity is also critical to prevent the COVID-19 pandemic. More greenery in the short term could also slash the prevalence and mortality. Our research highlights that societies could prevent other pandemics similar to the COVID-19 and improve public health in the future by adding natural land cover.

Keywords:

COVID-19; Prevalence; Mortality; Natural Land Cover; NDVI

Introduction

Coronavirus disease 2019 (COVID-19) has raised serious and urgent concerns globally (Chen et al., 2020; Verity et al., 2020). As of Nov. 1st, 2021, there were almost 246.69 million confirmed cases and 5.00 million deaths due to COVID-19 worldwide (Data from WHO COVID-19 Dashboard, see <https://covid19.who.int/>). In the United States (U.S.), the cumulative numbers of confirmed and death cases owing to COVID-19 are 45,665,006 and 741,650, respectively, as of Nov. 1st, 2021 (Data from U.S. Centers for Disease Control and Prevention, CDC). Moreover, the county-level COVID-19 health outcomes, including the mortality and prevalence, vary dramatically in the United States, ranging from 0 to 10.77 deaths/1,000 capita and from 19.62 to 543.05 cases/1,000 capita. A high prevalence means that more residuals get infected by the COVID-19, and a high mortality indicates that more people die due to the COVID-19. The local population with the lower prevalence and mortality may have few other chronic diseases (Mollalo et al., 2021; Nishiga et al., 2020) and relatively good mental health (Duan and Zhu, 2020; Dzhambov et al., 2020). Therefore, from a positive perspective, the COVID-19 is a cruel and dangerous filter that can identify healthy and unhealthy populations and helps people to detect more factors affecting public health.

Investigations regarding the relationships between COVID-19 health outcomes and geographical factors are urgently needed to locate the high-risk areas, to slow the disease's devastation, and to slash the risk of similar infectious diseases outbreaks (Mollalo et al., 2020a; Mollalo et al., 2020b; Yoo and Managi, 2020). Considering the current situation, globally, eliminating the COVID-19 is impossible in a short time (Lee et al., 2020). Thus, how to reduce the negative impacts and the deaths is a critical focus. Previous studies indicate that a healthy immune system is the main reason for the

patients surviving from the coronavirus (Combes et al., 2021; Williamson et al., 2020). Chronic diseases, such as cardiovascular disease, are associated with an increased risk of death in COVID-19 patients (Nishiga et al., 2020). Besides, people experience less depression and anxiety, exposed to more green space during the COVID-19 pandemic (Dzhambov et al., 2020). Natural environments, mainly based on natural land cover, provide ecosystem services to improve physical and mental health (Bratman et al., 2019; Costanza et al., 2014; Hartig and Kahn, 2016; Li and Managi, 2021b). Hence, the analyses on the associations of natural land cover with the COVID-19 outcomes may help identify the high-death-risk areas in the COVID-19 and develop optimal land-use policies to deal with other similar public health emergencies in the future (Nakamura and Managi, 2020).

Natural environments are positively related to public health (Rook, 2013; Sandifer et al., 2015; Ugolini et al., 2020). The severity of the COVID-19 symptoms might be correlated with people's living environments. People living with less greenness have more medical conditions (Williamson et al., 2020), like cardiovascular disease (Maas et al., 2009; Mitchell and Popham, 2008; Rahmani and Mirmahaleh, 2021), which would ultimately exacerbate the symptom of COVID-19 (Nishiga et al., 2020). Numerous researchers point out that the natural land cover in the local communities is associated with health outcomes by promoting physical exercise and social connections, relieving stress, and removing air pollution, noise, and heat exposure (Alcock et al., 2015; James et al., 2015; Li and Managi, 2021a; Richardson et al., 2013). It is hypothesized that living with more natural land cover reduces the prevalence and mortality by alleviating the COVID-19 symptoms. In other words, an increase in exposure to green spaces is associated with decreased risks of clinical diseases. Furthermore, living with more greenery is linked with improved mental health during

the pandemic (Dzhambov et al., 2020). To conclude, natural environments might improve health and ultimately reach a lower COVID-19 prevalence and mortality.

There is a trade-off between the health benefits and economic costs of an increase in natural land cover. In this way, an evaluation of the value of natural land cover on the COVID-19 is necessary. To estimate the monetary valuation of the environmental goods, stated preferences and revealed preference methods are widely used (Frey et al., 2010; Tsurumi and Managi, 2017). Stated preferences methods need surveys to directly ask the respondents to evaluate the monetary values of environmental goods, which is obviously inconsistent with the current topic. However, the revealed preference methods are more straightforward and do not need surveys. These methods only need to investigate the relationships among variables and then utilize the estimated coefficient to calculate the marginal substitution rate between environmental goods and income, based on the micro-econometric public health functions (Alcock et al., 2015; Frey et al., 2010; Li and Managi, 2021c). In other words, these methods assume that an alteration in income is critical to compensate for the change in environmental goods, vice versa.

Materials and Methods

Materials

Health Outcomes of COVID-19

Two variables, county-level prevalence and mortality, are used as the proxies for the COVID-19 health outcomes. The county-level prevalence is the ratio of the confirmed cases to the total population in a certain county over a specific period, and the county-level mortality is the ratio of the deaths to the total population (Porta, 2014).

The unit of these two indicators is cases per 1,000 capita (*cases/1,000 cap*). The accumulated numbers of the confirmed cases and deaths, and population are from the CDC. To detect the impacts of long-term living with nature on the COVID-19 health outcomes, the total prevalence and the total mortality are employed, based on the accumulated numbers from the first confirmed case recorded to Nov. 1st, 2021. **Figure 1** illustrates the spatial distribution of the total prevalence and mortality. Additionally, we calculated the quarterly prevalence and the quarterly mortality from the first quarter of 2020 to the third quarter of 2021 to examine the short-term effects of greenery. Due to some events, such as the Election, in certain months in the U.S., a tremendous monthly variation exists. To grasp the actual impacts of the environment, we therefore use the more stable data set, which is quarterly. It must be noted that there are reductions of the accumulated numbers of the confirmed case and deaths in several counties on some days, which might be caused by misdiagnosis, duplicate recorded by different counties, or other reasons. In the quarterly data set with more than 20,000 records, there are only no more than 50 reductions. Therefore, to avoid the negative prevalence and mortality, we force to set those reductions zero. (**Table A1**: Cross-Sectional Data Statistic Summary, **Table A2**: Panel Data Statistic Summary, **Table A3**: Data Source)

Land Cover Data

We extract county-level land cover data from the National Land Cover Dataset (NLCD). In the NLCD archive, there are eight-year data set regarding the CONUS, including 2001, 2004, 2006, 2008, 2011, 2013, 2016, and 2019. These data sets include 20 land types, but there are four land types, namely, dwarf scrub, sedge, lichens, and moss, only in Alaska. In other words, in the Contiguous United States (CONUS), there are 16 other land types: open water, perennial ice, developed open space, low intensity

developed area, medium intensity developed area, high intensity developed area, barren land, deciduous forest, evergreen forest, mixed forest, shrub, grassland, pasture, cultivated crops, woody wetlands, and emergent herbaceous wetlands (Detailed classification description, see <https://www.mrlc.gov/data/legends/national-land-cover-database-2019-nlcd2019-legend>). In this study, high intensity developed area and medium intensity developed area are considered as urban centers and urban areas, respectively. The difference among the four types of developed areas is the proportion of impervious surfaces in every grid. High intensity developed area has over 80% impervious surface and less than 20% greenery or water. Medium intensity developed area, low intensity developed area, and developed open space have 50% – 80%, 20% – 50%, less than 20% impervious surface, respectively.

The average percentages of each land type in the counties from the eight-year data set are taken as the land cover data in the analyses. At first, the total areas of each land type in the counties are obtained by tool in ArcGIS Pro 2.5.0, Tabulate Area, using the boundary shapefile from the U.S. Census Bureau. Then, they are converted into percentages, and every county in the CONUS has eight-year values. Finally, we average the values of each county. To probe the impacts of natural land cover on the COVID-19 health outcome and to get around the multicollinearity in the analyses, land cover variables, including open water, developed open space, low intensity developed area, medium intensity developed area, high intensity developed area, deciduous forest, evergreen forest, mixed forest, shrub, grassland, woody wetlands, and emergent herbaceous wetlands, are put into the cross-sectional regressions. (**Table A1: Cross-Sectional Data Statistic Summary**)

Normalized Difference Vegetation Index (NDVI) Data

To examine the short-term impact of the natural environment, we use the monthly NDVI data produced by the U.S. National Aeronautics and Space Administration (NASA). The NDVI is a graphical index to describe whether the observed pixel contains live green vegetation. This index range from -1 (no live green vegetation, -100%) to 1 (rife with live green vegetation, 100%). Although the panel land cover data set is desired in our study, the high-resolution raster is created every two years or longer, and there is a delay. Moreover, we also try to obtain the land cover data provided by NASA. However, NASA's land cover products are yearly low-resolution, and 2021's is not available. For these reasons, we eventually take the NDVI data set as the natural land cover variable in the panel regressions. We extract the monthly NDVI value based on NASA's products, MOD13A3 (<https://lpdaac.usgs.gov/products/mod13a3v006/>) and MYD13A3 (<https://lpdaac.usgs.gov/products/myd13a3v006/>), with a 1-km resolution. Then, the quarterly average values of each county are calculated. (**Table A2:** Panel Data Statistic Summary)

Other Potential Confounders

Twenty-eight other county-level potential confounders are obtained and controlled in the cross-sectional regressions. They are divided into five classes: political, demographic, socio-economic, clinical, and meteorological aspects. First, the political aspect includes five variables: the days of gatherings restrictions, the days of transport closing, the days of staying home restrictions, the days of international movement restrictions (international MoRe), and the days of internal movement restrictions (internal MoRe), from the first confirmed case recorded to Nov. 1st, 2021. All the political aspect confounders are acquired from the R package "COVID19" (Guidotti

and Ardia, 2020), based on the Oxford COVID-19 Government Response Tracker. Secondly, the demographic aspect includes six variables: the percentages of the population within specific age ranges, the percentage of black people, the percentage of Hispanic people, and the percentage of males. The U.S. Census Bureau provides the demographic data. Thirdly, the socio-economic aspect contains four variables: the unemployment rate, the median household income of counties, the poverty rate, and the percentage of the population without a high school diploma. These variables are obtained from the U.S. Department of Agriculture. Fourthly, there are eight variables in clinical aspect: poor health rate in 2019, the average days of poor physical health in 2019, the average days of poor mental health in 2019, smoker rate in 2019, the obesity rate in 2019, physical inactivity rate in 2019, exercise opportunity rates in 2019, and the numbers of hospital beds. All of these data are acquired from the University of Wisconsin, School of Medicine and Public Health. Finally, the meteorological aspect contains five variables: the mean of PM_{2.5} value during 2000-2016, and means of daily temperature and relative humidity in summer (June to September) and winter (December to February) during 2000-2016. The U.S. Environmental Protection Agency provides PM_{2.5} values, and other meteorological data are downloaded and extracted from Google Climatology Lab. (**Table A1:** Cross-Sectional Data Statistic Summary)

Three other variables, including surface temperature, nighttime light (NTL) index, and restriction stringency score, are controlled in the panel regression to detect the relationship between the county-level NDVI and the quarterly mortality and prevalence from the first quarter in 2020 to the third quarter in 2021. The surface temperature is an average value of the monthly day-time and nighttime surface temperature extracted from another NASA's products, MOD11C2 (<https://lpdaac.usgs.gov/products/mod11c2v006/>) and MOD11C2

(<https://lpdaac.usgs.gov/products/myd11c2v006/>) with a 0.05-arc-degree resolution. The NTL data are extracted from NASA's products, VNP46A3 (<https://ladsweb.modaps.eosdis.nasa.gov/missions-and-measurements/products/VNP46A3/>). The NTL data are widely used to represent the economic status of a specific region, according to the assumption that the brighter places are generally more affluent. Because the quarterly county-level economic status and prosperity are difficult to acquire, we take the NTL index as the substitution. The restriction stringency scores of each country are obtained from the Oxford COVID-19 Government Response Tracker, which is calculated based on the restriction policies, including gathering restrictions, transport closing policies, staying home policies, internal MoRe, and international MoRe. (**Table A2**: Panel Data Statistic Summary, **Table A3**: Data Source)

Methods

Ordinary Least Square (OLS) Model

According to several statistical tests, the Spatial Simultaneous Autoregressive (SAC) Model is regarded as the most rational model in our cross-sectional analysis for estimating the long-term effect of the natural environment on the COVID-19 health outcome. However, to make the calculations easy to understand, we show the basic model, OLS, and explain its shortcomings at first. The following equation is built to analyze the relationships between land cover variables and the county-level health outcomes while controlling for other county-level characteristics, based on the OLS model:

$$CHO_i = \beta_0 + \beta_1 LC_i + \beta_2 CON_i + \varepsilon_i \quad (1)$$

where CHO_i represents the COVID-19 health outcome, either prevalence or mortality, of county i , LC_i represents a vector of land cover data of county i , CON_i represents a vector of control variables of county i as control variables, and ε_i represents the error term. When mortality is taken as the dependent variable, CON_i would contain prevalence of county i . In this model β_0 , β_1 and β_2 are parameters to be estimated.

To examine whether the OLS is a reasonable model, we perform a series of statistical tests. At first, we perform the Breusch-Pagan test to detect heteroscedasticity. According to the significant test results, heteroscedasticity exists in the models. Furthermore, because the COVID-19 is an infectious disease that spreads spatially, the residuals of the OLS might also not be spatially randomly distributed. We utilize the Moran's I test for residual spatial autocorrelation. According to the test results, the significant positive values indicate that spatial autocorrelation exists in the OLS results. In other words, the residuals of the OLS are spatially clustered. In this case, a reasonable spatial model is required (Bivand and Piras, 2015).

Spatial Simultaneous Autoregressive (SAC) Model

Spatial models assume that the spatial correlations between a specific observation and its neighbors. The correlations might exist in the dependent variables, the independent variables, and/or the error terms (Bivand and Piras, 2015). If all these three correlations are significant, the OLS model, **Equation 1**, will change to Manski model, as follows:

$$CHO_i = \rho W_i NECHO_i + \beta_0 + \beta_1 LC_i + \beta_2 CON_i + \beta_3 W_i NELC_i + \beta_4 W_i NECON_i + (\lambda W_i u_i + \varepsilon_i) \quad (2)$$

where \mathbf{W}_i is the spatial weight vector, ρ is the spatial autocorrelation coefficient, \mathbf{NECHO}_i is a vector of the COVID-19 health outcome of the neighboring counties of county i , \mathbf{NELC}_i and \mathbf{NECON}_i are vectors of land cover variables and control variables of the neighboring counties of county i , λ is the error spatial dependence coefficient, u_i is the part of error term with spatial dependence of the neighboring counties of county i , and ϵ_i is the part without spatial dependence. β_3 and β_4 are the parameters to be estimated, similar to β_0 , β_1 and β_2 . \mathbf{W}_i is built as follows:

$$W_{ij} = \frac{1}{NE_i}, \text{ and } W_{ij} \in \mathbf{W}_i \quad (3)$$

where W_{ij} is the spatial weight of the neighboring county j to county i , and NE_i is the number of neighboring counties of county i . Our spatial data are contiguous boundaries of each county. In this situation, if they share one boundary point, they are deemed as the neighbors of each other.

Manski model is too complicated to use. We therefore perform a portmanteau spatial test to judge whether the spatial items are necessary. This test is based on the Lagrange multiplier for the spatially lagged dependence and the robust Lagrange multiplier for the error dependence. The test result shows that both spatially lagged dependence and error dependence are significant, so the items, $\rho\mathbf{W}_i\mathbf{NECHO}_i$ and $\lambda\mathbf{W}_iu_i$, is needed in the regression. If $\beta_3\mathbf{W}_i\mathbf{NELC}_i$ and $\beta_4\mathbf{W}_i\mathbf{NECON}_i$ are put into regression, the number of parameters and Akaike information criterion (AIC) will dramatically increase, eventually leading to overfitting. Thus, the SAC model is employed:

$$CHO_i = \rho\mathbf{W}_i\mathbf{NECHO}_i + \beta_0 + \beta_1\mathbf{LC}_i + \beta_2\mathbf{CON}_i + (\lambda\mathbf{W}_iu_i + \epsilon_i) \quad (4)$$

In the SAC model, the neighboring counties' COVID-19 status affects county i 's, but the county i 's also reversely influences these neighboring counties'. Therefore, the

spatial spillover function is an infinite iterative calculation. However, this function is a monotonically concave function and converges towards a certain value. According to the theory and previous studies, 500 iterations are enough to estimate the parameter correctly (Li and Managi, 2021a; Millo and Piras, 2012). In our calculation, we set the number of iterations 1000. Furthermore, the parameters estimated by the models involving a spatially lagged term contain two parts, direct impacts and indirect impacts. The direct impacts are the coefficients of relationships between the independent and dependent variables of a given county, which can be directly calculated by **Equation 4** without any iteration (Millo and Piras, 2012). The indirect impacts are estimated during the iterations (Lesage and Fischer, 2008; Li and Managi, 2021a). Avoiding too mathematical speaking, we provide a simple instance of the indirect impacts. For example, it is assumed New York county environment is good, and a resident from Kings county is attracted to have a trip to New York county. However, unfortunately, this resident gets infected by COVID-19. After this resident goes back, more people are infected. If the patients infected by that resident come and make the people infected in New York county, these infections in New York county are the indirect impact of the New York county environment. The sums of indirect impacts and direct impacts are the total impacts of the estimated parameters, regarded as the actual coefficients (Lesage and Fischer, 2008).

Panel SAC Model

To examine the short-term impacts of the natural environments on the COVID-19 health outcome, we apply the panel SAC model on the panel data set. The panel model selection process is similar to the cross-sectional model selection. The only difference is that the panel model selection process needs to choose the basic model among pooled

regression model (PRM), fixed effects model (FEM), random effects model (REM). According to the F test for individual effects, FEM is better than PRM, as the test result is significant. The significant result of the Hausman test for the panel model demonstrates that FEM is also more rational than REM. Additionally, people infected by COVID-19 are contagious. Even though having gone to hospitals, they are still able to infect other people in the relatively short term. Thus, the health outcomes are associated with the situation in the previous period. Adding a time-lagged term to the panel models is required. Accordingly, the FEM is taken as the basic model:

$$CHO_{it} = \beta_1 LC_{it} + \beta_2 CON_{it} + \beta_3 CHO_{it-1} + a_i + \varepsilon_{it} \quad (5)$$

where CHO_{it} is the COVID-19 health outcome of county i over period t , LC_{it} is the NDVI indicator of county i over period t , CON_{it} is a vector of control variables of county i over period t , CHO_{it-1} is a vector of the COVID-19 health outcome of county i over period $t - 1$ (if the dependent variable is the mortality, the CHO_{it-1} should encompass both the mortality and the prevalence; otherwise, only prevalence should be included), a_i is the time-invariant variable of county i , and ε_{it} is an error term. β_1 , β_2 and β_3 are the parameters to be estimated. Similar to OLS, FEM also assumes that the variables are spatially independent. However, the locally robust panel Lagrange multiplier tests for spatial dependence show that both spatially lagged dependence and error dependence significantly exist. Hence, the panel SAC model is applied:

$$CHO_{it} = \rho W_i NECHO_{it} + \beta_1 LC_{it} + \beta_2 CON_{it} + \beta_3 CHO_{it-1} + a_i + (\lambda W_i u_{it} + \varepsilon_{it}) \quad (6)$$

where $NECHO_{it}$ is a vector of the COVID-19 health outcome of the neighboring counties of county i over period t , u_{it} is the part of error term with spatial dependence of the neighboring counties of county i over period t , and ε_{it} is the part without spatial dependence.

310

311 *Monetary Value of Natural Land Cover on COVID-19*

312 The monetary value of natural land cover on COVID-19 is estimated to illustrate
313 the value of the natural environment. An increase in natural land cover, such as open
314 water and deciduous forest, is associated with a decrease in the COVID-19 mortality
315 and prevalence. However, there might be a trade-off between environmental benefits
316 and economic costs. In this study, the monetary valuation of an increase in natural land
317 cover on COVID-19 health outcomes can be estimated. Assuming no other variables
318 change, except land cover area per capita and household income, the household income
319 change to offset the variation of land cover area per capita is deemed its monetary value.
320 This method is widely used in the implicit evaluation of environmental goods, taking
321 health evaluation as the dependent variable (Alcock et al., 2014; Li and Managi, 2021b;
322 Tsurumi and Managi, 2017). The monetary value evaluation is therefore provided by
323 the marginal rate of substitution between the household income and the ratios of land
324 cover:

$$\frac{\Delta LC_k \cdot \beta_{1k}}{\Delta Income_k \cdot \beta_{Income}} = 1 \quad (7)$$

325 where ΔLC_k is the change of the k land cover, β_{1k} represents the coefficient of the land
326 cover k , $\Delta Income_k$ is the change of household income to offset the shift in land cover,
327 and β_{Income} represents the coefficient of the median household income in counties.
328 Assuming that the change of land cover is one unit, **Equation 7** is transformed as
329 follows:

$$MV_k = \frac{\beta_{1k}}{\beta_{Income}} \cdot Income \quad (8)$$

where MV_k represents the monetary value of the k land cover, and *Income* is the median household income in counties. It must be noted that the income variable in the regression is a natural logarithm, so the monetary values should be different in each county.

Results

Long-term Relationships between Natural Land Cover and the COVID-19 Health Outcomes

Table 1 demonstrates the result of the SAC model taking mortality as the dependent variable (**Model 1**). The spatially lagged dependence coefficient (ρ) is negative, indicating that a specific county's mortality is negatively correlated with its neighbors'. A high mortality might threaten a region to make people carefully prevent the COVID-19. The spatially error dependence coefficient (λ) is positive, proposing that the ignored variables are positively associated. The pseudo R^2 of the SAC model is 0.598, better than the OLS's, 0.483. Natural land cover, including open water, deciduous forest, evergreen forest, is negatively related to the COVID-19 mortality, whose total impacts are -0.004, -0.006, -0.004, respectively. The county with more natural land cover in the long term has a lower mortality. Long-term living in the natural environment might improve physical health and immune system. However, the developed open space is positively associated with mortality, and its total impact is 0.020. The counties with a high proportion of developed open space usually are rural, where medical systems are relatively weak. A relatively higher proportion of natural land cover could decrease the COVID-19 mortality to some degree.

Additionally, several other variables listed in **Table 1** are significant. Prevalence, unemployment rate, the ratio of adults without a high school diploma, poor health rate, average temperature in summer, average temperature in winter, and average relative humidity in summer are positively correlated with mortality. The positive relationship between prevalence and mortality is reasonable. More COVID-19 patients cause huge pressure on the medical systems, and this eventually leads to more deaths. The meteorological variables mentioned here depict the long-term situation rather than short-term variations. Moreover, transport closing restrictions, staying home restrictions, the proportion of population ages 15 to 44, the ratio of Hispanic people, the ratio of male, median household income, average poor mental health days, adult smoking rate, and the PM_{2.5} concentration are negatively associated with the mortality. The adult smoking rate is absolutely not the reason for the mortality reduction. We are concerned that its impacts are masked by other variables. According to the correlation test between median household income and adult smoking rate, their correlation is strongly negative (-0.661). The total impact of median household income is much greater than the adult smoking rate's. In this way, the result is still acceptable. The situation of PM_{2.5} concentration is similar. The high PM_{2.5} concentration is poisonous, which could not decrease the COVID-19 mortality. There is a significantly strong correlation (0.516) between PM_{2.5} and average relative humidity in summer.

Table 1 might be located here.

Table 2 illustrates the result of the SAC model taking prevalence as the dependent variable (**Model 2**). The spatially lagged dependence coefficient (ρ) is negative, and the spatially error dependence coefficient (λ) is positive. The pseudo R^2 of the SAC model is 0.599, better than the OLS's, 0.441. The relationships between open water, deciduous forest, and mixed forest and the prevalence are negative, whose coefficients are -0.164,

-0.099, -0.278, respectively, tallying with the previous study (You and Pan, 2020). However, another type of natural land cover, emergent herbaceous wetlands, is positively associated with the prevalence, and its total impact is 0.342. We delve into the spatial distribution of this land type. This land type is mainly distributed in Florida, Louisiana, Texas, and Minnesota, severely suffering from high COVID-19 prevalence (**Figure A1**). Additionally, the medium intensity developed area strongly prevents the spread of the COVID-19 since its total impact of the prevalence is -1.364. A rational development intensity could ensure sufficient medical resources in the region and enable residents to connect with the natural environment.

Several other control variables are significant and make sense in the cross-sectional analysis on the relationship between natural land cover and prevalence. Among the significant control variables, the days of transport closing, the ratio of Hispanic people, the ratio of male, obesity rate, physical inactivity rate, the ratio of people who have access to exercise opportunities, the numbers of hospital beds, average temperature in summer, and average relative humidity in summer are positively associated with the prevalence, which might favor the dispersal of the virus. However, the transport closing policies and the number of hospital beds do not increase the prevalence but inhibit the COVID-19 pandemic. Since this analysis is cross-sectional, the transport closing policies is indeed affected by the COVID-19 prevalence. The counties with more hospital beds generally have more population, which is likely to cause community transmission without strict prevention policies. Obesity and physical inactivity could lead to physical health issues and reduce immunity, increasing infection likelihood. High temperature and high humidity might be conducive to keeping the virus alive. It must be mentioned that geographical differences cause variations in temperature and humidity here. Furthermore, staying home policies, the ratios of population ages 15 to

44, 45 to 64, and over 65, the ratio of black people, median household income, and poverty rate are negatively linked with the prevalence. Although the associations of the prevalence with the ratios of population ages 15 to 44, 45 to 64, and over 65 are all negative, the total impact of the ratio of population ages over 65 is the largest, followed by the ratio of population ages 45 to 64. Older people may pay more attention to COVID-19 prevention, because they are more likely to die after the infection (Williamson et al., 2020). The richer counties also have a lower prevalence. The poverty rate is significantly correlated with the median household income (correlation coefficient: 0.846), so its real impact is masked by the median household income.

Table 2 might be located here.

Short-term Relationships between NDVI and the COVID-19 Health Outcomes

Table 3 shows the result of the panel SAC model taking mortality as the dependent variable (**Model 3**). The spatially lagged dependence coefficient is negative, while the spatial error dependence coefficient is positive. The R^2 of the panel SAC model is 0.290, lower than the FEM's (0.425). However, the spatially lagged dependence and error dependence tests point out that the panel SAC model is required. We therefore still use the panel SAC model here as the primary model. The NDVI is negatively related to mortality, whose total impact is -0.003. In other words, if the live green vegetation in the counties increases, the COVID-19 mortality would decrease, and more people could survive, consistent with the previous study (He et al., 2020). The prevalence in the current and previous periods causes more deaths, whose coefficients are 0.007 and 0.003, respectively. The mortality in the previous period is negatively associated with the mortality in the current period, whose coefficient is -0.084, indicating that the public could notice the caveats from the high mortality, and the governments might reallocate

medical resources to prevent a further increase of deaths. The strict restriction is seemingly linked with more deaths. Of course, the higher mortality also leads to more stringent restrictions. The short-term average temperature is negatively correlated with mortality. It must be noted the short-term average temperature here is different from the temperatures used in **Model 1**. The variations of the temperatures used in **Model 1** are mainly caused by geographical and spatial differences of the counties. For example, a county in Florida is generally warmer than a county in North Dakota. Yet, the variation of the temperatures shown here is induced by temporal differences. For instance, in the same county, the average temperature in summer is normally higher than it in winter. Thus, the negative relationship between temperature and mortality could be explained that the COVID-19 patients are more likely to die in the winter. The NTL indicates the prosperity of counties. The NTL should be higher if the counties return to the lifestyles of pre-COVID-19 from the restrictions. If the community transmission exists in those counties, it would be a disaster and the mortality would dramatically increase.

Table 3 might be located here.

Table 4 lists the result of the panel SAC model taking prevalence as the dependent variable (**Model 4**). Different from the abovementioned results, in this result, the spatially lagged dependence coefficient is positive. In the short term, the COVID-19 virus spreads spatially without too much prevention because the short-term prevalence of a specific county is strongly positively correlated with its neighbors'. Moreover, the negative spatially error dependence coefficient indicates a negative association of ignored variables. The R^2 of the panel SAC model is 0.867, much better than the FEM's (0.470). The negative NDVI total impact (-0.076) means the natural environment is associated with fewer patients. The high mortality leads to strict restriction, so their

relationship is positive (0.716). The low temperature reduces the infected people, whose coefficient is and -2.544, aligning with the previous study (Notari, 2021). The lagged prevalence is also negatively linked with the current prevalence. The high prevalence warns people to prevent the disease actively. The busy counties have a high NTL, and their residents have more chance to get infected due to loose prevention.

Table 4 might be located here.

Impacts and Monetary Values of Natural Land Cover

With adequate natural land cover, the mortality is lower. According to **Model 1**, a 1% increase in the ratio of open water in a county is associated with a 0.004-death decrease in the deaths due to the COVID-19 per 1,000 capita in that county, shown in **Table 5**. Moreover, a 1% increase in the ratio of deciduous or evergreen forests is linked with a 0.006- or 0.004-death decrease in the deaths per 1,000 capita, respectively. However, a 1% increase in the ratio of developed open space is related to 0.020-death in deaths per 1,000 capita. Moreover, natural land cover and rational development intensity are also associated with fewer confirmed cases, aligning with the previous study (You and Pan, 2020). A 1% increase in the ratio of open water, medium intensity developed area, deciduous forest, or mixed forest is correlated with a 0.164-, 1.364-, 0.099-, or 0.278-case(s) decrease in the COVID-19 confirmed cases per 1,000 capita, based on **Model 2**. But a 1% increase in the ratio of emergent herbaceous wetlands is associated with a 0.342-case increase. Furthermore, a 1% short-term increase in NDVI, namely greenery, leads to a 0.003-death decrease in the deaths per 1,000 capita over a certain period, according to **Model 3**. A 1% short-term increase in NDVI causes a 0.229-case decrease in the confirmed cases per 1,000 capita.

Table 5 might be located here.

The value of improving natural environments is still challenging to understand by the public without professional knowledge. To make well inform them, the coefficients of natural land cover are converted into monetary values. According to **Model 1**, a 1% increase in the ratio of open water, deciduous forest, or evergreen forest in a county is equivalent to an about 212-USD, 313-USD, or 219-USD increase in household income in that county, respectively, listed in **Table 6**. A 1% increase in the ratio of developed open space is associated with a rough 1045-USD decrease in household income. Based on **Model 2**, a 1% increase in the ratio of open water, medium intensity developed area, deciduous forest, or mixed forest is related to an approximate 382-USD, 3183-USD, 230-USD, or 650-USD, respectively. However, a 1% increase in emergent herbaceous wetlands correlates with a 799-USD decrease in household income.

Table 6 might be located here.

Discussion

Several natural land types are negatively associated with the spread of the COVID-19 and the reduction of deaths due to the COVID-19. Our results show that living with more open water and deciduous forest could prevent the spread of the COVID-19 and decrease deaths in the long term. Moreover, a county with more evergreen forests and mixed forests tends to have fewer confirmed cases or deaths. However, emergent herbaceous wetlands are seemingly positively correlated with the prevalence because of their spatial distribution. The development intensity also affects the county-level prevalence and mortality. To confirm whether these relationships happen to be statistical correlations, we further delve into the relationship between the COVID-19

health outcome and short-term average greenery, namely NDVI. The consequences indicate that a high NDVI in a county leads to lower prevalence and mortality in the short term, verifying that natural environments improve COVID-19 health outcomes. Furthermore, the relatively good status, such as more greenery, high household income, among others, in a county, could prevent the disease, but it might also attract more people, including patients, because their direct and indirect impacts are in the opposite direction. Therefore, effective restriction policies are needed to reduce the COVID-19's impacts, even though the basic conditions in a specific place are better than the average level.

Several recent studies argue that green space may be a critical factor in the COVID-19 pandemic. An increase in urban vegetation is associated with a decrease in cumulative COVID-19 infected cases in the U.S. (You and Pan, 2020). The presence of parks and green spaces encourages physical activity, positively associated with human health (Slater et al., 2020). COVID-19 infection is related to the ecological environment in South Korea (Kim, 2021).

The importance of natural environments in the COVID-19 pandemic is illustrated in our research. According to our analyses, the natural environment may play a practical part in cutting down the COVID-19 prevalence and mortality. In a way, our finding provides evidence for the previous perspectives and studies (Diaz et al., 2018; Hartig and Kahn, 2016; Lachowycz et al., 2012; Slater et al., 2020). Greenspace has effects on physical activity, obesity, mental health, cardiovascular outcomes (James et al., 2015), air pollution (Hoek et al., 2008), and even human well-being (Diener et al., 2018; Li and Managi, 2021b), especially in people living or working in high intensity developed area. These factors are associated with the possibility of several medical conditions, like cardiovascular and respiratory diseases, which may ultimately aggravate the

severity of symptoms after infected COVID-19. Based on these findings, policymakers could make the prevention and control measures more flexible to reduce the negative impacts of those strategies (Yoo and Managi, 2020). An increase in natural land cover may improve public health but negatively affect the economics. To provide more information for the trade-off between health benefit and economic cost, we estimate the monetary of the land cover on health outcome. To sum up, adding more green spaces to living environments should be considered in the future urban planning, achieving several Sustainable Development Goals (SDGs).

An increase of natural land cover in living environments might not directly prevent the spread of COVID-19, but it improves public health status. In other words, with more natural land cover, people may have fewer clinical factors associated with a high risk of death infected by COVID-19 (Williamson et al., 2020). Therefore, these strategies would also prevent outbreaks of other diseases in the future. In this way, an increase in green space and reducing development intensity, at least, help achieve SDG 3 (good health and well-being) and 11 (sustainable cities and communities).

There are some limitations worth noting in this study. Firstly, some potential factors may be overlooked or unable to be obtained, although we have already controlled 28 county-level variables in the cross-sectional analyses. Secondly, the resolution of land cover and the lag of these data increase the uncertainty because the latest land cover data are about 2019 with a resolution of 30 meters (Becker et al., 2019; Wickham et al., 2013). Thirdly, the COVID-19 data are county-level data, possibly resulting in an ecological fallacy. Fourthly, since the models are based on FEM, all variables should be panel data with temporal variations in the panel analyses. So only fewer variables are controlled. Future studies are better to use finer-scale or even individual-level data, to detect the casual interpretation of associations discussed in this article. The physical

mechanisms that natural land cover affects the spread of contiguous diseases, similar to the COVID-19, should be deeply investigated. Additionally, the specific costs and benefits of increasing natural land cover to achieve SDGs need further estimations.

Conclusion

Our results indicate that natural land cover could reduce the COVID-19 prevalence and mortality in both long and short terms. A 1% increase of the ratio of open water, deciduous forest, or evergreen forest is linked with a 0.004-, 0.006-, or 0.004-death decrease in the mortality, equivalent to a 212-, 313-, or 219-USD increase in the household income, respectively. Moreover, in terms of prevalence, a 1% increase of the ratio of open water, deciduous forest, or evergreen forest are worth 382, 230, or 650 USD, respectively. A rational development intensity of residential areas is also an effective approach to cut down deaths and confirmed cases. The relationships between short-term variations of greenery and the COVID-19 health outcomes strength that natural environments could help prevent the spread of COVID-19 and slash mortality. Our research highlights that governments could prevent other pandemics in the future and improve public health by increasing natural land cover, such as open water and deciduous forest.

Data Availability:

All data sources used in the analyses, along with fully reproducible code, are publicly available at https://github.com/MichaelChaoLi-cpu/COVID-19_and_Land_Cover_NDVI.git

Figure




Figure 1(a): The County-level COVID-19 Prevalence (As of Nov. 1st, 2021)




Figure 1(b): The County-level COVID-19 Mortality (As of Nov. 1st, 2021)

Table 1: Result of the SAC Model Taking Mortality as the Dependent Variable (Model 1)

	Direct Impacts	Indirect Impacts	Total Impacts
Open Water (%)	-0.005646** (0.002608)	0.001527** (0.000749)	-0.004119** (0.001891)
Developed Open Space (%)	0.027805** (0.013005)	-0.007519** (0.003622)	0.020286** (0.00955)
Low Intensity Developed Area (%)	-0.011578 (0.014244)	0.003131 (0.003925)	-0.008447 (0.010375)
Medium Intensity Developed Area (%)	-0.019557 (0.022297)	0.005289 (0.006103)	-0.014268 (0.016285)
High Intensity Developed Area (%)	0.037524 (0.026059)	-0.010147 (0.007163)	0.027377 (0.019078)
Deciduous Forest (%)	-0.008322*** (0.002366)	0.00225*** (0.000702)	-0.006071*** (0.001731)
Evergreen Forest (%)	-0.005833** (0.002968)	0.001577** (0.000819)	-0.004256** (0.00218)
Mixed Forest (%)	-0.00109 (0.004576)	0.000295 (0.001245)	-0.000795 (0.003342)
Shrub (%)	0.002565 (0.002802)	-0.000694 (0.000764)	0.001871 (0.00205)
Grassland (%)	0.00028 (0.002454)	-7.6e-05 (0.000665)	0.000205 (0.001795)
Woody Wetlands (%)	0.000195 (0.004057)	-5.3e-05 (0.001101)	0.000142 (0.002966)
Emergent Herbaceous Wetlands (%)	-0.012675 (0.007617)	0.003428 (0.002109)	-0.009247 (0.005571)
Prevalence (cap/1000)	0.013011*** (0.000635)	-0.003519*** (0.000428)	0.009493*** (0.000589)
Gathering Restrictions (days)	-0.000167 (0.000348)	4.5e-05 (9.6e-05)	-0.000122 (0.000254)
Transport Closing (days)	-0.000712*** (0.000193)	0.000193*** (5.8e-05)	-0.00052*** (0.000141)
Staying Home (days)	-0.002503*** (0.000677)	0.000677*** (0.000207)	-0.001826*** (0.000491)
Internal MoRe (days)	-0.00295 (0.006905)	0.000798 (0.001871)	-0.002152 (0.005053)
International MoRe (days)	0.008052 (0.006843)	-0.002178 (0.001864)	0.005875 (0.005017)
Population 15-44 (%)	-0.049464*** (0.010133)	0.013376*** (0.003108)	-0.036088*** (0.007561)
Population 45-64 (%)	-0.004694 (0.013275)	0.001269 (0.003608)	-0.003425 (0.009702)
Population >= 65 (%)	0.005111	-0.001382	0.003729

	(0.009735)	(0.002657)	(0.0071)
Black People (%)	-0.004495	0.001216	-0.003279
	(0.002839)	(0.000796)	(0.002063)
Hispanic People (%)	-0.016814***	0.004547***	-0.012267***
	(0.003767)	(0.001187)	(0.002739)
Male (%)	-0.036678***	0.009919***	-0.02676***
	(0.010031)	(0.002966)	(0.007377)
Unemployment Rate	0.04301**	-0.011631**	0.031379**
	(0.020824)	(0.005788)	(0.015261)
Median Household Income (logarithm)	-1.401255***	0.378932***	-1.022323***
	(0.213768)	(0.073119)	(0.159606)
Poverty Rate (%)	0.012989	-0.003512	0.009476
	(0.00787)	(0.002165)	(0.005765)
Adults Without High School Diploma (%)	0.019924***	-0.005388***	0.014536***
	(0.005513)	(0.001581)	(0.004098)
Poor Health Rate (%)	0.063922***	-0.017286***	0.046636***
	(0.017658)	(0.00532)	(0.012865)
Poor Physical Health (days)	-0.184678	0.049941	-0.134737
	(0.13012)	(0.035925)	(0.095069)
Poor Mental Health (days)	-0.230416*	0.06231*	-0.168106*
	(0.115121)	(0.032536)	(0.083802)
Adult Smoking Rate (%)	-0.039115***	0.010577***	-0.028537***
	(0.014334)	(0.00408)	(0.010518)
Obesity Rate (%)	-0.006043	0.001634	-0.004409
	(0.006535)	(0.001789)	(0.004774)
Physical Inactivity Rate (%)	0.008252	-0.002231	0.00602
	(0.006747)	(0.001842)	(0.004942)
Having Access To Exercise Opportunities (%)	-0.000961	0.00026	-0.000701
	(0.000967)	(0.000265)	(0.000706)
Hospital Beds (bed/1000)	0.004987	-0.001349	0.003638
	(0.003772)	(0.001033)	(0.002762)
Average Temperature In Summer	0.090952***	-0.024596***	0.066357***
	(0.025327)	(0.007782)	(0.018243)
Average Temperature In Winter	0.029361*	-0.00794*	0.021421**
	(0.015092)	(0.004276)	(0.010973)
Average Relative Humidity In Summer	0.019894***	-0.00538**	0.014514***
	(0.007288)	(0.002139)	(0.005277)
Average Relative Humidity In Winter	-0.010229	0.002766	-0.007463
	(0.009529)	(0.002606)	(0.006965)
PM2.5	-0.056104*	0.015172*	-0.040932*
	(0.03291)	(0.009117)	(0.024096)
Spatially lagged dependence coefficient (ρ)	-0.34506***	Log likelihood	-4093.374
Spatially error dependence coefficient (λ)	0.68623***	AIC	8276.7
Number of observations	3103	R ²	0.5984

Note: the standard errors of the estimated parameters are list in the parentheses.

***: $p < 0.01$; **: $p < 0.05$; *: $p < 0.1$

583

584

Table 2: Result of the SAC Model Taking Prevalence as the Dependent Variable (Model 2)

	Direct Impacts	Indirect Impacts	Total Impacts
Open Water (%)	-0.231923*** (0.072038)	0.068225*** (0.022512)	-0.163698*** (0.050807)
Developed Open Space (%)	-0.172223 (0.360558)	0.050663 (0.107107)	-0.12156 (0.254136)
Low Intensity Developed Area (%)	0.480323 (0.383814)	-0.141298 (0.113723)	0.339026 (0.271906)
Medium Intensity Developed Area (%)	-1.931861*** (0.590928)	0.5683*** (0.180937)	-1.363562*** (0.422798)
High Intensity Developed Area (%)	0.8496 (0.696589)	-0.249929 (0.206108)	0.599672 (0.493748)
Deciduous Forest (%)	-0.139624** (0.066787)	0.041074** (0.0202)	-0.098551** (0.047203)
Evergreen Forest (%)	-0.080339 (0.084448)	0.023634 (0.024909)	-0.056706 (0.059795)
Mixed Forest (%)	-0.394239*** (0.133787)	0.115974*** (0.041794)	-0.278265*** (0.094123)
Shrub (%)	-0.110582 (0.087635)	0.03253 (0.026256)	-0.078052 (0.061774)
Grassland (%)	-0.03096 (0.073891)	0.009108 (0.021978)	-0.021853 (0.05206)
Woody Wetlands (%)	-0.025476 (0.122301)	0.007494 (0.0362)	-0.017982 (0.086342)
Emergent Herbaceous Wetlands (%)	0.485201** (0.224188)	-0.142733** (0.066288)	0.342469** (0.160146)
Gathering Restrictions (days)	-0.000967 (0.010591)	0.000285 (0.003107)	-0.000683 (0.007506)
Transport Closing (days)	0.017704*** (0.005734)	-0.005208*** (0.001755)	0.012496*** (0.004088)
Staying Home (days)	-0.127064*** (0.019363)	0.037379*** (0.00683)	-0.089685*** (0.013912)
Internal MoRe (days)	0.157108 (0.190666)	-0.046217 (0.056357)	0.110891 (0.134936)
International MoRe (days)	0.052411 (0.188339)	-0.015418 (0.055714)	0.036993 (0.133019)
Population 15-44 (%)	-1.532032*** (0.282925)	0.450681*** (0.092565)	-1.081351*** (0.20586)
Population 45-64 (%)	-1.875023*** (0.356314)	0.551579*** (0.117443)	-1.323444*** (0.257433)
Population >= 65 (%)	-2.350508*** (0.27225)	0.691454*** (0.099596)	-1.659054*** (0.207722)
Black People (%)	-0.463024*** (0.080368)	0.136209*** (0.027638)	-0.326815*** (0.057384)
Hispanic People (%)	0.198655* (0.080368)	-0.058439* (0.027638)	0.140216* (0.057384)

	(0.103038)	(0.030637)	(0.073239)
Male (%)	3.143065***	-0.924602***	2.218463***
	(0.259911)	(0.112796)	(0.206971)
Unemployment Rate	0.541061	-0.159165	0.381896
	(0.559721)	(0.165544)	(0.396054)
Median Household Income (logarithm)	-31.956257***	9.400638***	-22.555619***
	(6.330123)	(2.037901)	(4.587318)
Poverty Rate (%)	-0.422528*	0.124296*	-0.298232*
	(0.229069)	(0.068364)	(0.162612)
Adults Without High School Diploma (%)	0.128418	-0.037777	0.090641
	(0.14871)	(0.044178)	(0.105023)
Poor Health Rate (%)	0.416833	-0.122621	0.294212
	(0.506967)	(0.150881)	(0.357505)
Poor Physical Health (days)	2.526838	-0.743325	1.783513
	(3.594016)	(1.060744)	(2.542089)
Poor Mental Health (days)	-2.528876	0.743925	-1.784952
	(3.319548)	(0.981407)	(2.34709)
Adult Smoking Rate (%)	0.119984	-0.035296	0.084688
	(0.402139)	(0.118423)	(0.284535)
Obesity Rate (%)	0.363426**	-0.10691**	0.256516**
	(0.180932)	(0.054371)	(0.128235)
Physical Inactivity Rate (%)	0.559892***	-0.164705***	0.395188***
	(0.180067)	(0.055336)	(0.128256)
Having Access To Exercise Opportunities (%)	0.254658***	-0.074913***	0.179745***
	(0.025743)	(0.010531)	(0.019218)
Hospital Beds (bed/1000)	0.514691***	-0.151408***	0.363284***
	(0.103005)	(0.033162)	(0.074523)
Average Temperature In Summer	4.442997***	-1.307006***	3.135992***
	(0.774849)	(0.28612)	(0.528971)
Average Temperature In Winter	-0.593354	0.174548	-0.418806
	(0.485631)	(0.144772)	(0.343102)
Average Relative Humidity In Summer	0.510341**	-0.150128**	0.360213**
	(0.230012)	(0.069217)	(0.163197)
Average Relative Humidity In Winter	-0.350458	0.103095	-0.247363
	(0.294551)	(0.087423)	(0.208419)
PM2.5	1.657234	-0.487512	1.169722
	(1.05481)	(0.316816)	(0.744562)
Spatially lagged dependence coefficient (ρ)	-0.38423***	Log likelihood	-14416.31
Spatially error dependence coefficient (λ)	0.76938***	AIC	28921
Number of observations	3103	R ²	0.5990

Note: the standard errors of the estimated parameters are list in the parentheses.

***: $p < 0.01$; **: $p < 0.05$; *: $p < 0.1$

Table 3: Result of the Panel SAC Model Taking Mortality as the Dependent Variable (Model 3)

	Direct Impacts	Indirect Impacts	Total Impacts
Prevalence (cases/1000)	0.012166*** (0.000291)	-0.00549*** (0.000177)	0.006676*** (0.00017)
Restriction Stringency	-0.000899* (0.000477)	0.000406* (0.000216)	-0.000493* (0.000261)
NDVI (%)	-0.005547*** (0.001112)	0.002503*** (0.000507)	-0.003044*** (0.000608)
Temperature (°C)	-0.028789*** (0.002458)	0.012991*** (0.001142)	-0.015798*** (0.001359)
NTL	0.022852*** (0.003825)	-0.010312*** (0.001748)	0.01254*** (0.002094)
Time Lag of Prevalence (cases/1000)	0.004716*** (0.000319)	-0.002128*** (0.000152)	0.002588*** (0.000176)
Time Lag of Mortality (cases/1000)	-0.153623*** (0.006957)	0.069323*** (0.003522)	-0.084301*** (0.003843)
Spatially lagged dependence coefficient (ρ)	-0.692590***	R^2	0.2895
Spatially error dependence coefficient (λ)	0.8325688***		
Number of observations	18612 (N:3102)		

Note: the standard errors of the estimated parameters are list in the parentheses.

***: $p < 0.01$; **: $p < 0.05$; *: $p < 0.1$

Table 4: Result of the Panel SAC Model Taking Prevalence as the Dependent Variable (Model 4)

	Direct Impacts	Indirect Impacts	Total Impacts
Restriction Stringency	0.074928*** (0.00503)	0.640949*** (0.047759)	0.715876*** (0.052521)
NDVI (%)	-0.023987*** (0.007182)	-0.205187*** (0.061738)	-0.229173*** (0.068894)
Temperature (°C)	-0.266258*** (0.01654)	-2.277629*** (0.160078)	-2.543887*** (0.175634)
NTL	0.268543*** (0.023367)	2.297171*** (0.21224)	2.565714*** (0.234801)
Time Lag of Prevalence (cases/1000)	-0.068076*** (0.004417)	-0.582332*** (0.042491)	-0.650408*** (0.046665)
Spatially lagged dependence coefficient (ρ)	0.8325688***	R^2	0.8673
Spatially error dependence coefficient (λ)	-0.692590***		
Number of observations	18612 (N:3102)		

Note: the standard errors of the estimated parameters are list in the parentheses.

***: $p < 0.01$; **: $p < 0.05$; *: $p < 0.1$

587

588

Table 5: The Impacts of Natural Land Cover Change on the Health Outcomes

Model	Land Cover Variable	Health Outcome	Impacts of Health Outcome	95% Confidence Interval
Model 1	Open Water (%)	Mortality	-0.00412	(-0.00782 - -0.00041)
	Developed Open Space (%)		0.02029	(0.00157 - 0.039)
	Deciduous Forest (%)		-0.00607	(-0.00946 - -0.00268)
	Evergreen Forest (%)		-0.00426	(-0.00853 - 2e-05)
Model 2	Open Water (%)	Prevalence	-0.1637	(-0.26328 - -0.06412)
	Medium Intensity Developed Area (%)		-1.36356	(-2.19225 - -0.53488)
	Deciduous Forest (%)		-0.09855	(-0.19107 - -0.00603)
	Mixed Forest (%)		-0.27826	(-0.46275 - -0.09378)
	Emergent Herbaceous Wetlands (%)		0.34247	(0.02858 - 0.65636)
Model 3	NDVI (%)	Mortality	-0.00304	(-0.00424 - -0.00185)
Model 4	NDVI (%)	Prevalence	-0.22917	(-0.3642 - -0.09414)

589

590

Table 6: The Monetary Values of Natural Land Cover Change on the Health Outcomes

Model	Land Cover Variable	Health Outcome	Monetary Value	95% Confidence Interval
Model 1	Open Water (%)	Mortality	212	(-274 - 698)
	Developed Open Space (%)		-1045	(-1531 - -559)
	Deciduous Forest (%)		313	(-173 - 799)
	Evergreen Forest (%)		219	(-267 - 705)
Model 2	Open Water (%)	Prevalence	382	(-104 - 868)
	Medium Intensity Developed Area (%)		3183	(2697 - 3669)
	Deciduous Forest (%)		230	(-256 - 716)
	Mixed Forest (%)		650	(164 - 1136)
	Emergent Herbaceous Wetlands (%)		-799	(-1286 - -313)

591

592

Reference

- Alcock, I., White, M.P., Lovell, R., Higgins, S.L., Osborne, N.J., Husk, K., Wheeler, B.W., 2015. What accounts for 'England's green and pleasant land'? A panel data analysis of mental health and land cover types in rural England. *Landsc. Urban Plan.* 142, 38-46.
- Alcock, I., White, M.P., Wheeler, B.W., Fleming, L.E., Depledge, M.H., 2014. Longitudinal Effects on Mental Health of Moving to Greener and Less Green Urban Areas. *Environ. Sci. Technol.* 48, 1247-1255.
- Becker, D.A., Browning, M.H.E.M., Kuo, M., Van Den Eeden, S.K., 2019. Is green land cover associated with less health care spending? Promising findings from county-level Medicare spending in the continental United States. *Urban For. Urban Green.* 41, 39-47.
- Bivand, R., Piras, G., 2015. Comparing implementations of estimation methods for spatial econometrics. American Statistical Association.
- Bratman, G.N., Anderson, C.B., Berman, M.G., Cochran, B., de Vries, S., Flanders, J., Folke, C., Frumkin, H., Gross, J.J., Hartig, T., Kahn, P.H., Kuo, M., Lawler, J.J., Levin, P.S., Lindahl, T., Meyer-Lindenberg, A., Mitchell, R., Ouyang, Z.Y., Roe, J., Scarlett, L., Smith, J.R., van den Bosch, M., Wheeler, B.W., White, M.P., Zheng, H., Daily, G.C., 2019. Nature and mental health: An ecosystem service perspective. *Sci. Adv.* 5, eaax0903.
- Chen, H., Guo, J., Wang, C., Luo, F., Yu, X., Zhang, W., Li, J., Zhao, D., Xu, D., Gong, Q., Liao, J., Yang, H., Hou, W., Zhang, Y., 2020. Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *The Lancet* 395, 809-815.
- Combes, A.J., Courau, T., Kuhn, N.F., Hu, K.H., Ray, A., Chen, W.S., Chew, N.W., Cleary, S.J., Kushnoor, D., Reeder, G.C., Shen, A., Tsui, J., Hiam-Galvez, K.J., Muñoz-Sandoval, P., Zhu, W.S., Lee, D.S., Sun, Y., You, R., Magnen, M., Rodriguez, L., Im, K.W., Serwas, N.K., Leligidowicz, A., Zamecnik, C.R., Loudermilk, R.P., Wilson, M.R., Ye, C.J., Fragiadakis, G.K., Looney, M.R., Chan, V., Ward, A., Carrillo, S., Matthay, M., Erle, D.J., Woodruff, P.G., Langelier, C., Kangelaris, K., Hendrickson, C.M., Calfee, C., Rao, A.A., Krummel, M.F., 2021. Global absence and targeting of protective immune states in severe COVID-19. *Nature* 591, 124-130.

Costanza, R., De Groot, R., Sutton, P., Van Der Ploeg, S., Anderson, S.J., Kubiszewski, I., Farber, S., Turner, R.K., 2014. Changes in the global value of ecosystem services. *Global Environmental Change* 26, 152-158.

Diaz, S., Pascual, U., Stenseke, M., Martin-Lopez, B., Watson, R.T., Molnar, Z., Hill, R., Chan, K.M.A., Baste, I.A., Brauman, K.A., Polasky, S., Church, A., Lonsdale, M., Larigauderie, A., Leadley, P.W., van Oudenhoven, A.P.E., van der Plaats, F., Schroter, M., Lavorel, S., Aumeeruddy-Thomas, Y., Bukvareva, E., Davies, K., Demissew, S., Erpul, G., Failler, P., Guerra, C.A., Hewitt, C.L., Keune, H., Lindley, S., Shirayama, Y., 2018. Assessing nature's contributions to people. *Science* 359, 270-272.

Diener, E., Oishi, S., Tay, L., 2018. Advances in subjective well-being research. *Nat. Hum. Behav.* 2, 253-260.

Duan, L., Zhu, G., 2020. Psychological interventions for people affected by the COVID-19 epidemic. *The Lancet Psychiatry* 7, 300-302.

Dzhambov, A.M., Lercher, P., Browning, M.H.E.M., Stoyanov, D., Petrova, N., Novakov, S., Dimitrova, D.D., 2020. Does greenery experienced indoors and outdoors provide an escape and support mental health during the COVID-19 quarantine? *Environ. Res.*, 110420.

Frey, B.S., Luechinger, S., Stutzer, A., 2010. The Life Satisfaction Approach to Environmental Valuation, in: Rausser, G.C., Smith, V.K., Zilberman, D. (Eds.), *Annual Review of Resource Economics*, Vol 2, 2010. Annual Reviews, Palo Alto, pp. 139-160.

Guidotti, E., Ardia, D., 2020. COVID-19 Data Hub. *Journal of Open Source Software* 5, 2376.

Hartig, T., Kahn, P.H., 2016. Living in cities, naturally. *Science* 352, 938-940.

He, H., Lin, X., Yang, Y., Lu, Y., 2020. Association of street greenery and physical activity in older adults: A novel study using pedestrian-centered photographs. *Urban For. Urban Green.* 55, 126789.

Hoek, G., Beelen, R., De Hoogh, K., Vienneau, D., Gulliver, J., Fischer, P., Briggs, D., 2008. A review of land-use regression models to assess spatial variation of outdoor air pollution. *Atmos. Environ.* 42, 7561-7578.

James, P., Banay, R.F., Hart, J.E., Laden, F., 2015. A Review of the Health Benefits of Greenness. *Current Epidemiology Reports* 2, 131-142.

Kim, D., 2021. Exploratory study on the spatial relationship between emerging infectious diseases and urban characteristics: Cases from Korea. *Sustainable Cities and Society* 66, 102672.

Lachowycz, K., Jones, A.P., Page, A.S., Wheeler, B.W., Cooper, A.R., 2012. What can global positioning systems tell us about the contribution of different types of urban greenspace to children's physical activity? *Health Place* 18, 586-594.

Lee, A., Thornley, S., Morris, A.J., Sundborn, G., 2020. Should countries aim for elimination in the covid-19 pandemic? *BMJ*, m3410.

Lesage, J.P., Fischer, M.M., 2008. Spatial Growth Regressions: Model Specification, Estimation and Interpretation. *Spatial Economic Analysis* 3, 275-304.

Li, C., Managi, S., 2021a. Contribution of on-road transportation to PM2.5. *Sci Rep* 11.

Li, C., Managi, S., 2021b. Land cover matters to human well-being. *Sci Rep* 11.

Li, C., Managi, S., 2021c. Spatial Variability of the Relationship between Air Pollution and Well-being. *Sustainable Cities and Society*, 103447.

Maas, J., Verheij, R.A., De Vries, S., Spreeuwenberg, P., Schellevis, F.G., Groenewegen, P.P., 2009. Morbidity is related to a green living environment. *Journal of Epidemiology & Community Health* 63, 967-973.

Millo, G., Piras, G., 2012. splm: Spatial Panel Data Models in R. *J. Stat. Softw.* 47, 1-38.

Mitchell, R., Popham, F., 2008. Effect of exposure to natural environment on health inequalities: an observational population study. *Lancet* 372, 1655-1660.

Mollalo, A., Rivera, K.M., Vahabi, N., 2021. Spatial statistical analysis of pre-existing mortalities of 20 diseases with COVID-19 mortalities in the continental United States. *Sustainable Cities and Society*, 102738.

715
716 Mollalo, A., Rivera, K.M., Vahedi, B., 2020a. Artificial Neural Network
717 Modeling of Novel Coronavirus (COVID-19) Incidence Rates across the
718 Continental United States. *Int. J. Environ. Res. Public Health* 17, 4204.

719
720 Mollalo, A., Vahedi, B., Rivera, K.M., 2020b. GIS-based spatial modeling of
721 COVID-19 incidence rate in the continental United States. *Science of The Total*
722 *Environment* 728, 138884.

723
724 Nakamura, H., Managi, S., 2020. Airport risk of importation and exportation of
725 the COVID-19 pandemic. *Transport Policy* 96, 40-47.

726
727 Nishiga, M., Wang, D.W., Han, Y., Lewis, D.B., Wu, J.C., 2020. COVID-19 and
728 cardiovascular disease: from basic mechanisms to clinical perspectives. *Nature*
729 *Reviews Cardiology* 17, 543-558.

730
731 Notari, A., 2021. Temperature dependence of COVID-19 transmission. *Science*
732 *of The Total Environment* 763, 144390.

733
734 Porta, M., 2014. *A dictionary of epidemiology*. Oxford university press.

735
736 Rahmani, A.M., Mirmahaleh, S.Y.H., 2021. Coronavirus disease (COVID-19)
737 prevention and treatment methods and effective parameters: A systematic
738 literature review. *Sustainable Cities and Society* 64, 102568.

739
740 Richardson, E.A., Pearce, J., Mitchell, R., Kingham, S., 2013. Role of physical
741 activity in the relationship between urban green space and health. *Public Health*
742 127, 318-324.

743
744 Rook, G.A., 2013. Regulation of the immune system by biodiversity from the
745 natural environment: An ecosystem service essential to health. *Proceedings of*
746 *the National Academy of Sciences* 110, 18360-18367.

747
748 Sandifer, P.A., Sutton-Grier, A.E., Ward, B.P., 2015. Exploring connections
749 among nature, biodiversity, ecosystem services, and human health and well-
750 being: Opportunities to enhance health and biodiversity conservation. *Ecosyst.*
751 *Serv.* 12, 1-15.

752
753 Slater, S.J., Christiana, R.W., Gustat, J., 2020. Recommendations for Keeping
754 Parks and Green Space Accessible for Mental and Physical Health During
755 COVID-19 and Other Pandemics. *Preventing Chronic Disease* 17.

- Tsurumi, T., Managi, S., 2017. Monetary Valuations of Life Conditions in a Consistent Framework: The Life Satisfaction Approach. *J. Happiness Stud.* 18, 1275-1303.
- Ugolini, F., Massetti, L., Calaza-Martínez, P., Cariñanos, P., Dobbs, C., Ostoić, S.K., Marin, A.M., Pearlmutter, D., Saaroni, H., Šaulienė, I., Simoneti, M., Verlič, A., Vuletić, D., Sanesi, G., 2020. Effects of the COVID-19 pandemic on the use and perceptions of urban green space: An international exploratory study. *Urban For. Urban Green.* 56, 126888.
- Verity, R., Okell, L.C., Dorigatti, I., Winskill, P., Whittaker, C., Imai, N., Cuomo-Dannenburg, G., Thompson, H., Walker, P.G.T., Fu, H., Dighe, A., Griffin, J.T., Baguelin, M., Bhatia, S., Boonyasiri, A., Cori, A., Cucunubá Z., Fitzjohn, R., Gaythorpe, K., Green, W., Hamlet, A., Hinsley, W., Laydon, D., Ndejati-Gilani, G., Riley, S., Van Elsland, S., Volz, E., Wang, H., Wang, Y., Xi, X., Donnelly, C.A., Ghani, A.C., Ferguson, N.M., 2020. Estimates of the severity of coronavirus disease 2019: a model-based analysis. *The Lancet Infectious Diseases* 20, 669-677.
- Wickham, J.D., Stehman, S.V., Gass, L., Dewitz, J., Fry, J.A., Wade, T.G., 2013. Accuracy assessment of NLCD 2006 land cover and impervious surface. *Remote Sens. Environ.* 130, 294-304.
- Williamson, E.J., Walker, A.J., Bhaskaran, K., Bacon, S., Bates, C., Morton, C.E., Curtis, H.J., Mehrkar, A., Evans, D., Inglesby, P., Cockburn, J., McDonald, H.I., Mackenna, B., Tomlinson, L., Douglas, I.J., Rentsch, C.T., Mathur, R., Wong, A.Y.S., Grieve, R., Harrison, D., Forbes, H., Schultze, A., Croker, R., Parry, J., Hester, F., Harper, S., Perera, R., Evans, S.J.W., Smeeth, L., Goldacre, B., 2020. Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 584, 430-436.
- Yoo, S., Managi, S., 2020. Global mortality benefits of COVID-19 action. *Technological Forecasting and Social Change* 160, 120231.
- You, Y., Pan, S., 2020. Urban Vegetation Slows Down the Spread of Coronavirus Disease (COVID-19) in the United States. *Geophysical Research Letters* 47.

Table A1: Cross-sectional Data Statistic Summary

Statistic	N	Mean	St. Dev.	Min	Max
Mortality Rate(cases/1000)	3,103	2.700	1.341	0.000	10.767
Prevalence (cases/1000)	3,103	149.992	36.592	19.622	543.046
Open Water (%)	3,103	4.795	11.413	0.000	90.899
Developed Open Space (%)	3,103	4.285	3.636	0.119	33.574
Low Intensity Developed Area (%)	3,103	2.438	4.067	0.018	42.512
Medium Intensity Developed Area (%)	3,103	1.129	2.822	0.001	34.421
High Intensity Developed Area (%)	3,103	0.465	1.653	0.00002	37.377
Deciduous Forest (%)	3,103	15.980	19.463	0.000	86.896
Evergreen Forest (%)	3,103	8.729	13.767	0.000	80.504
Mixed Forest (%)	3,103	5.254	7.212	0.000	56.211
Shrub (%)	3,103	8.431	18.226	0.000	97.995
Grassland (%)	3,103	9.385	17.050	0.001	97.665
Woody Wetlands (%)	3,103	5.458	9.039	0.000	67.105
Emergent Herbaceous Wetlands (%)	3,103	1.371	3.403	0.000	55.545
Gathering Restrictions (days)	3,103	372.568	118.105	0	583
Transport Closing (days)	3,103	207.508	195.640	0	578
Staying Home (days)	3,103	418.358	68.703	140	571
Internal MoRe (days)	3,103	559.356	27.756	332	584
International MoRe (days)	3,103	560.193	28.407	332	629
Population 15-44 (%)	3,103	35.881	5.053	15.853	64.848
Population 45-64 (%)	3,103	26.198	2.579	9.645	37.642
Population >= 65 (%)	3,103	19.812	4.751	4.859	58.174
Black People (%)	3,103	9.424	14.504	0.000	86.593
Hispanic People (%)	3,103	9.786	13.914	0.648	96.353
Male (%)	3,103	50.085	2.210	42.992	73.486
Unemployment Rate	3,103	3.960	1.388	0.700	18.300
Median Household Income (logarithm)	3,103	10.841	0.241	10.142	11.852
Poverty Rate (%)	3,103	15.165	6.072	2.600	48.400
Adults Without High School Diploma (%)	3,103	13.437	6.338	1.200	66.300
Poor Health Rate (%)	3,103	17.481	4.679	8.289	40.732
Poor Physical Health (days)	3,103	3.921	0.710	2.324	6.430
Poor Mental Health (days)	3,103	3.936	0.611	2.440	5.964
Adult Smoking Rate (%)	3,103	17.833	3.555	6.735	39.080
Obesity Rate (%)	3,103	32.085	4.578	13.600	49.500
Physical Inactivity Rate (%)	3,103	25.753	5.167	8.400	45.100
Having Access To Exercise Opportunities (%)	3,103	62.774	22.988	0.000	100.000
Hospital Beds (bed/1000)	3,103	3.037	4.487	0.000	99.470
Average Temperature In Summer	3,103	303.129	3.176	290.456	313.873
Average Temperature In Winter	3,103	280.409	6.602	264.694	298.340
Average Relative Humidity In Summer	3,103	88.977	9.697	31.643	99.779
Average Relative Humidity In Winter	3,103	87.484	4.789	58.160	97.673
PM2.5	3,103	9.608	1.966	4.612	14.990

Table 2: Panel Data Statistic Summary

Period	Statistic	N	Mean	St. Dev.	Min	Max
2020 Q1	Mortality Rate(cases/1000)	3,102	0.004	0.021	0	0
	Prevalence (cases/1000)	3,102	0.150	0.474	0.000	10.292
	NDVI (%)	3,102	38.572	16.125	-0.524	78.896
	Temperature (°C)	3,102	4.187	7.010	-13.666	21.746
	NTL Index	3,102	9.551	3.356	0.337	21.705
2020 Q2	Mortality Rate(cases/1000)	3,102	0.174	0.341	0.000	4.131
	Prevalence (cases/1000)	3,102	5.030	7.693	0.000	132.134
	NDVI (%)	3,102	57.576	15.466	13.831	82.701
	Temperature (°C)	3,102	18.212	4.344	2.635	34.910
	NTL Index	3,102	13.893	2.743	4.593	26.558
2020 Q3	Mortality Rate(cases/1000)	3,102	0.275	0.378	0.000	4.950
	Prevalence (cases/1000)	3,102	14.653	10.937	0.000	142.959
	NDVI (%)	3,102	68.887	17.221	12.307	90.035
	Temperature (°C)	3,102	23.317	3.511	13.413	41.152
	NTL Index	3,102	17.887	2.760	5.555	28.704
2020 Q4	Mortality Rate(cases/1000)	3,102	0.756	0.734	0	8
	Prevalence (cases/1000)	3,102	47.908	22.928	0.000	259.528
	NDVI (%)	3,102	48.627	14.988	12.764	81.384
	Temperature (°C)	3,102	9.329	5.141	-5.382	22.934
	NTL Index	3,102	15.326	3.450	6.040	26.162
2021 Q1	Mortality Rate(cases/1000)	3,102	0.725	0.525	0.000	5.384
	Prevalence (cases/1000)	3,102	28.651	11.966	0.000	154.811
	NDVI (%)	3,102	37.627	14.443	6.448	79.555
	Temperature (°C)	3,102	3.499	6.899	-12.106	21.592
	NTL Index	3,102	10.785	3.386	1.326	23.234
2021 Q2	Mortality Rate(cases/1000)	3,102	0.222	0.332	0.000	5.553
	Prevalence (cases/1000)	3,102	8.802	6.087	0.000	96.544
	NDVI (%)	3,102	57.677	15.363	11.526	80.733
	Temperature (°C)	3,102	18.687	3.642	4.388	35.383
	NTL Index	3,102	14.118	2.830	5.425	27.505
2021 Q3	Mortality Rate(cases/1000)	3,102	0.392	0.346	0.000	2.692
	Prevalence (cases/1000)	3,102	35.594	17.387	2.160	113.394
	NDVI (%)	3,102	69.420	16.966	11.639	89.066
	Temperature (°C)	3,102	23.355	3.087	12.340	39.154

	NTL Index	3,102	17.684	2.797	7.145	27.032
799	<hr/>					
800						

Table A3: Data Sources

	Source	Note
Mortality	R Package “COVID19” The U.S. CDC	
Prevalence	R Package “COVID19” The U.S. CDC	
Boundaries of Counties	US Census Bureau https://www2.census.gov/geo/tiger/GENZ2017/shp/	Land cover data are calculated by tool Tabulated Area in ArcGIS Pro 2.5.0
Land Cover Data	Multi-Resolution Land Characteristics (MRLC) consortium https://www.mrlc.gov/	
Population	R Package “COVID19”	
Gathering Restrictions Transport Closing Staying Home	R Package “COVID19” the Oxford COVID-19 Government Response Tracker	
Percentage Of Population 45-64		
Percentage Of Population ≥ 65	U.S. Census Bureau https://www.census.gov/data/tables/time-series/demo/popest/2010s-counties-detail.html	
Percentage Of Black People		
Percentage Of Hispanic People		
Percentage Of Males		
Natural Logarithm Of Median Household Income In 2018		
Unemployment Rate In 2019	United States Department of Agriculture https://www.ers.usda.gov/data-products/county-level-data-sets/download-data/	
Natural Logarithm Of Median House Value		
Poverty Rate In 2018		
Percentage Of The Adults With Less Than High School Diploma		
Adult Smoking Rate In 2019		
Population With Obesity Rate	County Health Rankings & Roadmaps https://www.countyhealthrankings.org/explore-health-rankings/rankings-data-documentation	
Physical Inactivity Rate In 2019		
Having Access To Exercise Opportunities In 2019		

Mean Of Daily Temperature In Summer		
Mean Of Daily Temperature In Winter	Gridmet via Google Earth engine http://www.climatologylab.org/gridmet.html	4km * 4km temperature and relative humidity predictions, summer and winter averaged during 2000-2016
Mean Of Relative Humidity In Summer		
Mean Of Relative Humidity In Winter		
PM2.5	U.S. Environmental Protection Agency	
	NASA	
NDVI	https://lpdaac.usgs.gov/products/mod13a3v006/	MOD13A3 MYD13A3
	https://lpdaac.usgs.gov/products/myd13a3v006/	1-km resolution monthly raster
Temperature	NASA	
	https://lpdaac.usgs.gov/products/mod11c2v006/	MOD11C3 MYD11C3
	https://lpdaac.usgs.gov/products/myd11c2v006/	0.05-arc-degree resolution monthly raster
NTL	NASA	
	https://ladsweb.modaps.eosdis.nasa.gov/missions-and-measurements/products/VNP46A3/	VNP46A3 15-arc-second resolution monthly raster

801

802




Figure A1: The Spatial Distribution of Emergent Herbaceous Wetlands Ratio