

Tutorial for Introductory Analysis of Daily Precipitation Data with hydroTSM

Mauricio Zambrano-Bigiarini*

version 0.8, DD-MMM-2022

1 Installation

Installing the latest stable version (from CRAN):

```
install.packages("hydroTSM")
```

Alternatively, you can also try the under-development version (from Github):

```
if (!require(devtools)) install.packages("devtools")
library(devtools)
install_github("hzambran/hydroTSM")
```

2 Setting up the environment

- Loading the *hydroTSM* library, which contains data and functions used in this analysis.

```
library(hydroTSM)
```

```
## Loading required package: zoo
```

```
##
```

```
## Attaching package: 'zoo'
```

```
## The following objects are masked from 'package:base':
```

```
##
```

```
##      as.Date, as.Date.numeric
```

```
## Loading required package: xts
```

- Loading daily precipitation data at the station San Martino di Castrozza, Trento Province, Italy, with data from 01/Jan/1921 to 31/Dec/1990.

```
data(SanMartinoPPts)
```

- Selecting only a 6-years time slice for the analysis

```
x <- window(SanMartinoPPts, start=as.Date("1985-01-01"))
```

- Monthly values of precipitation

```
( m <- daily2monthly(x, FUN=sum) )
```

*mauricio.zambrano@ufrontera.cl

```
## 1985-01-01 1985-02-01 1985-03-01 1985-04-01 1985-05-01 1985-06-01 1985-07-01
##      141.2      7.0      140.6      72.0      175.6      131.4      85.4
## 1985-08-01 1985-09-01 1985-10-01 1985-11-01 1985-12-01 1986-01-01 1986-02-01
##      159.4      27.2      58.4      101.8      54.8      75.8      131.6
## 1986-03-01 1986-04-01 1986-05-01 1986-06-01 1986-07-01 1986-08-01 1986-09-01
##      59.6      237.8      108.2      144.8      81.2      141.0      69.8
## 1986-10-01 1986-11-01 1986-12-01 1987-01-01 1987-02-01 1987-03-01 1987-04-01
##      38.2      44.4      20.4      46.8      111.0      45.6      98.4
## 1987-05-01 1987-06-01 1987-07-01 1987-08-01 1987-09-01 1987-10-01 1987-11-01
##      212.0      153.8      221.8      175.0      90.6      278.8      164.8
## 1987-12-01 1988-01-01 1988-02-01 1988-03-01 1988-04-01 1988-05-01 1988-06-01
##      29.8      118.0      49.8      22.4      100.6      187.4      193.0
## 1988-07-01 1988-08-01 1988-09-01 1988-10-01 1988-11-01 1988-12-01 1989-01-01
##      120.4      149.2      61.2      136.4      10.0      59.4      0.0
## 1989-02-01 1989-03-01 1989-04-01 1989-05-01 1989-06-01 1989-07-01 1989-08-01
##      152.6      46.2      365.4      77.4      241.6      302.8      114.4
## 1989-09-01 1989-10-01 1989-11-01 1989-12-01 1990-01-01 1990-02-01 1990-03-01
##      65.4      12.8      145.0      110.6      51.6      12.4      65.8
## 1990-04-01 1990-05-01 1990-06-01 1990-07-01 1990-08-01 1990-09-01 1990-10-01
##      127.0      74.4      175.0      143.8      90.8      106.0      153.0
## 1990-11-01 1990-12-01
##      326.6      106.0
```

- Dates of the daily values of 'x'

```
dates <- time(x)
```

- Amount of years in 'x' (needed for computations)

```
( nyears <- yip(from=start(x), to=end(x), out.type="nmbr" ) )
```

```
## [1] 6
```

3 Basic exploratory data analysis (EDA)


1) Summary statistics

```
smry(x)
```

```
##           Index      x
## Min.      1985-01-01  0.0000
## 1st Qu.    1986-07-02  0.0000
## Median     1988-01-01  0.0000
## Mean       1988-01-01  3.7470
## 3rd Qu.    1989-07-01  2.6000
## Max.       1990-12-31 122.0000
## IQR        <NA>      2.6000
## sd         <NA>     10.0428
## cv         <NA>      2.6800
## Skewness   <NA>      5.3512
## Kurtosis   <NA>     39.1619
## NA's       <NA>      0.0000
## n          <NA>    2191.0000
```

- Using the *hydroplot* function, which (by default) plots 9 different graphs: 3 ts plots, 3 boxplots and 3 histograms summarizing 'x'. For this example, only daily and monthly plots are produced, and only data starting on 01-Jan-1987 are plotted.

```
hydroplot(x, var.type="Precipitation", main="at San Martino",
          pfreq = "dm", from="1987-01-01")
```



2) Amount of days with information (not NA) per year

```
dwi(x)
```

```
## 1985 1986 1987 1988 1989 1990
## 365 365 365 366 365 365
```

3) Amount of days with information (not NA) per month per year

```
dwi(x, out.unit="mpy")
```

```
##      Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov Dec
## 1985  31  28  31  30  31  30  31  31  30  31  30  31
## 1986  31  28  31  30  31  30  31  31  30  31  30  31
## 1987  31  28  31  30  31  30  31  31  30  31  30  31
## 1988  31  29  31  30  31  30  31  31  30  31  30  31
## 1989  31  28  31  30  31  30  31  31  30  31  30  31
## 1990  31  28  31  30  31  30  31  31  30  31  30  31
```

4) Plotting the monthly precipitation values for each year, useful for identifying dry/wet months.

```

# Daily zoo to monthly zoo
m <- daily2monthly(x, FUN=sum, na.rm=TRUE)


# Creating a matrix with monthly values per year in each column
M <- matrix(m, ncol=12, byrow=TRUE)
colnames(M) <- month.abb
rownames(M) <- unique(format(time(m), "%Y"))

# Plotting the monthly precipitation values
require(lattice)

## Loading required package: lattice
print(matrixplot(M, ColorRamp="Precipitation",
  main="Monthly precipitation at San Martino st., [mm/month]"))

```

Monthly precipitation at San Martino st., [mm/month]



4 Annual analysis

Annual values of precipitation

```
daily2annual(x, FUN=sum, na.rm=TRUE)
```

```
## 1985-01-01 1986-01-01 1987-01-01 1988-01-01 1989-01-01 1990-01-01
##      1154.8      1152.8      1628.4      1207.8      1634.2      1432.4
```

Average annual precipitation

Obvious way:

```
mean( daily2annual(x, FUN=sum, na.rm=TRUE) )
```

```
## [1] 1368.4
```

Another way (more useful for streamflows, where FUN=mean):

The function *annualfunction* applies FUN twice over x:

(i) firstly, over all the elements of x belonging to the same year, in order to obtain the corresponding annual values, and (ii) secondly, over all the annual values of x previously obtained, in order to obtain a single annual value.

```
annualfunction(x, FUN=sum, na.rm=TRUE) / nyears
```

```
## value
```

```
## 1368.4
```

5 Monthly analysis

Median of the monthly values at station 'x'. Not needed, just for looking at these values in the boxplot.

```
monthlyfunction(m, FUN=median, na.rm=TRUE)
```

```
## Jan Feb Mar Apr May Jun Jul Aug Sep Oct Nov Dec
## 63.7 80.4 52.9 113.8 141.9 164.4 132.1 145.1 67.6 97.4 123.4 57.1
```

Vector with the three-letter abbreviations for the month names

```
cmonth <- format(time(m), "%b")
```


Creating ordered monthly factors

```
months <- factor(cmonth, levels=unique(cmonth), ordered=TRUE)
```

Boxplot of the monthly values

```
boxplot( coredata(m) ~ months, col="lightblue", main="Monthly Precipitation",
        ylab="Precipitation, [mm]", xlab="Month")
```

Monthly Precipitation



6 Seasonal analysis

Average seasonal values of precipitation

```
seasonalfunction(x, FUN=sum, na.rm=TRUE) / nyears
```

```
##      DJF      MAM      JJA      SON
## 213.1333 369.4000 470.8000 315.0667
```

Extracting the seasonal values for each year

```
( DJF <- dm2seasonal(x, season="DJF", FUN=sum) )
```

```
## 1985 1986 1987 1988 1989 1990
## 148.2 262.2 178.2 197.6 212.0 174.6
```

```
( MAM <- dm2seasonal(m, season="MAM", FUN=sum) )
```

```
## 1985 1986 1987 1988 1989 1990
## 388.2 405.6 356.0 310.4 489.0 267.2
```

```
( JJA <- dm2seasonal(m, season="JJA", FUN=sum) )
```


```
## 1985 1986 1987 1988 1989 1990
## 376.2 367.0 550.6 462.6 658.8 409.6
```

```
( SON <- dm2seasonal(m, season="SON", FUN=sum) )
```

```
## 1985 1986 1987 1988 1989 1990
## 187.4 152.4 534.2 207.6 223.2 585.6
```

Plotting the time evolution of the seasonal precipitation values

```
hydroplot(x, pfreq="seasonal", FUN=sum, stype="default")
```



7 Some extreme indices

Common steps for the analysis of this section:

Loading daily precipitation data at the station San Martino di Castrozza, Trento Province, Italy, with data from 01/Jan/1921 to 31/Dec/1990.


```
data(SanMartinoPPts)
```

Selecting only a three-year time slice for the analysis

```
x <- window(SanMartinoPPts, start=as.Date("1988-01-01"))
```

Plotting the selected time series

```
hydroplot(x, ptype="ts", pfreq="o", var.unit="mm")
```



7.1 Heavy precipitation days (R10mm)

Counting and plotting the number of days in the period where precipitation is > 10 [mm]

```
( R10mm <- length( x[x>10] ) )
```

```
## [1] 127
```

7.2 Very wet days (R95p)

- Identifying the wet days (daily precipitation ≥ 1 mm):

```
wet.index <- which(x >= 1)
```

- Computing the 95th percentile of precipitation on wet days (PR_{wn95}):

```
( PRwn95 <- quantile(x[wet.index], probs=0.95, na.rm=TRUE) )
```

```
## 95%
```

```
## 39.75
```

Note 1: this computation was carried out for the three-year time period 1988-1990, not the 30-year period 1961-1990 commonly used.

Note 2: missing values are removed from the computation.

- Identifying the very wet days (daily precipitation $\geq PR_{wn95}$)

```
(very.wet.index <- which(x >= PRwn95))
```

```
## [1] 30 92 234 287 422 423 461 550 551 674 676 719 939 950 998
```

```
## [16] 1058 1061 1075
```

- Computing the total precipitation on the very wet days:


```
( R95p <- sum(x[very.wet.index]) )
```

```
## [1] 1196.4
```

Note 3: this computation was carried out for the three-year time period 1988-1990, not the 30-year period 1961-1990 commonly used


7.3 5-day total precipitation

Computing the 5-day total (accumulated) precipitation

```
x.5max <- rollapply(data=x, width=5, FUN=sum, fill=NA, partial= TRUE,
                    align="center")
```

```
hydroplot(x.5max, ptype="ts+boxplot", pfreq="o", var.unit="mm")
```

```
## [Note: pfreq='o' => ptype has been changed to 'ts']
```



Maximum annual value of 5-day total precipitation

```
(x.5max.annual <- daily2annual(x.5max, FUN=max, na.rm=TRUE))
```

```
## 1988-10-14 1989-07-04 1990-11-25
```

```
##      113.2      170.8      237.2
```

Note 1: for this computation, a moving window centred in the current day is used. If the user wants the 5-day total precipitation accumulated in the 4 days before the current day + the precipitation in the current day, the user have to modify the moving window.

Note 2: For the first two and last two values, the width of the window is adapted to ignore values not within the time series

8 Climograph


Since v0.5-0, `hydroTSM` includes a function to plot a climograph, considering not only precipitation but air temperature data as well.

```
# Loading daily ts of precipitation, maximum and minimum temperature  
data(MaquehueTemuco)
```

```
# extracting individual ts of precipitation, maximum and minimum temperature  
pcp <- MaquehueTemuco[, 1]  
tmx <- MaquehueTemuco[, 2]  
tmn <- MaquehueTemuco[, 3]
```


Plotting a full climograph:

```
m <- climograph(pcp=pcp, tmx=tmx, tmn=tmn, na.rm=TRUE,  
  main="Maquehue Temuco Ad (Chile)", lat=-38.770, lon=-72.637)
```




Plotting a climograph with uncertainty bands around mean values, but with no labels for tmx and tmn:

```
m <- climograph(pcp=pcp, tmx=tmx, tmn=tmn, na.rm=TRUE, tmx.labels=FALSE, tmn.labels=FALSE,
  main="Maquehue Temuco Ad (Chile)", lat=-38.770, lon=-72.637)
```




Plotting a climograph with uncertainty bands around mean values, but with no labels for tmx, tmn and pcp:

```
m <- climograph(pcp=pcp, tmx=tmx, tmn=tmn, na.rm=TRUE,
  pcp.labels=FALSE, tmean.labels=FALSE, tmx.labels=FALSE, tmn.labels=FALSE,
  main="Maquehue Temuco Ad (Chile)", lat=-38.770, lon=-72.637)
```



To better represent the hydrological year in Chile (South America), the following figure will plot a full climograph starting in April (`start.month=4`) instead of January (`start.month=1`):

```
m <- climograph(pcp=pcp, tmx=tmx, tmn=tmn, na.rm=TRUE,
               start.month=4, temp.labels.dx=c(rep(-0.2,4), rep(0.2,6),rep(-0.2,2)),
               main="Maquehue Temuco Ad (Chile)", lat=-38.770, lon=-72.637)
```



9 Software details

This tutorial was built under:

```
## [1] "x86_64-pc-linux-gnu (64-bit)"
## [1] "R version 4.2.0 (2022-04-22)"
## [1] "hydroTSM 0.6-6"
```

10 Version history

- v0.8: XXX 2022
- v0.7: Mar 2020

- v0.6: Aug 2017
- v0.5: May 2013
- v0.4: Aug 2011
- v0.3: Apr 2011
- v0.2: Oct 2010
- v0.1: 30-May-2013

11 Appendix

In order to make easier the use of **hydroTSM** for users not familiar with R, in this section a minimal set of information is provided to guide the user in the R world.

11.1 Editors, GUI

- **GNU/Linux only:** ESS (<https://ess.r-project.org/>)
- **Windows only :** NppToR (<https://sourceforge.net/projects/npptor/>)
- **Multi-platform:** Sublime Text (<https://sublime.weberup.com/>) ; RStudio (<https://www.rstudio.com/>)

11.2 Importing data

- `?read.table`, `?write.table`: allow the user to read/write a file (in table format) and create a data frame from it. Related functions are `?read.csv`, `?write.csv`, `?read.csv2`, `?write.csv2`.
- `?zoo::read.zoo`, `?zoo::write.zoo`: functions for reading and writing time series from/to text files, respectively.
- **R Data Import/Export:** <https://cran.r-project.org/doc/manuals/r-release/R-data.html>
- **foreign** R package: read data stored in several R-external formats (dBase, Minitab, S, SAS, SPSS, Stata, Systat, Weka, ...)
- **readxl** R package: Import MS Excel files into R.
- **some examples:** <https://www.statmethods.net/input/importingdata.html>

11.3 Useful Websites

- **Quick R:** <https://www.statmethods.net/>
- **Time series in R:** <https://cran.r-project.org/web/views/TimeSeries.html>
- **Quick reference for the zoo package:** <https://cran.r-project.org/web/packages/zoo/vignettes/zoo-quickref.pdf>