


## Problem 1: Flipping and shifting


Let  $x[n] = \{\cdots, 0, \underset{\uparrow}{1}, 2, 3, 4, 5, 0, \cdots\}$  and  $y[n] = x[2 - n]$ . There are two ways that we can construct  $y[n]$  from  $x[n]$ .



**Method 1:** Shift  $x[n]$  to the left and then flip it.

Assume  $z[n]$  is created by shifting  $x[n]$  to the left by 2, then

$$z[n] = x[n + 2].$$




$y[n]$  is the flipped version of  $z[n]$ , so

$$y[n] = z[-n].$$

Since  $z[n] = x[n + 2] \Rightarrow z[-n] = x[-n + 2]$ ,


$$y[n] = z[-n] = x[-n + 2] = x[2 - n].$$



**Method 2:** Flip  $x[n]$  and then shift it to the right.

Assume  $z[n]$  is the flipped version of  $x[n]$ , then

$$z[n] = x[-n].$$



$y[n]$  is created by shifting  $z[n]$  to the right by 2 (delaying by 2), so

$$z[n] = z[n - 2].$$

Since  $z[n] = x[-n] \Rightarrow z[n - 2] = x[-(n - 2)]$ ,

$$y[n] = z[n - 2] = x[-(n - 2)] = x[2 - n].$$

**Note:** Given  $y[n] = x[n + k]$ ,

- If  $k < 0$ :  $y[n]$  is a right-shifted (delayed) version of  $x[n]$ .
- If  $k > 0$ :  $y[n]$  is a left-shifted version of  $x[n]$ .

## Problem 2: Linearity, shift-invariance, and causality

(a)  $y[n] = \max(0, x[n])$

- Non-linear: superposition does not hold
  - SI: no operation on indices that makes output shift-variant
  - Causal: no access of future indices
- (b)  $y[n] = y[n-1] + 3^{x[n]}$
- Non-linear: output does not scale with scaled input ( $3^{x[n]}$  term)
  - SI: no operation on indices that makes output shift-variant
  - Causal: no access of future indices
- (c)  $y[n] = x[|n| - n]$
- Linear: scaled input yields scaled output
  - SV: shifted input does not yield shifted output due to access function  $|n| - n$
  - Not causal: access of future indices
- (d)  $y[n] = \sum_{m=-\infty}^{n+2} x[m-2]$
- Linear: simply adding previous inputs
  - SI: no operation on indices that makes output shift-variant
  - Causal: no access to future values / indices

### Problem 3: Stability

- (a)  $y[n] = 3x[n]e^n$   
 If  $|x[n]| < M_0$  then  $y[n]$  is not bounded because  $e^n$  goes to infinity. Hence, the system is not stable.
- (b)  $y[n] = x[n] + x[n-1]$   
 If  $|x[n]| \leq M_0$  then the largest output is

$$x[n] + x[n+1] = M_0 + M_0$$

Therefore,

$$|y[n]| \leq 2M_0 < \infty$$

The system is stable because we have a bound.

- (c)  $y[n] = \sum_{m=-\infty}^n x[m]$   
 If  $|x[m]| < M_0$ , there is no bound on  $y[n]$  because the output is a summed accumulation of previous inputs. Bound cannot be calculated. Hence, the system is not stable.

### Problem 4: Convolution

Consider the finite duration sequences  $x[n] = u[n] - u[n-N]$  and  $h[n] = n(u[n] - u[n-M])$ ,  $M \leq N$ . Find an analytical expression for the sequence  $y[n] = h[n] * x[n]$ .

**Solution:** We can rewrite  $x[n]$  and  $h[n]$  as

$$x[n] = \begin{cases} 1, & 0 \leq n < N \\ 0, & \text{else} \end{cases} = \{\cdots, 0, \underset{\uparrow}{1}, 1, \cdots, 1, 0, \cdots\}$$

$$h[n] = \begin{cases} n, & 0 \leq n < M \\ 0, & \text{else} \end{cases} = \{\cdots, 0, \underset{\uparrow}{0}, 1, 2, \cdots, M-1, 0, \cdots\}$$


By definition,

$$(x * h)[n] = \sum_{k=0}^{M-1} h[n-k]x[k] = \sum_{k=0}^{M-1} h[k]$$

(Here,  $x[n]$  can be seen as a windowing function for  $h[n]$ ).

**Question:** For which values of  $n$  that  $(x * h)[n]$  receives non-zero values?

The convolution response only has non-zero values if there is overlap between  $h[n]$  and sliding  $x[n]$  (or  $x[n]$  and sliding  $h[n]$ ). It is more interesting to look at the beginning and the end of this sequence:



The arrow here shows the end of the window. It is clear that the operation does not stop when  $n = M - 1$  but it continues until  $n = (M - 1) + N - 1$ , where  $M - 1$  is the end of  $h[n]$ ,  $N$  is the “length” of  $x[n]$ , and 1 is from overlap condition.

Consider  $N = 5, M = 4$ , then

$$x[n] = \{\cdots, 0, \underset{\uparrow}{1}, 1, 1, 1, 1, 0, 0, \cdots\}$$

$$h[n] = \{\cdots, 0, \underset{\uparrow}{0}, 1, 2, 3, 0, \cdots\}$$

Unrolling the operation gives us:

$n$	-4	-3	-2	-1	0	1	2	3	4	5	6	7
					↓							
$h[n]$					0	1	2	3				
$x[n]$	1	1	1	1	1							
		1	1	1	1	1						
			1	1	1	1	1					
				1	1	1	1	1				
					1	1	1	1	1			
						1	1	1	1	1		
							1	1	1	1	1	
								1	1	1	1	1

The last non-zero index is  $n = (M - 1) + N - 1 = 7$ . The first non-zero index should usually be 0, however in this case  $(x * h)[0] = 0$  because  $h[0] = 0$ . Therefore,  $(x * h)[n] \neq 0$  for  $n \in [1, 7], n \in \mathbb{Z}$ .