

Class 3: Integrated models and ARIMA

Andrew Parnell
andrew.parnell@ucd.ie



Learning outcomes

- ▶ Understand how differencing works to help make data stationary
- ▶ Know the basics of the ARIMA(p, d, q) framework
- ▶ Understand how to fit an ARIMA(p, d, q) model in a realistic setting

1 / 23

2 / 23

Reminder: stationarity

- ▶ A time series is said to be weakly stationary if:
 - ▶ The mean is stable
 - ▶ The variance is stable
 - ▶ The autocorrelation doesn't depend on where you are in the series

Reminder: ARMA models

- ▶ Combine the autoregressive and the moving average framework into one
- ▶ The general equation for an ARMA(p, q) model is:

$$y_t = \alpha + \sum_{i=1}^p \beta_i y_{t-i} + \sum_{j=1}^q \theta_j \epsilon_{t-j} + \epsilon_t$$

3 / 23

4 / 23

Combining ARMA with the random walk to produce ARIMA

- ▶ There is one other time series model we have already met, that of the random walk:

$$y_t = y_{t-1} + \epsilon_t$$

where $\epsilon_t \sim N(0, \sigma^2)$

- ▶ We could re-write this as:

$$y_t - y_{t-1} = \epsilon_t$$

i.e. the *differences* are random normally-distributed noise

5 / 23

Differencing

- ▶ Differencing is a great way of getting rid of a trend
- ▶ If $y_t \approx y_{t-1} + b$ then there will be an increasing linear slope in the time series
- ▶ Creating $y_t - y_{t-1}$ will remove it and all values will hover around the value b
- ▶ Even when the trend is non-linear differencing might help
- ▶ Differencing twice will remove a quadratic trend for the same reasons
- ▶ You can do even higher levels of differencing but this starts to cause problems
- ▶ The twice differenced series is:

$$(y_t - y_{t-1}) - (y_{t-1} - y_{t-2}) = y_t - 2y_{t-1} + y_{t-2}$$

6 / 23

Idea: combine differencing into the ARMA framework

- ▶ We can combine these ideas into the ARMA framework to produce an ARIMA model (the I stands for integrated, i.e. differenced)
- ▶ An ARIMA model isn't really stationary as the differences are actually removing part of the trend
- ▶ The ARIMA model is written as ARIMA(p, d, q) where p and q are as before and d is the number of differences.

7 / 23

Example: the ARIMA(1,1,1) model

- ▶ If we want to fit an ARIMA(1,1,1) model we first let $z_t = y_t - y_{t-1}$ then fit the model:

$$z_t \sim N(\alpha + \beta z_{t-1} + \theta \epsilon_{t-1}, \sigma^2)$$

- ▶ This is equivalent to an ARMA model on the first differences

8 / 23

Fitting an ARIMA(1, 1, 1) model to the wheat data

- ▶ Recall that the ARMA(2,1) fit wasn't very good to the wheat data
- ▶ Instead try an ARIMA(1, 1, 0) model (i.e. AR(1) on the first differences)

```
wheat = read.csv('../data/wheat.csv')
Arima(wheat$wheat, order = c(1, 1, 0))

## Series: wheat$wheat
## ARIMA(1,1,0)
##
## Coefficients:
##          ar1
##        -0.0529
## s.e.    0.1520
##
## sigma^2 estimated as 10076177:  log likelihood=-492.55
## AIC=989.1   AICc=989.34   BIC=993
```

9 / 23

General format: the ARIMA(p,d,q) model

- ▶ First take the d th difference of the series y_t , and call this z_t .
- ▶ If you want to do this by hand in R you can use the `diff` function, e.g. `diff(y, differences = 2)`
- ▶ Then fit the model:

$$z_t \sim N \left(\alpha + \sum_{i=1}^p \beta_i z_{t-i} + \sum_{j=1}^q \theta_j \epsilon_{t-j}, \sigma^2 \right)$$

10 / 23

Choosing p , d and q

- ▶ It's much harder to have an initial guess at all of p , d and q in one go.
- ▶ We can usually guess at the number of differences d from the time series and ACF plots. If there is a very high degree of autocorrelation it's usually a good idea to try a model with $d=1$ or 2
- ▶ I've never met a model where you needed to difference more than twice. Beware of over-differencing

11 / 23

Revisiting the real-world example

12 / 23

Steps in an ARIMA time series analysis


1. Plot the data and the ACF/PACF
2. Decide if the data look stationary or not. If not, perform a suitable transformation and return to 1. **If the data has a strong trend or there is a high degree of autocorrelation try 1 or 2 differences**
3. Guess at values of p , d , and q for an ARIMA(p , d , q) model
4. Fit the model
5. Try a few models around it by increasing/decreasing p , d and q and checking the AIC (or others)
6. Check the residuals
7. Forecast into the future

13 / 23

A real example: wheat data


- Plot reminder

```
wheat = read.csv('../data/wheat.csv')
plot(wheat$year, wheat$wheat, type = 'l')
```



ACF and PACF

```
par(mfrow = c(1, 2))
acf(wheat$wheat)
pacf(wheat$wheat)
```




- Suggest looking at first differences

15 / 23

Plot of first differences


```
plot(diff(wheat$wheat), type = 'l')
```



16 / 23

ACF/PACF of first differences

```
par(mfrow = c(1, 2))
acf(diff(wheat$wheat))
pacf(diff(wheat$wheat))
```



- Interesting peaks in ACF at lag 2, and PACF at lag 2.

17 / 23

First model

```
Arima(wheat$wheat, order = c(0, 1, 0))
```

```
## Series: wheat$wheat
## ARIMA(0,1,0)
##
## sigma^2 estimated as 9905911: log likelihood=-492.61
## AIC=987.22 AICc=987.3 BIC=989.17
```

- This is just a random walk model. Can also get these from forecast with the function naive

18 / 23

Next models

- Try ARIMA(1, 1, 1), ARIMA(1, 1, 0), ARIMA(0, 1, 1)

```
Arima(wheat$wheat, order = c(1, 1, 1))$aic
```

```
## [1] 985.4555
```

```
Arima(wheat$wheat, order = c(1, 1, 0))$aic
```

```
## [1] 989.0983
```

```
Arima(wheat$wheat, order = c(0, 1, 1))$aic
```

```
## [1] 986.0402
```


- Best one seems to be ARIMA(1, 1, 1). (though BIC suggests others)

19 / 23

Check residuals

- Check the residuals of this model


```
my_model_ARIMA111 = Arima(wheat$wheat, order = c(1, 1, 1))
qqnorm(my_model_ARIMA111$residuals)
qqline(my_model_ARIMA111$residuals)
```



20 / 23

Forecast into the future

```
plot(forecast(my_model_ARIMA111,h=20))
```



21 / 23

Why does the difference not continue into the future?

- ▶ You might have expected the forecasts to continue rising into the future due to the difference
- ▶ The MA part of the model is obviously flat as previously discussed as there are no further errors to correct
- ▶ The AR part of the model reverts back to the estimated mean of the last data point because the β parameter is less than 1 - it dampens out the future predictions and stops them from going crazy

22 / 23

Summary

- ▶ ARIMA models extend the ARMA framework to further add in differencing
- ▶ ARIMA models are no longer stationary as soon as $d > 0$
- ▶ A single difference will remove a linear trends, a second difference quadratic trends
- ▶ Can spot the need for differencing from the time series plot and the ACF
- ▶ Do not over-difference your data!

23 / 23