

Numerical Methods for Elliptic Problems – Exercise Sheet 1

Due date: October 13, 2026, 10:15-11:45, Room SP2-0346

Exercise 1 (Vector identities, integration-by-parts):

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with piecewise smooth boundary, $\vec{b} \in C^1(\bar{\Omega}; \mathbb{R}^d)$ and $u, v \in C^2(\bar{\Omega})$. Let $\vec{n}$ denote the outward unit normal and write $\partial_n u := \nabla u \cdot \vec{n}$. Show the following identities:

- (a) $\operatorname{div}(\vec{b} u) = \vec{b} \cdot \nabla u + u \operatorname{div}(\vec{b})$
- (b) $\int_{\Omega} \vec{b} \cdot \nabla u \, dx = - \int_{\Omega} u \operatorname{div} \vec{b} \, dx + \int_{\partial\Omega} u \vec{b} \cdot \vec{n} \, ds$
- (c) $\int_{\Omega} \nabla u \cdot \nabla v \, dx = - \int_{\Omega} \Delta u v \, dx + \int_{\partial\Omega} \partial_n u v \, ds$

Hint: Use the product rule of differentiation and the Gauß integral theorem.

Exercise 2 (Fundamental lemma 1):

Prove the following assertion: Let $\Omega \subset \mathbb{R}^d$ be an open set and $f \in C(\bar{\Omega})$. Then

$$\int_{\omega} f(x) \, dx = 0 \quad \text{for every bounded measurable set } \omega \subset \Omega \quad \Longleftrightarrow \quad f \equiv 0.$$

Show also that if the assumption $f \in C(\bar{\Omega})$ is replaced by $f \in L^1(\Omega)$, then the equivalence remains valid with $f = 0$ almost everywhere in Ω .

Hint: It suffices to test closed balls contained in Ω .

Exercise 3 (Fundamental lemma, 2nd version):

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain and $f \in C(\bar{\Omega})$. Show that

$$\int_{\Omega} f(x) v(x) \, dx = 0 \quad \text{for every } v \in C(\bar{\Omega}) \text{ with } v|_{\partial\Omega} = 0 \quad \Longleftrightarrow \quad f(x) = 0 \quad \text{for every } x \in \Omega.$$

Show moreover that for $f \in L^2(\Omega)$

$$\int_{\Omega} f v \, dx = 0 \quad \text{for all } v \in C_c^\infty(\Omega) \quad \Leftrightarrow \quad \int_{\Omega} f v \, dx = 0 \quad \text{for all } v \in L^2(\Omega) \quad \Leftrightarrow \quad f = 0 \quad \text{a.e. in } \Omega.$$

Hint: In the continuous case, assume $f(x_0) > 0$ at some $x_0 \in \Omega$ and construct a test function supported in a small ball around x_0 . In the L^2 case, use the density of $C_c^\infty(\Omega)$ in $L^2(\Omega)$.

Exercise 4:

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with Lipschitz boundary, $a \in L^\infty(\Omega)$ satisfy $a(x) \geq a_0 > 0$ almost everywhere for some constant a_0 , and let $f \in L^2(\Omega)$.

(a) Derive the necessary optimality condition (Euler–Lagrange equation) for

$$J(u) := \frac{1}{2} \int_{\Omega} a |\nabla u|^2 dx - \int_{\Omega} f u dx \longrightarrow \min_{u: u|_{\partial\Omega}=0}$$

in the form of a variational problem. Compare with the minimal surface problem.

(b) Use integration by parts and the fundamental lemma to derive the corresponding PDE and state its boundary condition assuming that $a \in C^1(\overline{\Omega})$, $f \in C(\overline{\Omega})$, and $u \in C^2(\overline{\Omega})$.

Exercise 5:

Let $\Omega \subset \mathbb{R}^d$ be a domain and $g = u_g|_{\partial\Omega}$ for some $u_g \in C^2(\overline{\Omega})$.

Show that any classical solution $u \in C^2(\Omega) \cap C(\overline{\Omega})$ of

$$-\Delta u = 0 \quad \text{in } \Omega, \quad u = g \quad \text{on } \partial\Omega$$

can be decomposed as $u = u_g + w$, where w solves

$$-\Delta w = f \quad \text{in } \Omega, \quad w = 0 \quad \text{on } \partial\Omega.$$

Determine the right-hand side f , which depends only on u_g .

Exercise 6:

Let $\Omega \subset \mathbb{R}^d$ be a bounded domain with piecewise smooth boundary. Recall the diffusion–convection–reaction equation

$$-\operatorname{div}(a \nabla u - \vec{b} u) + cu = f \quad \text{in } \Omega.$$

Assume sufficient smoothness for the following classical calculations, in particular $\vec{b} \in C^1(\overline{\Omega}; \mathbb{R}^d)$. Show that for $a \equiv 0$ this can be rewritten as a transport equation

$$\vec{b}' \cdot \nabla u + c' u = f' \quad \text{in } \Omega.$$

Assume $f' \equiv c' \equiv 0$ and $\vec{b}'$ constant. Describe the solutions along connected characteristic segments in Ω .

Extra: Use this description and a sketch to formulate boundary conditions that potentially lead to a well-posed problem.

Exercise 7:

Try to install NGSolve on your system by using pip or by compiling it from source such that the `CppExtensionTest.ipynb` notebook can be run successfully. Installation information can be found in `getting-started/ngsolve-installation.pdf`.